

Date:
01.12.15

1 Date: 1.12.15

Def: let y be a function of x and $y^{(n)}(x) = d^n y / dx^n$ for $n \in \mathbb{N} \setminus \{1, 2, 3, \dots\}$ where F is a given function

$$\begin{aligned} 2xy + x^2 y' &= 1 \\ 2xy + x^2 y' - 1 &= 0 \end{aligned}$$

This is an ODE, where

$$F(x, y, y') = 2xy + x^2 y' - 1$$

2 Date: 1.13.15

Date:
01.13.15

ex: Given f on an interval I find F st $F(x)f'(x)$ for all $x \in I$ Here F is by def, an antiderivative of f on I . we denote F as $\int f(x)dx$. All solns F to (1) are of

the form $\int f(x)dx + c$, where c is an arbitray constant, (1) is an ODE, i.i.,

$$y; = f(x) \iff y' - f(x) = 0$$

the latter has the form

$$f(x, y') = 0$$

A little calc III

Given $F : \mathbb{R}^n \rightarrow \mathbb{R}$, say $\vec{a} = (x_1, \dots, x_n) \in \mathbb{R}^n$, the derivative of F at \vec{x} is

$$= F(x) = (F_{x_1}(\vec{x}), F_{x_2}(\vec{x}), \dots, F_{x_n}(\vec{x})) \in Mat_{1 \times n}(\mathbb{R}),$$

where

$$F_{x_i}(\vec{x}) = \lim_{h \rightarrow 0} \frac{F(x_1, \dots, x_i + h, \dots, x_n)}{h}$$

F_{x_i} is called the partial derivative of F with respect to x_i .

we call $\nabla F = F'$ the gradient of F .

given $r : I \subset \mathbb{R} \rightarrow \mathbb{R}^n$, where I subset \mathbb{R} is an interval in \mathbb{R} , the derivative of r at t is

$$r'(t) = (x'_1(t), x'_2(t), \dots, x'_n(t)),$$

$$r(t) = (x_1(t), x_2(t), \dots, x_n(t)),$$

here $r'(t)$ is a tangent vector to r at $r'(t)$ in \mathbb{R}^n .

3 Date: 01.14.15

Date:
01.14.15

Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$, say $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and $F(x) = (f_1)$ thrm(Chain Rule). if \mathbb{R}^n :

Recall: separable ODE

Date:
01.26.15.

$$y' = g(x)h(y) \text{ or } \frac{dy}{dx} = g(x)h(y)$$

"separate" the variable as

$$\frac{1}{h(y)}y' = g(x) \text{ or } \frac{1}{h(y)}\frac{dy}{dx} = g(x)$$

$$\int \frac{1}{h(y)}\frac{dy}{dx}dx = \int g(x)dx$$

Now, $y = y(x)$ then by
by change of vars,

$$\int \frac{1}{h(y)}dy = \int \frac{1}{h(y)}\frac{dy}{dx}dx = \int g(x)dx$$

in short,

$$\int \frac{1}{h(y)}dy = \int g(x)dx$$

Now, once these integrals are evaluated,
if possible, then the resulting eqn
is one of y on the left and x on the
right, which at least implicitly defines
solns $y(x)$ to $y' = h(y)g(x)$. this
resulting eqn may or may not be possible
to solve for y explicitly in terms of x

Example

$x' = kx$, think $x = x(t)$ this is separable; so,

$$\frac{1}{x}\frac{dx}{dt} = k$$

$$\int \frac{1}{x}\frac{dx}{dt}dt = k \int dt \implies$$

(by change of vars)

$$\int \frac{1}{x}dx = \int \frac{1}{x}\frac{dx}{dt}dt = k \int dt$$

$$\implies \ln|x|kt + C,$$

$$\implies |x| = e^{kt+c} = Ce^{kt}, C = e^c > 0$$

$$\implies x(t) = Ce^{kt}, C \neq 0, \text{ any } k \in \mathbb{R}$$

Example

$$T' = k(A - T), \quad k > 0$$

this is separable; so,

$$\begin{aligned}\frac{1}{A - T} \frac{dT}{dt} &= k \implies \\ \int \frac{1}{A - T} dT &= kt + C \implies \\ -\ln|A - T| &= kt + C \implies \\ \frac{1}{A - T} &= Ce^{kt} \implies \\ A - T &= Ce^{-kt} \implies \\ T &= A - Ce^{-kt}\end{aligned}$$

Aside

substitution (change of vars)

$$\int f(g(x))g'(x)dx = \int f(v)dv$$

$v = g(x)$, or

$$\int f(v)\frac{v}{x}dx = \int f(v)dv$$

Date:
01.27.15.

Ex(p.43) 35

$$x(t) = ce^{kt} \text{ (form } x = kx)$$

C14 has a decay rate constant of

$$k = -0.0001216$$

Notice that

$$x(0) = C$$

so, C is called the initial value.
that notation ' x_0 ' is used for C
i.e., $x_0 = x(0)$. thus,

$$x(t) = x_0 e^{kt}$$

in # 35, $x(t) = x_0/6$.
so,

$$\frac{x_0}{6} = x_0 e^{kt}$$

solve for t. thus,

$$t = \frac{1}{6} \ln(1/6) = \frac{1}{|k|} \ln(6)$$

Torricelli's law

Think of $x = x(t)$ and $h = h(t)$
we want $x(t)$, say in particular,
we want t st $x(t) = 0$, so called
"drain time."
recall # 35, p.18, that "ground speed"
is given by $|v| = \sqrt{2gx}$
from "free-fall" a height x.
in contex,

$$\frac{dh}{dt} = \sqrt{2gx}$$

In "the spout" $V = ah$; so

$$\frac{dV}{dt} = a \frac{dh}{dt} \sqrt{2gx}$$

in the tank

$$\frac{dV}{dt} = -a \sqrt{2gx}$$

let $A(x)$ be the ? cros=sectinal
area of the tank at height x, then

$$V = \int_x^0 A(t) dt$$

by ftc(1),

$$\frac{dV}{dx} = A(x)$$

by the chain rule

$$\frac{dV}{dt} = dv/dx dx/dt = A(x)x'$$

$$a(x) * x' = -a\sqrt{2gx}$$

i.e.

$$x' A(x) = -a\sqrt{2gx}$$

which is a separable ODE. thus

$$\frac{A(x)}{\sqrt{x}} dx = -a\sqrt{2g} \implies$$

(int w.r.t t and Δ vars)

$$\int \frac{A(x)}{\sqrt{x}} dx = -at\sqrt{2g}$$

Date:
01.29.15

4 Date: 01.29.2015

Ex (p.45) # 59

revolve $x^2 = by$ about y-axis
depth is 4ft at noon, $y(0) = 4$

$$a = \pi r^2$$

depth is 1ft at 1pm same day
Recall: $\int y^{-1/2} A(y) dx = -8at$ in ft and s.
thus, by torciullis law,

$$-8\pi r^2 t = \pi b \int y^{1/2} dy \implies$$

$$-8r^2 t = \frac{2b}{3} y^{3/2} + C$$

$$y(0) = 4 \implies C = -16b/3$$

\therefore

$$\frac{2b}{3} y^{3/2} = \frac{16b}{3} - 8r^2 t$$

Now, in 3600s (1hr), $y=1$, i.e., $y(3600) = 1$,

$$\frac{2b}{3} = \frac{16b}{3} - 8r^2(3600)$$

$$r^2 = \frac{14b}{3 * 8 * 3600} = \frac{7b}{12 * 3600} \implies$$

$$r = \frac{1}{60} \sqrt{\frac{7b}{12}}$$

Drain time t_0 is

$$0 = \frac{16b}{3} - 8r^2 t_0 \implies t_0 = \frac{2b}{3r^2}$$

Now, in particuler, if
 $y = 4$, then the radius
of $A(y)$ is 2, i.e., $x = 2$.

Thus,

$$x^2 = by \implies 4 = 4b \implies b = 1 \therefore$$

$$r = \frac{1}{60} \sqrt{\frac{17}{12}} \text{ \& } t_0 = \frac{2}{3r^2}$$

Date:
02.02.15

5 Date: 02.02.2015

Thrm existence - uniqueness thrm, $\exists!$ thrm


If $f : D, \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ and $f_y : D_2 \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$


are cts on $R = (a, b) \times (c, d)$ and $(x_0, y_0) \in R$ and

then there exist an interval I st $x_0 \in I$, $I \subseteq (a, b)$ and
the initial value problem

$$\frac{dy}{dx} = f(x, y) \text{ and } y_0 = y(x_0)$$

has a unique $y = y(x)$ for all $x \in I$.





$$R = (a, b) \times (c, d)$$

$$= \{(x, y) \in \mathbb{R}^2 \mid a < x < b \text{ \& } c < y < d\}$$

The $\exists!$ thrm is a "local" result, local to x_0 , more precisely, it just says that there is a unique soln in I not necessarily outside of I .

Ex(p.29) # 27

$$y' = s\sqrt{y} \text{ \& } y(0) = 0$$

consider


$$y(x) = \begin{cases} 0 & x \leq c \\ (x - c)^2 & x \geq c \end{cases}$$

which is ctn on \mathbb{R}

notice that both parts

$y(x)$ satisfy the initial value problem if


$c \geq 0$ Note:



Notice that

$$f(x, y) = 2\sqrt{y} \text{ \& } f_y(x, y) = \frac{1}{\sqrt{y}}$$

which is cts on $R = \{(x, y) \in \mathbb{R}^2 | y > 0\}$



notice that $(0, 0)$ is not in R , not
 "interior" to R . $\therefore \exists!$ thrm does not apply
 Notice that if $y = f(x)$ is a soln to
 an ODE on a interval I
 then so is $y = f(x - c)$ a soln to

the ODE on $I - c = \{x \in R \mid x + c \in I\}$.

Date:
02.03.15.

ex p.28

15. $y' = \sqrt{x-y}$, $y(2) = 2$, no
 16. $y' = \sqrt{x-y}$, $y(2) = 1$, yes
 here $f(x, y) = \sqrt{x-y}$, which has
 cts iff $x - y \geq 0 \iff x \geq y$
 $f_y(x, y) = -\frac{1}{2\sqrt{x-y}}$, which is cts
 on $\{(x, y) \in \mathbb{R}^2 \mid y < x\}$.

Notes

Def A first order linear ODE
 has the form

$$(1) \ y' + p(x)y = q(x)$$

where p & q are cts on some interval \mathcal{I} .

Notice that if $q(x) = 0$ then (1) is
 separable.

Key observation: the left side of
 (1) resembles the product rule.

this motivates a question: Is there
 a cst $I(x)$, say on the interval \mathcal{I} ,
 st

(1') $y'I(x) + yp(x)I(x) = q(x)I(x)$,
 where the left side of (1') is the derivative
 of a product?

if there is such an $I(x)$
 then
 (2) $I'(x) = p(x)I(x)$.
 put $v = I(x)$, then (2) becomes

$$v' = p(x)v$$

which is separable. thus

$$\begin{aligned} \frac{v'}{v} &= p(x) \implies \\ \int \frac{1}{v} dv &= \int p(x) dx \implies \\ \ln |v| &= \int p(x) dx + C \implies \\ |v| &= e^{\int p(x) dx + C} = Ke^{\int p(x) dx} \end{aligned}$$

where $k > 0 \therefore$

$$v = Ke^{\int p(x) dx}, K \neq 0$$

Def. the integrating factor of

$$y' + p(x)y = q(x) \text{ is}$$

$$I(x) = e^{\int p(x)dx}$$

Finally, from (1')

$$(yI(x))' = q(x)I(x) \implies$$

$$yI(x) = \int q(x)I(x)dx \implies$$

$$\boxed{y = \frac{1}{I(x)} \int q(x)I(x)dx}$$

Date:
02.04.15.

Single Tanking Problem

c_i = concentration coming into the tank (constant)
 r_i = rate of flow into the tank (constant)
 $c_0(t)$ = concentration coming out of the tank
 r_0 = rate of flow out of the tank (constant)
 $x(t)$ = amount of salute in tank at time t
 $V(t)$ volume of tank at time t

Units:

$$\text{Concentration} = \frac{\text{amount of solute}}{\text{unit volume}}$$

$$\text{Rate} = \frac{\text{volume}}{\text{unit time}}$$

$$\text{amount} = (\text{concentration})(\text{rate})(\text{time})$$

Notice that the rate of change fo the volume is constant and
 is $m = r_i - r_0$; where, $V(t) = (r_i - r_0)t + V_0 = mt + V_0$, where $V_0 = V(0)$
 For a small Δt
 $x(t + \Delta t) = x(t) + \text{amount in} - \text{amount out over time } \Delta t$

amount in over $\Delta t = c_i r_i \Delta t$;
 amount out over $\Delta t \approx c_0(t) r_0 \Delta t$.
 thus,

$$\Delta x = x(t + \Delta t) - x(t) \approx (c_i r_i - c_0(t) r_0) \Delta t \implies$$

$$\frac{\Delta x}{\Delta t} \approx c_i r_i - c_0(t) r_0$$

this suggest that

$$\frac{dy}{dx} = c_i r_i - c_0(t) r_0$$

Now

$$c_0(t) = \frac{x(t)}{V(t)} \implies$$

$$\frac{dx}{dt} = c_i r_i - \frac{x(t)}{V(t)} r_0$$

which is a 1st order linear ODE.
 more consiely, put $x' = \frac{dx}{dt}$ and
 $x = x(t)$, then

$$\boxed{x' + \frac{r_0}{V(t)} x = c_i r_i}$$

Recall that $V(t) = mt + v_0$, $m = r_i - r_0$; so,

$$x' + \frac{r_0}{mt + v_0} x = c_i r_i$$

Here

$$p(t) = \frac{r_0}{mt + V_0} \implies \int p(x)dx = \frac{r_0}{m} \ln(mt + V_0) + C$$

where in context, $V(t) > 0$. Choose, of ease, $C = 0$, then

$$I(t) = e^{\int p(t)dt} = (mt + V_0)^{r_0/m}$$

So,

$$\begin{aligned} x'(mt + V_0)^{r_0/m} + r_0(mt + V_0)^{r_0/m-m} &= c_i r_i (mt + V_0)^{r_0/m} \implies \\ (x(mt + V_0)^{r_0/m})' &= c_i r_i + (mt + V_0)^{r_0/m} \implies \\ x(mt + V_0)^{r_0/m} &= c_i r_i \int (mt + V_0)^{r_0/m} dt \end{aligned}$$

If $r_0/m = -1$ then $r_0 = -m = -r_i + r_0$
 $\implies r_i = 0$, which is not of interest in
 context ("mixing"). Thus, if $r_0/m \neq -1$
 then

$$\begin{aligned} \int (mt + V_0)^{r_0/m} dt &= \frac{1}{m} (mt + V_0)^{r_0/m+1} \frac{m}{r_0 + m} + C \\ &= \frac{1}{r_i} (mt + V_0)^{r_i/m} + C \\ &\quad \vdots \end{aligned}$$

$$x(mt + v_0)^{r_0/m} = c_i (mt + V_0)^{r_i/m} + C$$

at $t = 0$

$$\begin{aligned} x_0 V_0^{r_0/m} &= c_i V_0^{r_i/m} + C \implies \\ C &= c_i V_0^{r_i/m} - x_0 V_0^{r_0/m} \\ &= V_0^{r_0/m} (c_i V_0 - x_0) \end{aligned}$$

thus,

$$\begin{aligned} x(mt + V_0)^{r_0/m} &= c_i (mt + V_0)^{r_0/m} + V_0^{r_0/m} (c_i V_0 - x_0) \\ \implies x &= c_i (mt + V_0) + (c_i V_0 - x_0) \left(\frac{V_0}{mt + V_0} \right)^{r_0/m} \end{aligned}$$

$$x = c_i V + (c_i V_0 - x_0) \left(\frac{V_0}{V} \right)^{r_0/(r_i - r_0)}$$

where $x = x(t)$ & $V = (r_i - r_0)t + V_0$

Date:
02.09.15.

6 Date: 02.09.2015

1.6 Substitutions in ODEs
Consider a slope field

$$(1) \frac{dy}{dx} = f(y, x)$$

i.e., a 1st order normal ODE. If

$$\alpha(x, y)$$

appears in (1), then we are compelled
to make the substitution

$$v = \alpha(x, y)$$

("alpha" for auxiliary variable)
By the calc III chain rule

$$\begin{aligned} \frac{dv}{dx} &= \frac{\partial \alpha}{\partial x} \frac{dx}{dx} + \frac{\partial \alpha}{\partial y} \frac{dy}{dx} \\ &= \alpha_x + \alpha_y \frac{dy}{dx} \end{aligned}$$

If $v = \alpha(x, y)$ can be solved
for y in terms of x and v, say

$$y = \beta(x, v)$$

then from (1), we have that

$$\frac{dv}{dx} = \alpha_x + \alpha_y \frac{dy}{dx} = \alpha_x + \alpha_y f(x, y)$$

where,

$$\boxed{\frac{dv}{dx} = \alpha_x + \alpha_y f(x, \beta(x, v))}$$

which is a new ODE with dependent variable v and independent variable x.

6.1 ex

$$\frac{dy}{dx} = f(x, y, ax + by + c)$$

put

$$v = ax + by + c$$

then

$$\frac{dv}{dx} = a + b \frac{dy}{dx}$$

thus,

$$\frac{dv}{dx} = a + b \frac{dy}{dx} = a + bf(x, y, ax + by + c)$$

$$\implies \boxed{\frac{dv}{dx} = a + bf(x, (v - ax - c)/b, v)}$$

where

$$y = \beta(x, v) = \frac{v - ax - c}{b}, b \neq 0$$

6.2 p.74 16

$$y' = \sqrt{x + y + 1}$$

$$v = x + y + 1 \implies y = v - x - 1 \text{ \&}$$

$$\frac{dv}{dx} = 1 + \frac{dy}{dx} \implies$$

$$\frac{dv}{dx} = \sqrt{v} + 1 \text{ (separable)}$$

Def. A first order normal homogenous ODE has the form

$$\frac{dy}{dx} = f(y/x)$$

6.3 Ex

$$y' = \frac{xy}{x^2 + y^2}$$

In general, put

$$v = \alpha(x, y) = y/x \text{ (slope)}$$

so, $y = xv$ implies that

$$f(v) = f(y/x) = \frac{dy}{dx} = v + x \frac{dv}{dx} \implies$$

$$x \frac{dv}{dx} = f(v) - v \text{ (separable)}$$

$$\frac{1}{f(v) - v} \frac{dv}{dx} = \frac{1}{x} \implies$$

$$\int \frac{1}{f(v) - v} dv = \ln |x| + c$$

6.4 Ex (Revisited)

$$y' = \frac{xy}{x^2 + y^2} = \frac{y/x}{1 + (y/x)^2}, v = y/x \implies$$

$$\int \frac{1}{\frac{v}{1+v^2} - v} dv = \ln |x| + c \implies$$

$$-\int \frac{1+v^2}{v^3} dv = \ln |x| + c \implies \dots$$

Date:
02.10.15

7 Date: 02.10.2015

Thrm. If $p(x, y) = \sum a_{i_1 i_2} x^{i_1} y^{i_2}$ and $Q(x, y) = \sum a_{j_1 j_2} x^{j_1} y^{j_2}$ are polynomials over \mathbb{R} , then if there is a $k \in \mathbb{Z}^+$ st for all i_1, i_2, j_1, j_2 ,

$$i_1 + i_2 = d = j_1 + j_2$$

then $p(x, y)y' = Q(x, y)$ is a 1st order linear homogenous ODE.

Proof. notice that

$$\begin{aligned} y' \sum a_{i_1 i_2} x^{i_1} y^{i_2} &= \sum a_{j_1 j_2} x^{j_1} y^{j_2} \implies \\ \frac{1}{x^d} y' \sum a_{i_1 i_2} x^{i_1} y^{i_2} &= \frac{1}{x^d} \sum a_{j_1 j_2} x^{j_1} y^{j_2} \implies \\ y' \sum a_{i_1 i_2} \frac{y^{i_2}}{x^{d-i_1}} &= \sum b_{j_1 j_2} \frac{y^{j_2}}{x^{d-j_1}} \implies \\ y' \sum a_{i_1 i_2} \left(\frac{y}{x}\right)^{i_2} &= \sum b_{j_1 j_2} \left(\frac{y}{x}\right)^{j_2} \implies \\ y' &= \frac{\sum a_{i_1 i_2} \left(\frac{y}{x}\right)^{i_2}}{\sum b_{j_1 j_2} \left(\frac{y}{x}\right)^{j_2}} \end{aligned}$$

which is hom. endproof

Def. (i) $\deg a_{i_1, i_2 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} = \sum_{k=1}^n i_{ki}$
(ii) $\deg (p(x_i))$
where $p(x_1, x_2, \dots, x_n) = \sum a_{i_1 i_2 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}$

7.1 ex p.74 2

$$\begin{aligned} 2xyy' &= x^2 + 2y^2 \implies \\ y' &= \left(\frac{1}{2} \frac{1}{y/x} + 2(y/x)\right), v = y/x \implies \\ y = vx \implies y' &= v + xv' \text{ \& } xv' + v = \frac{1}{2v} + v \implies \\ v' &= \frac{1}{2xv} \text{ (separable)} \implies \\ vv' &= \frac{1}{2x} \implies \\ \int v dv &= \frac{1}{2} \int \frac{1}{x} dx = \frac{1}{2} \ln |x| \implies \\ \frac{v^2}{2} &= \frac{1}{2} \ln |x| + c \implies \\ v^2 &= \ln |x| + c \implies \end{aligned}$$

$$\begin{aligned}
v &= + - \sqrt{\ln|x| + c} \implies \\
\frac{y}{x} &+ - \sqrt{\ln|x| + c} \implies \\
y &= + - x \sqrt{\ln|x| + c}
\end{aligned}$$

Def. A first order (normal) Bernoulli ODE has the

$$y' + yp(x) = y^n q(x)$$

side note(good book) Asmlov PDE

8 Date: 02.11.2015

Recall: Bernoulli ODE

$$y' + yp(x) = y^n q(x)$$

put $v = y^m$ then

$$\begin{aligned}
\frac{dv}{dx} &= my^{m-1} \frac{dy}{dx} \implies \\
my^{m-1} \frac{dy}{dx} + my^m p(x) &= my^{m+n-1} q(x) \implies \\
\frac{dv}{dx} vmp(x) &= my^{m+n-1} q(x)
\end{aligned}$$

want: $m + n - 1 = 0$. this requires
that $m = 1 - n$. $\therefore v = y^{1-n}$ reduces
a Bernoulli ODE to a 1st order linear ODE.

8.1 p.74 25

$$\begin{aligned}
&y^2(xy' + y)(1 + x^4)^{1/2} = x \\
\implies &(xy^2y' + y^3)\sqrt{1 + x^4} = x \\
\implies &xy^2y'\sqrt{1 + x^4} + y^3\sqrt{1 + x^4} = x \\
\implies &y'\sqrt{1 + x^4} + y\frac{\sqrt{1 + x^4}}{x} = y^{-2} \\
\implies &y' + y\frac{1}{x} = y^{-2}\frac{1}{\sqrt{1 + x^4}}
\end{aligned}$$

put $v = y^3$ then

$$\begin{aligned}
\frac{dv}{dx} &= 3y^2 \frac{dy}{dx} \implies \\
3y^2y' + \frac{3y^3}{x} &= \frac{3}{\sqrt{1 + x^4}} \implies \\
v' + \frac{3v}{x} &= \frac{3}{\sqrt{1 + x^4}} \text{ (linear)}
\end{aligned}$$

Here $p(x) = 3/x$; so

$$I(x) = e^{\int p(x)dx} = e^{3 \ln|x|} = x^3$$

Thus,

$$\begin{aligned}
 v'x^3 + v3x^2 &= \frac{3x^3}{\sqrt{1+x^4}} \implies \\
 (vx^3)' &= \frac{3x^3}{\sqrt{1+x^4}} \implies \\
 vx^3 &= 3 \int \frac{x^3}{\sqrt{1+x^4}} dx = \frac{3}{4} \int w^{-1/2} dw \\
 &= \frac{3}{4} \cdot \frac{2}{1} w^{1/2} + c \\
 &= \frac{3}{2} (\sqrt{1+x^4} + c) \\
 w &= 1+x^4 \\
 w' &= 4x^3 \\
 y^3 &= \frac{3}{2} \left(\frac{\sqrt{1+x^4} + c}{x^3} \right) \implies \\
 y &= \left(\frac{3}{2} \left(\frac{\sqrt{1+x^4} + c}{x^3} \right) \right)^{1/3}
 \end{aligned}$$

Exam 1 1. Given an ODE and a soln to it, verify it is a soln, then find a particular soln givin an initial cond.

2. Given a description of an ODE, write down the ODE.
3. Dropping ball from some h, find , ground time and speed.
4. high jump on earth given, find high jump on jupiter.
5. (a) solve ODEs
- (b)

one is separable and the other is linear or bornulli

6. Torricelli problem.