

UNOFFICIAL SOLUTIONS BY TheLongCat

B3: ATOMIC AND LASER PHYSICS

TRINITY TERM 2021

Last updated: 30th May 2025

Disclaimer: due to its unofficial nature, the author does not warrant the accuracy of the presented solutions in any form. However, the author is happy to discuss the typos and errors should one arises.

Turn over as you please – we are NOT under exam conditions here.

1. (DRAFT)

(a) Fine structure Hamiltonian:

$$\begin{aligned}\Delta\hat{H}_{\text{SO}} &= \frac{\hbar^2}{2m_e^2c^2} \underbrace{\left\langle \frac{1}{r^3} \right\rangle}_{\frac{1}{l(l+\frac{1}{2})(l+1)} \left(\frac{Z}{na_0}\right)^3} \cdot \frac{Ze^2}{4\pi\epsilon_0} \cdot \langle \mathbf{s} \cdot \mathbf{l} \rangle \quad \text{for hydrogen} \\ \Rightarrow \Delta E_{\text{SO}} &= \frac{\hbar^2}{2m_e^2c^2} \cdot \frac{e^2}{4\pi\epsilon_0} \cdot \frac{1}{l(l+\frac{1}{2})(l+1)} \left(\frac{1}{na_0}\right)^3 \cdot \frac{1}{2} [j(j+1) - l(l+1) - s(s+1)] \\ &\sim \frac{\hbar^2}{2m_e^2c^2} \cdot \frac{e^2}{4\pi\epsilon_0} \cdot \frac{1}{a_0^3}\end{aligned}$$

Zeeman:

$$\begin{aligned}\Delta\hat{H}_z &= -\boldsymbol{\mu} \cdot \mathbf{B} \\ &= \mu_B \mathbf{L} \cdot \mathbf{B} + g_s \mu_B \mathbf{S} \cdot \mathbf{B} \\ &= g_J \mu_B \mathbf{J} \cdot \mathbf{B} \sim \mu_B B\end{aligned}$$

Equating both energies gives:

$$\begin{aligned}\mu_B B &= \frac{\hbar^2}{2m_e^2c^2} \cdot \underbrace{\frac{e^2}{4\pi\epsilon_0}}_{\frac{\alpha\hbar c}{a_0}} \cdot \frac{1}{a_0^3} \\ &= \frac{(\hbar c)^3}{2m_e^2c^4} \cdot \alpha \cdot \frac{1}{a_0^3} \\ &= \frac{(197.33 \text{ MeV fm})^3}{2(0.511 \text{ MeV})^2} \cdot \frac{1}{137.06} \cdot \frac{1}{(5.292 \times 10^4 \text{ fm})^3} \\ &= 7.24 \times 10^{-10} \text{ MeV} \\ &= 7.24 \times 10^{-4} \text{ eV} = 1.16 \times 10^{-22} \text{ J} \\ \Rightarrow B &= 11.9 \text{ T}\end{aligned}$$

So within 12 T, Zeeman perturbation works on top of spin-orbit interaction.

(b) From above, $\Delta\hat{H}_z = \mu_B [\hat{\mathbf{L}} \cdot \hat{\mathbf{B}} + g_s \hat{\mathbf{S}} \cdot \hat{\mathbf{B}}]$.By Wigner-Eckart theorem, we may substitute \mathbf{L} and \mathbf{S} with \mathbf{J} with constant of proportionality akin to vector projection:

$$\begin{aligned}\Delta\hat{H}_z &= \mu_B \left[\frac{\hat{\mathbf{L}} \cdot \hat{\mathbf{J}}}{\hat{J}^2} \hat{\mathbf{J}} \cdot \mathbf{B} + g_s \frac{\hat{\mathbf{S}} \cdot \hat{\mathbf{J}}}{\hat{J}^2} \hat{\mathbf{J}} \cdot \mathbf{B} \right] \\ &= g_J \mu_B \hat{\mathbf{J}} \cdot \mathbf{B}\end{aligned}$$

with $g_J = \frac{3}{2} + \frac{S(S+1) - L(L+1)}{2J(J+1)}$ approximating $g_s \simeq 2$.

(c)

$$\begin{aligned}V_{\text{ext}} &= \alpha(x^2 + y^2) + \beta z^2 \\ &= \alpha r^2 \cos^2 \theta + \beta r^2 \sin^2 \theta\end{aligned}$$

Energy shift $\Delta E = \langle n | \hat{V}_{\text{ext}} | n \rangle$ for state $|n\rangle$.

For $1s$,

$$\begin{aligned}\Delta E_{1s} &= \int_0^{2\pi} d\phi \int_0^\pi d\theta \int_0^\infty \frac{1}{\pi a_0^3} e^{-\frac{2r}{a_0}} (\alpha r^2 \cos^2 \theta + \beta r^2 \sin^2 \theta) \times r^2 \sin \theta dr \\ &= 2\pi \cdot \frac{1}{\pi a_0^3} \int_0^\pi \alpha \cos^2 \theta + \beta \sin^2 \theta \sin \theta d\theta \int_0^\infty r^4 e^{-\frac{2r}{a_0}} dr\end{aligned}$$

Radial integral:

$$\begin{aligned}F = r^2 &\Rightarrow dF = 2r dr \\ dG = e^{-\frac{2r}{a_0}} dr &\Rightarrow G = -\frac{a_0}{2} e^{-\frac{2r}{a_0}}\end{aligned}$$

$$\rightarrow \int_0^\infty a_0 r e^{-\frac{2r}{a_0}} dr$$

$$\begin{aligned}F = r &\Rightarrow dF = dr \\ dG = e^{-\frac{2r}{a_0}} dr &\Rightarrow G = -\frac{a_0}{2} e^{-\frac{2r}{a_0}}\end{aligned}$$

$$\begin{aligned}&\rightarrow \int_0^\infty \frac{a_0^2}{2} e^{-\frac{2r}{a_0}} dr \\ &= \left[-\frac{a_0^3}{4} e^{-\frac{2r}{a_0}} \right]_0^\infty \\ &= \frac{a_0^3}{4} \cdot \frac{12a_0^2}{4} = \frac{3a_0^5}{4}\end{aligned}$$

θ integral:

$$\begin{aligned}&\int_0^\pi \alpha \cos^2 \theta \sin \theta + \beta (1 - \cos^2 \theta) \sin \theta d\theta \\ &= \int_1^{-1} -\alpha \cos^2 \theta - \beta (1 - \cos^2 \theta) d(\cos \theta) \\ &= \left[-\frac{\alpha \cos^3 \theta}{3} - \beta \left(\cos \theta - \frac{\cos^3 \theta}{3} \right) \right]_1^{-1} \\ &= \frac{2\alpha}{3} + \beta \left(2 - \frac{2}{3} \right) = \frac{2\alpha}{3} + \frac{4\beta}{3}\end{aligned}$$

So:

$$\begin{aligned}\Delta E_{1s} &= \frac{2}{a_0^3} \cdot \left(\frac{2\alpha}{3} + \frac{4\beta}{3} \right) \cdot \frac{3a_0^5}{4} \\ &= (\alpha + 2\beta) a_0^2 \\ &= 0 \quad \text{for } \beta = -\frac{\alpha}{2}\end{aligned}$$

For $2s$,

$$\begin{aligned}\Delta E_{2s} &= \int_0^{2\pi} d\phi \int_0^\pi d\theta \int_0^\infty \frac{1}{32\pi a_0^3} \left(2 - \frac{r}{a_0}\right)^2 e^{-\frac{r}{a_0}} r^2 \sin\theta (\alpha r^2 \cos^2\theta + \beta r^2 \sin^2\theta) dr \\ &= \dots\end{aligned}$$

Repeat for $2p$, $3s$, $3p$, $3d$. When $\beta = -\frac{\alpha}{2}$, shifting is null.

(d) **(TO EXPAND)** $\Delta\hat{H}_z = g_J\mu_B\hat{\mathbf{J}} \cdot \mathbf{B}$

For $\mathbf{B} \parallel \hat{\mathbf{z}}$, $\hat{\mathbf{J}} \cdot \mathbf{B} \rightarrow M_J B$

For $\mathbf{B} \parallel \hat{\mathbf{x}}$, $\hat{\mathbf{J}} \cdot \mathbf{B} \rightarrow \hat{\mathbf{J}}_x \cdot B$ where $\hat{\mathbf{J}}_x = \frac{1}{2} [\hat{\mathbf{J}}_+ + \hat{\mathbf{J}}_-]$

Energy shift $\Delta E = \langle n | \Delta\hat{H}_z | n \rangle$

So $\Delta E = 0$ for $\mathbf{B} \parallel \hat{\mathbf{x}}$?

2. (DRAFT)

- (a) The differential equations are the result of time-dependent perturbation theory of monochromatic light-matter interaction – c_1, c_2 are the amplitudes of each stationary state, $\hbar\Omega = \langle 2 | e\mathcal{E}_0 x | 1 \rangle$ is Rabi frequency which characterises the transition frequency in resonance. Rotating wave approximation is the dropping of term with $\omega + \omega_0$, since $|\omega - \omega_0| \ll (\omega + \omega_0)$ for optical transitions, causing the fast mode to be negligible.

(b)

$$\begin{aligned}
 c_2(t) &= \int_0^t -ie^{-i\delta t} \frac{\Omega}{2} dt \quad \text{since } \tau_p \text{ is very short} \Rightarrow c_1 \simeq 1 \\
 &= -\frac{i\Omega}{2} \left[\frac{e^{-i\delta t}}{-i\delta} \right]_0^t \\
 &= -\frac{i\Omega}{2} \left(\frac{e^{-i\delta t} - 1}{-i\delta} \right) \\
 &= \frac{i\Omega}{\delta} e^{-i\frac{\delta}{2}t} \left(\frac{e^{-i\frac{\delta}{2}t} - e^{i\frac{\delta}{2}t}}{2i} \right) \\
 &= -i \frac{i\Omega}{\delta} e^{-i\frac{\delta}{2}t} \sin \frac{\delta}{2}t \\
 \Rightarrow |c_2(\tau_p)|^2 &= \frac{\Omega^2}{\delta^2} \sin^2 \left(\frac{\delta\tau_p}{2} \right)
 \end{aligned}$$

In absence of radiation, $c_2 \rightarrow c_2 e^{i\delta t}$ since it is stationary (only relative phase matter so choose c_1 to have 0 phase).

After T , $c_2 \rightarrow c_2 e^{i\delta T} = c_2(T)$.

Repeating the same calculation above gives:

$$\begin{aligned}
 c_2(T + \tau_p) &= c_2(T) + \frac{-i\Omega}{\delta} \sin \left(\frac{\delta\tau_p}{2} \right) e^{-i\frac{\delta}{2}t} \\
 &= -\frac{i\Omega}{\delta} \sin \left(\frac{\delta\tau_p}{2} \right) e^{i\frac{\delta}{2}\tau_p} [1 + e^{i\delta T}] \\
 \Rightarrow |c_2(t)|^2 &= \frac{\Omega^2}{\delta^2} \sin^2 \left(\frac{\delta\tau_p}{2} \right) \cos^2 \left(\frac{\delta T}{2} \right)
 \end{aligned}$$

- (c) **(TO EXPAND)** Hyperfine $\Delta E = -g_I \mu_N \mathbf{I} \cdot \mathbf{B}_e$

Assuming the Interval Rule holds, transition energy $A \propto g_I \mu_N$

$$\begin{aligned}
 \Rightarrow \omega_0 &\simeq \frac{g_I \mu_N}{\hbar} \\
 &= 1.92 \times 10^4 \text{ T}^{-1} \text{ s}^{-1}
 \end{aligned}$$