

---

# CS273A



## Lecture 6: Genes Enrichment, Gene Regulation I

MW 1:30-2:50pm in Clark **S361\*** (behind Peet's)

Profs: Serafim Batzoglou & Gill Bejerano



CAs: Karthik Jagadeesh & Johannes Birgmeier

\* Mostly: track on website/piazza

# Announcements




- <http://cs273a.stanford.edu/>
  - Lecture slides, problem sets, etc.
- Course communications via Piazza
  - Auditors please sign up too



- PS1 is out.
- Last Tutorial this Fri. UCSC Browser. Bring your laptop!

# Genome Content

# Genes = coding + “non-coding”



# Genes


---

- Gene production is conceptually simple
  - Contiguous stretches of DNA transcribe (1 to 1) into RNA
  - Some (coding or non-coding) RNAs are further spliced
  - Some (m)RNAs are then translated into protein ( $4^3$  to  $20+1$ )
  - Other (nc)RNA stretches just go off to do their thing as RNA
- The devil is in the details, but by and large – this is it.

(non/coding) Gene finding - classical computational challenge:

1. Obtain experimental data
2. Find features in the data (eg, genetic code, splice sites)
3. Generalize from features (eg, predict genes yet unseen)
4. Link to biochemical machinery (eg, spliceosome)

# Coding and non-coding gene production



The cell is constantly making new proteins and ncRNAs.


These perform their function for a while,

And are then degraded.

Newly made coding and non coding gene products take their place.


The picture within a cell is constantly “refreshing”.

# Cell differentiation



To change its behavior a cell can change the repertoire of genes and ncRNAs it makes.



That is exactly what happens when cells differentiate during development from stem cells to their different final fates.



# Genes usually work in groups

Biochemical pathways, signaling pathways, etc.

Asking about the expression perturbation of groups of genes is both more appealing biologically, and more powerful statistically (you sum perturbations).





# Keyword lists are not enough



## Sheer number of terms too much to remember and sort


- Need standardized, stable, carefully defined terms
- Need to describe different levels of detail
- So...defined terms need to be related in a hierarchy

### Anatomy

#### keywords

Organ system  
Cardiovascular system  
Heart

### Anatomy Hierarchy




## With structured vocabularies/hierarchies


- Parent/child relationships exist between terms
- Increased depth -> Increased resolution
- Can annotate data at appropriate level
- May query at appropriate level



# annotate genes to most specific terms




# General Implementations for Vocabularies




1. Annotate at appropriate level, query at appropriate level
2. Queries for higher level terms include annotations to lower level terms


# Example Research Project (to be revisited)



# Let's first ask what is changing?




# Cluster all genes for differential expression



Experiment (replicates)    Control (replicates)

genes




Most significantly up-regulated genes

Unchanged genes


Most significantly down-regulated genes

# Determine cut-offs, examine individual genes

---



# Ask about whole gene sets





# Simplest way to ask: Hypergeometric

Genes measured

$N = 20,000$

Total genes in set 3

$K = 11$

I've picked the top

$n = 100$


diff. expressed genes.

Of them  $k = 8$





belong to gene set 3.

Under a null of randomly distributed genes, how surprising is it?


$P\text{-value} = \Pr_{\text{hyper}}(k \geq 8 \mid N, K, n)$



# GSEA (Gene Set Enrichment Analysis)



# Another popular approach: DAVID






















**DAVID Bioinformatics Resources 2007**  
National Institute of Allergy and Infectious Diseases (NIAID), NIH

## Functional Annotation Chart

Current Gene List: demolist2  
Current Background: Homo sapiens  
394 DAVID IDs  
[Options](#)

[Rerun Using Options](#) [Create Sublist](#) [Download File](#)


Popular site that apparently uses very old (2009?) GO vocabulary.  
Not rec'd by GO any more...

Sublist	Category	Term	RT	Genes	Count	%	P-Value	Benjamini
<input type="checkbox"/>	GOTERM_MF_ALL	<a href="#">protein binding</a>	RT		150	38.1	2.0E-12	4.9E-9
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">development</a>	RT		80	20.3	3.7E-9	1.3E-5
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">response to stress</a>	RT		54	13.7	6.3E-8	1.1E-4
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">regulation of biological process</a>	RT		125	31.7	4.0E-7	4.5E-4
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">negative regulation of biological process</a>	RT		40	10.2	1.1E-6	9.0E-4
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">inflammatory response</a>	RT		19	4.8	1.2E-6	7.9E-4
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">negative regulation of cellular process</a>	RT		38	9.6	1.2E-6	7.0E-4
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">cell communication</a>	RT		116	29.4	1.4E-6	6.8E-4
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">regulation of cellular process</a>	RT		116	29.4	2.3E-6	9.8E-4
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">negative regulation of physiological process</a>	RT		35	8.9	5.3E-6	2.0E-3
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">signal transduction</a>	RT		106	26.9	8.0E-6	2.7E-3
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">cell-cell signaling</a>	RT		30	7.6	9.7E-6	3.0E-3
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">response to external stimulus</a>	RT		29	7.4	1.3E-5	3.6E-3
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">response to pest, pathogen or parasite</a>	RT		30	7.6	2.4E-5	6.2E-3
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">response to wounding</a>	RT		24	6.1	2.7E-5	6.5E-3
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">regulation of physiological process</a>	RT		110	27.9	2.7E-5	6.1E-3
<input type="checkbox"/>	GOTERM_MF_ALL	<a href="#">transcription factor activity</a>	RT		40	10.2	3.2E-5	4.0E-2
<input type="checkbox"/>	GOTERM_MF_ALL	<a href="#">transcription regulator activity</a>	RT		50	12.7	4.2E-5	3.4E-2
<input type="checkbox"/>	GOTERM_BP_ALL	<a href="#">negative regulation of cellular physiological process</a>	RT		32	8.1	4.2E-5	8.9E-3

Input: list of genes of interest (without expression values).

# Multiple Testing Correction

---




Note that statistically you cannot just run individual tests on 1,000 different gene sets. You have to apply further statistical corrections, to account for the fact that even in 1,000 random experiments a handful may come out good by chance.

(eg experiment = Throw a coin 10 times. Ask if it is biased.  
If you repeat it 1,000 times, you will eventually get an all heads series, from a fair coin. Mustn't deduce that the coin is biased)


# RNA-seq

“Next” (2nd) generation sequencing.



# What will you test?

---



Also note that this is a very general approach to test gene lists. Instead of a microarray experiment you can do RNA-seq. Advantage: RNA-seq measures all genes (up to your ability to correctly reconstruct them). Microarrays only measure the probes you can fit on them. (Some genes, or indeed entire pathways, may be missing from some microarray designs).

# Single gene in situ hybridization

---




**Sall1**




# Spatial-temporal maps generation

AI:  
Robotics,  
Vision






# Cell differentiation




To change its behavior a cell can change the repertoire of coding and non-coding genes it makes.

But how?





# Closing the loop

Some proteins and non coding RNAs go “back” to bind DNA near genes, turning these genes on and off.



# Genes & Gene Regulation

- Gene = genomic substring that encodes HOW to make a protein (or ncRNA).
- Genomic switch = genomic substring that encodes WHEN, WHERE & HOW MUCH of a protein to make.



# Transcription Regulation

---

Conceptually simple:


1. The machine that transcribes (“RNA polymerase”)
2. All kinds of proteins and ncRNAs that bind to DNA and to each other to attract or repel the RNA polymerase (“transcription associated factors”).
3. DNA accessibility – making DNA stretches in/accessible to the RNA polymerase and/or transcription associated factors by un/wrapping them around nucleosomes.

(Distinguish DNA patterns from proteins they interact with)

# RNA Polymerase


---

- Transcription = Copying a segment of DNA into (non/coding) RNA
- Gene transcription starts at the (aptly named) TSS, or gene transcription start site
- Transcription is done by RNA polymerase, a complex of 10-12 subunit proteins.
- There are three types of RNA polymerases in human:
  - RNA pol I synthesizes ribosomal RNAs
  - RNA pol II synthesizes pre-mRNAs and most microRNAs
  - RNA pol III synthesizes tRNAs, rRNA and other ssRNAs




# RNA Polymerase is General Purpose

- RNA Polymerase is the general purpose transcriptional machinery.
- It generally does not recognize gene transcription start sites by itself, and requires interactions with multiple additional proteins.





# Terminology

- Transcription Factors (TF): Proteins that return to the nucleus, bind specific DNA sequences there, and affect transcription.
  - There are 1,200-2,000 TFs in the human genome (out of 20-25,000 genes)
  - Only a subset of TFs may be expressed in a given cell at a given point in time.
- Transcription Factor Binding Sites: 4-20bp stretches of DNA where TFs bind.
  - There are millions of TF binding sites in the human genome.
  - In a cell at a given point in time, a site can be either occupied or unoccupied.



# Terminology


- Promoter: The region of DNA 100-1,000bp immediately “upstream” of the TSS, which encodes binding sites for the general purpose RNA polymerase associated TFs, and at times some context specific sites.
  - There are as many promoters as there are TSS’s in the human genome. Many genes have more than one TSS.
- Enhancer: A region of 100-1,000bp, up to 1Mb or more, upstream or downstream from the TSS that includes binding sites for multiple TFs. When bound by (the right) TFs an enhancer turns on/accelerates transcription.
  - Note how an enhancer (E) very far away in sequence (1D) can in fact get very close to the promoter (P) in space (3D).





# TFBS Position Weight Matrix (PWM)



Sites	Alignment Matrix					Frequency weight Matrix					
ATGGCATG	Pos	A	C	G	T	Pos	A	C	G	T	Con
AGGGTGGG	1	9	0	0	1	1	0.9	0	0	0.1	A
ATCGCATG	2	0	1	2	7	2	0	0.1	0.2	0.7	T
TTGCCACG	3	0	1	7	2	3	0	0.1	0.7	0.2	G
ATGGTATT	4	1	1	8	0	4	0.1	0.1	0.8	0	G
ATTGCACG	5	0	7	1	2	5	0	0.7	0.1	0.2	C
AGGGCGTT	6	8	0	2	0	6	0.8	0	0.2	0	A
ATGACATG	7	0	3	0	7	7	0	0.3	0	0.7	T
ATGGCATG	8	0	0	8	2	8	0	0	0.8	0.2	G
ACTGGATG											



Note the strong independence assumption between positions.  
Holds for most transcription binding profiles in the human genome.



# Enhancers



# One nice hypothetical example

