


Mathematical Induction

Part Two

Let P be some property. The ***principle of mathematical induction*** states that if



Theorem: The sum of the first n powers of two is $2^n - 1$.

Proof: Let $P(n)$ be the statement “the sum of the first n powers of two is $2^n - 1$.” We will prove, by induction, that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

For our base case, we need to show $P(0)$ is true, meaning that the sum of the first zero powers of two is $2^0 - 1$. Since the sum of the first zero powers of two is zero and $2^0 - 1$ is zero as well, we see that $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ holds, meaning that

$$2^0 + 2^1 + \dots + 2^{k-1} = 2^k - 1. \quad (1)$$

We need to show that $P(k + 1)$ holds, meaning that the sum of the first $k + 1$ powers of two is $2^{k+1} - 1$. To see this, notice that

$$\begin{aligned} 2^0 + 2^1 + \dots + 2^{k-1} + 2^k &= (2^0 + 2^1 + \dots + 2^{k-1}) + 2^k \\ &= 2^k - 1 + 2^k && \text{(via (1))} \\ &= 2(2^k) - 1 \\ &= 2^{k+1} - 1. \end{aligned}$$

Therefore, $P(k + 1)$ is true, completing the induction. ■

Induction in Practice

- Typically, a proof by induction will not explicitly state $P(n)$.
- Rather, the proof will describe $P(n)$ implicitly and leave it to the reader to fill in the details.
- Provided that there is sufficient detail to determine
 - what $P(n)$ is,
 - that $P(0)$ is true, and that
 - whenever $P(k)$ is true, $P(k+1)$ is true,the proof is usually valid.

Theorem: The sum of the first n powers of two is $2^n - 1$.

Proof: By induction.

For our base case, we'll prove the theorem is true when $n = 0$. The sum of the first zero powers of two is zero, and $2^0 - 1 = 0$, so the theorem is true in this case.

For the inductive step, assume the theorem holds when $n = k$ for some arbitrary $k \in \mathbb{N}$. Then

$$\begin{aligned} 2^0 + 2^1 + \dots + 2^{k-1} + 2^k &= (2^0 + 2^1 + \dots + 2^{k-1}) + 2^k \\ &= 2^k - 1 + 2^k \\ &= 2(2^k) - 1 \\ &= 2^{k+1} - 1. \end{aligned}$$

So the theorem is true when $n = k+1$, completing the induction. ■

Variations on Induction: **Starting Later**

Induction Starting at 0

- To prove that $P(n)$ is true for all natural numbers greater than or equal to 0:
 - Show that $P(0)$ is true.
 - Show that for any $k \geq 0$, that if $P(k)$ is true, then $P(k+1)$ is true.
 - Conclude $P(n)$ holds for all natural numbers greater than or equal to 0.

Induction Starting at m

- To prove that $P(n)$ is true for all natural numbers greater than or equal to m :
 - Show that $P(m)$ is true.
 - Show that for any $k \geq m$, that if $P(k)$ is true, then $P(k+1)$ is true.
 - Conclude $P(n)$ holds for all natural numbers greater than or equal to m .

n^2 versus 2^n

$$0^2 = 0$$

$$2^0 = 1$$

$$1^2 = 1$$

$$2^1 = 2$$

$$2^2 = 4$$

$$2^2 = 4$$

$$3^2 = 9$$

$$2^3 = 8$$

$$4^2 = 16$$

$$2^4 = 16$$

$$5^2 = 25$$

$$2^5 = 32$$

$$6^2 = 36$$

$$2^6 = 64$$

$$7^2 = 49$$

$$2^7 = 128$$

$$8^2 = 64$$

$$2^8 = 256$$

$$9^2 = 81$$

$$2^9 = 512$$

n^2 versus 2^n

$$0^2 = 0 < 2^0 = 1$$

$$1^2 = 1 < 2^1 = 2$$

$$2^2 = 4 = 2^2 = 4$$

$$3^2 = 9 > 2^3 = 8$$

$$4^2 = 16 = 2^4 = 16$$

$$5^2 = 25 < 2^5 = 32$$

$$6^2 = 36 < 2^6 = 64$$

$$7^2 = 49 < 2^7 = 128$$

$$8^2 = 64 < 2^8 = 256$$

$$9^2 = 81 < 2^9 = 512$$

n^2 versus 2^n

$$0^2 = 0 < 2^0 = 1$$

$$1^2 = 1 < 2^1 = 2$$

$$2^2 = 4 = 2^2 = 4$$

$$3^2 = 9 > 2^3 = 8$$

$$4^2 = 16 = 2^4 = 16$$

$$5^2 = 25 < 2^5 = 32$$

$$6^2 = 36 < 2^6 = 64$$

$$7^2 = 49 < 2^7 = 128$$

$$8^2 = 64 < 2^8 = 256$$

$$9^2 = 81 < 2^9 = 512$$

2^n is much bigger here. Does the trend continue?

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.”

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$(k + 1)^2 = k^2 + 2k + 1$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && (\text{since } 1 < 5 \leq k)\end{aligned}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && (\text{since } 1 < 5 \leq k) \\ &= k^2 + 3k\end{aligned}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && (\text{since } 1 < 5 \leq k) \\ &= k^2 + 3k \\ &< k^2 + k \cdot k && (\text{since } 3 < 5 \leq k)\end{aligned}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\&< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\&= k^2 + 3k \\&< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\&= k^2 + k^2\end{aligned}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\&< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\&= k^2 + 3k \\&< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\&= k^2 + k^2 \\&= 2k^2\end{aligned}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\&< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\&= k^2 + 3k \\&< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\&= k^2 + k^2 \\&= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && (\text{since } 1 < 5 \leq k) \\ &= k^2 + 3k \\ &< k^2 + k \cdot k && (\text{since } 3 < 5 \leq k) \\ &= k^2 + k^2 \\ &= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$. Now, by our inductive hypothesis, we know $k^2 < 2^k$.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\ &= k^2 + 3k \\ &< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\ &= k^2 + k^2 \\ &= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$. Now, by our inductive hypothesis, we know $k^2 < 2^k$. Some further algebra gives us the following:

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\ &= k^2 + 3k \\ &< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\ &= k^2 + k^2 \\ &= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$. Now, by our inductive hypothesis, we know $k^2 < 2^k$. Some further algebra gives us the following:

$$(k + 1)^2 < 2k^2 \qquad \text{(from above)}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement " $n^2 < 2^n$." We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\ &= k^2 + 3k \\ &< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\ &= k^2 + k^2 \\ &= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$. Now, by our inductive hypothesis, we know $k^2 < 2^k$. Some further algebra gives us the following:

$$\begin{aligned}(k + 1)^2 &< 2k^2 && \text{(from above)} \\ &< 2(2^k) && \text{(from the inductive hypothesis)}\end{aligned}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\&< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\&= k^2 + 3k \\&< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\&= k^2 + k^2 \\&= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$. Now, by our inductive hypothesis, we know $k^2 < 2^k$. Some further algebra gives us the following:

$$\begin{aligned}(k + 1)^2 &< 2k^2 && \text{(from above)} \\&< 2(2^k) && \text{(from the inductive hypothesis)} \\&= 2^{k+1}\end{aligned}$$

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\ &= k^2 + 3k \\ &< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\ &= k^2 + k^2 \\ &= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$. Now, by our inductive hypothesis, we know $k^2 < 2^k$. Some further algebra gives us the following:

$$\begin{aligned}(k + 1)^2 &< 2k^2 && \text{(from above)} \\ &< 2(2^k) && \text{(from the inductive hypothesis)} \\ &= 2^{k+1}\end{aligned}$$

So $(k+1)^2 < 2^{k+1}$. Therefore, $P(k+1)$ is true, completing the induction.

Theorem: If n is a natural number and $n \geq 5$, then $n^2 < 2^n$.

Proof: Let $P(n)$ be the statement “ $n^2 < 2^n$.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$ where $n \geq 5$, from which the theorem follows.

As our base case, we prove $P(5)$, that $5^2 < 2^5$. Note that $5^2 = 25$ and that $2^5 = 32$, so $5^2 < 2^5$.

For the inductive step, assume that for some $k \geq 5$ that $P(k)$ is true, meaning that $k^2 < 2^k$. We will prove $P(k+1)$, that $(k+1)^2 < 2^{k+1}$.

First, recall that $(k + 1)^2 = k^2 + 2k + 1$. Starting with this equality and using the fact that $k \geq 5$ (that is, that $5 \leq k$), we derive the following:

$$\begin{aligned}(k + 1)^2 &= k^2 + 2k + 1 \\ &< k^2 + 2k + k && \text{(since } 1 < 5 \leq k\text{)} \\ &= k^2 + 3k \\ &< k^2 + k \cdot k && \text{(since } 3 < 5 \leq k\text{)} \\ &= k^2 + k^2 \\ &= 2k^2\end{aligned}$$

So we see that $(k + 1)^2 < 2k^2$. Now, by our inductive hypothesis, we know $k^2 < 2^k$. Some further algebra gives us the following:

$$\begin{aligned}(k + 1)^2 &< 2k^2 && \text{(from above)} \\ &< 2(2^k) && \text{(from the inductive hypothesis)} \\ &= 2^{k+1}\end{aligned}$$


So $(k+1)^2 < 2^{k+1}$. Therefore, $P(k+1)$ is true, completing the induction. ■

A Note on the Previous Proof


- I chose this proof for a few reasons:
 - It uses induction to prove a result that is false for small n , but true for all sufficiently large n .
 - It doesn't require us to introduce any new terms and definitions.
- That said, I'm not a fan of it for a few reasons:
 - This result is not particularly deep or interesting.
 - There's a lot of algebra.
 - It's not at all obvious how to come up with this line of reasoning.
- **Challenge:** Find an inductive proof with the two “good” qualities, but without the three “bad” qualities.

Variations on Induction: **Bigger Steps**


Subdividing a Square




Subdividing a Square




Subdividing a Square



Subdividing a Square



Subdividing a Square




For what values of n can a square be subdivided into n squares?

1 2 3 4 5 6 7 8 9 10 11 12

1 2 3 4 5 6 7 8 9 10 11 12

1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12



1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

1	2
4	3

1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

1		2
		3
6	5	4

1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

5	6	1
4	7	
3		2

1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

1	8		
2			
3			
4	5	6	7

1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

1	2	3
8	9	4
7	6	5

1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

1	2	3	
8	9		
7		10	4
		6	5


1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

1	10		9
2	11		8
3			
4	5	6	7


1 ~~2~~ ~~3~~ 4 ~~5~~ 6 7 8 9 10 11 12

1	2		3
8	9	10	4
	12	11	
7	6		5


The Key Insight




The Key Insight




The Key Insight




The Key Insight




The Key Insight




The Key Insight




The Key Insight




The Key Insight



The Key Insight



The Key Insight



The Key Insight

- If we can subdivide a square into n squares, we can also subdivide it into $n + 3$ squares.
- Since we can subdivide a bigger square into 6, 7, and 8 squares, we can subdivide a square into n squares for any $n \geq 6$:
 - For multiples of three, start with 6 and keep adding three squares until n is reached.
 - For numbers congruent to one modulo three, start with 7 and keep adding three squares until n is reached.
 - For numbers congruent to two modulo three, start with 8 and keep adding three squares until n is reached.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.”

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.




Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.




As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:



Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:






For the inductive step, assume that for some $k \geq 6$ that $P(k)$ is true and a square can be subdivided into k squares.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:






For the inductive step, assume that for some $k \geq 6$ that $P(k)$ is true and a square can be subdivided into k squares. We prove $P(k+3)$, that a square can be subdivided into $k+3$ squares.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:






For the inductive step, assume that for some $k \geq 6$ that $P(k)$ is true and a square can be subdivided into k squares. We prove $P(k+3)$, that a square can be subdivided into $k+3$ squares. To see this, start by obtaining (via the inductive hypothesis) a subdivision of a square into k squares.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:






For the inductive step, assume that for some $k \geq 6$ that $P(k)$ is true and a square can be subdivided into k squares. We prove $P(k+3)$, that a square can be subdivided into $k+3$ squares. To see this, start by obtaining (via the inductive hypothesis) a subdivision of a square into k squares. Then, choose any of the squares and split it into four equal squares.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:






For the inductive step, assume that for some $k \geq 6$ that $P(k)$ is true and a square can be subdivided into k squares. We prove $P(k+3)$, that a square can be subdivided into $k+3$ squares. To see this, start by obtaining (via the inductive hypothesis) a subdivision of a square into k squares. Then, choose any of the squares and split it into four equal squares. This removes one of the k squares and adds four more, so there will be a net total of $k+3$ squares.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.

As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:






For the inductive step, assume that for some $k \geq 6$ that $P(k)$ is true and a square can be subdivided into k squares. We prove $P(k+3)$, that a square can be subdivided into $k+3$ squares. To see this, start by obtaining (via the inductive hypothesis) a subdivision of a square into k squares. Then, choose any of the squares and split it into four equal squares. This removes one of the k squares and adds four more, so there will be a net total of $k+3$ squares. Thus $P(k+3)$ holds, completing the induction.

Theorem: For any $n \geq 6$, it is possible to subdivide a square into n smaller squares.

Proof: Let $P(n)$ be the statement “a square can be subdivided into n smaller squares.” We will prove by induction that $P(n)$ holds for all $n \geq 6$, from which the theorem follows.


As our base cases, we prove $P(6)$, $P(7)$, and $P(8)$, that a square can be subdivided into 6, 7, and 8 squares. This is shown here:



For the inductive step, assume that for some $k \geq 6$ that $P(k)$ is true and a square can be subdivided into k squares. We prove $P(k+3)$, that a square can be subdivided into $k+3$ squares. To see this, start by obtaining (via the inductive hypothesis) a subdivision of a square into k squares. Then, choose any of the squares and split it into four equal squares. This removes one of the k squares and adds four more, so there will be a net total of $k+3$ squares. Thus $P(k+3)$ holds, completing the induction. ■


Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:




Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:



Why This Works

- This induction has three consecutive base cases and takes steps of size three.
- Thinking back to our “induction machine” analogy:



Generalizing Induction

- When doing a proof by induction:
 - Feel free to use multiple base cases.
 - Feel free to take steps of sizes other than one.
- Just be careful to make sure you cover all the numbers you think that you're covering!

Time-Out for Announcements!

Problem Set One

- Problem Set 1 due Friday, October 3 at the start of class.
 - Stop by office hours with questions!
 - Ask questions on Piazza!
 - Email the staff list
(cs103-aut1415-staff@lists.stanford.edu)
with questions!

Checkpoint Problems

- Problem Set 1 Checkpoints graded; feedback available online at Scoryst.
 - Please review this feedback before submitting the rest of the problems – the point of the checkpoint is to get useful feedback!
- In the future, please only submit one copy of the checkpoint per group – it *dramatically* simplifies grading.
- Submitted in the wrong category! No worries! We've got it handled.

A Quick Apology

Solution Sets

- Solution sets for the discussion problems and for the checkpoint problem are available in hardcopy at lecture or in the Gates B wing open space near Keith's office.
- SCPD students – we'll send you an email with information about solutions soon.

Piazza Questions

- We've gotten a lot of private questions on Piazza that are really interesting and probably super useful to other students.
- If you're asking a Piazza question that doesn't give away hints or answers to problem set questions, consider making it public so that everyone can learn!

GTGTC

- Girls Teaching Girls to Code (GTGTC) has an opening on their admin team.
- They run day-long tutoring sessions where high-school girls get mentorship from Stanford students. At one of their events last year, they had 40 Stanford students mentor 200 high-schoolers!
- Interested? Apply at <http://bit.ly/gtgtc-app>, or contact Jessie Duan at jduan1@stanford.edu.

Your Questions

“What are some of the most interesting/unusual proof techniques?”

“Most of the proofs in the homework can be solved algebraically without the use of words – the algebra suffices as an explanation. Especially when you use proof by contradiction. How do you know you're being superfluous with the explanations?”

“Is a program a proof? If so, suppose we have a program that is guaranteed to prove or disprove some fact. Assume that it is guaranteed to terminate, but will only do so thousands of years from now. Is it still a proof?”


“Keith, why did you choose academia over industry?”

Back to CS103!

Example: Tournaments


Tournaments

- A ***tournament*** is a contest for $n \geq 1$ people.
- Each person plays exactly one game against each other person, and there are no ties.
- The result can be visualized in a picture like this one, which is called a ***tournament graph***.




Victory Chains

- A ***victory chain*** in a tournament is a way of lining up the players so that every player beat the player that comes after them.




Victory Chains

- A ***victory chain*** in a tournament is a way of lining up the players so that every player beat the player that comes after them.




Victory Chains

- A ***victory chain*** in a tournament is a way of lining up the players so that every player beat the player that comes after them.



Victory Chains

- A ***victory chain*** in a tournament is a way of lining up the players so that every player beat the player that comes after them.




Theorem: Every tournament, regardless of the outcome has a victory chain.


Thinking Inductively


- The inductive step in an inductive proof uses the fact that the result is true for a smaller number (k) to prove that the result is true for a larger number ($k+1$).
- In most inductive proofs, the proof that the result is true for $k+1$ explicitly tries to simplify the $k+1$ case into the k case.
 - Counterfeit coins: Turn $k+1$ weighings into k weighings.
 - **MU** puzzle: Turn a sequence of $k+1$ events into a sequence of k events.
 - Triangulation: Turn a polygon of $k+1$ vertices into one with k vertices.
 - Square subdivision: Use a subdivision into k to get one for $k+3$.


Thinking Inductively


- For our victory chain proof, we will simplify the problem by turning the larger tournament into two smaller tournaments.
- We'll inductively argue that, since those smaller tournaments each have victory chains, the larger tournament must have a victory chain.






































The Proof, Schematically




The Proof, Schematically




The Proof, Schematically



The Proof, Schematically



The Proof, Schematically



The Idea

- Suppose that every tournament with at most k players has a victory chain.
- Take a tournament T with $k+1$ players.
- Choose any one player p .
- Form the subtournaments T_0 and T_1 of all players who beat p and lost to p , respectively.
- Get victory chains from T_0 and T_1 .
- Splice those chains together through p .

The Idea

Suppose that every tournament with at most k players has a victory chain.

Take a tournament T with $k+1$ players.

Choose any one player p .

- Form the subtournaments T_0 and T_1 of all players who beat p and lost to p , respectively.
- Get victory chains from T_0 and T_1 .

Splice those chains together through p .

The Idea

Suppose that every tournament with k players has a victory chain.

Take a tournament T with n players.

Choose any one player p .

This is the key idea behind an inductive proof – we're reducing the problem to smaller copies of itself.

- Form the subtournaments T_0 and T_1 of all players who beat p and lost to p , respectively.
- Get victory chains from T_0 and T_1 .

Splice those chains together through p .

Writing the Proof: A First Attempt

We're going to run into trouble in the middle of this proof. Don't worry – we'll see how to fix it.

Theorem: Every tournament has a victory chain.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.”

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p .

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we **assume that any tournament with k players has a victory chain.** We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . **Both of these tournaments have between 0 and k players.**

Theorem: Every

Proof: Let $P(n)$ be the statement that every tournament with n players has a victory chain. We want to prove $P(n)$ for all $n \in \mathbb{N}$, from

As a base case, $P(0)$ is true. A tournament with 0 players has a victory chain consisting of all 0 of the players. For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

At this point, we're stuck. We know that tournaments with exactly k players must have a victory chain, but we're not assuming anything about tournaments with $0, 1, 2, \dots, k-1$ players. Therefore, we can't necessarily say anything about these subtournaments.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will show that $P(n)$ is true for all $n \in \mathbb{N}$, from

What if we made some additional assumptions so that we can say something about these smaller tournaments?

As a base case, consider a tournament with 0 players. This tournament has a victory chain consisting of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we **assume that any tournament with k players has a victory chain.** We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . **Both of these tournaments have between 0 and k players.**

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(k)$ is true. In other words, we assume that any tournament with k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0)$, $P(1)$, $P(2)$, ..., and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0)$, $P(1)$, $P(2)$, ..., and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from $n = 0$ onwards.

We're now assuming that the result is true for $0, 1, 2, 3, \dots, k$. Now, we can continue to make progress!

As a base case, consider a tournament with 0 players. This tournament vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0), P(1), P(2), \dots$, and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0)$, $P(1)$, $P(2)$, ..., and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0)$, $P(1)$, $P(2)$, ..., and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players. Therefore, by our inductive hypothesis, these two subtournaments have victory chains, call them V_0 and V_1 .

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0)$, $P(1)$, $P(2)$, ..., and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players. Therefore, by our inductive hypothesis, these two subtournaments have victory chains, call them V_0 and V_1 . If we then splice together these chains to form the chain V_0, p, V_1 , then we have a victory chain for T : every player is present, and every player beat the player immediately after them.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0)$, $P(1)$, $P(2)$, ..., and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players. Therefore, by our inductive hypothesis, these two subtournaments have victory chains, call them V_0 and V_1 . If we then splice together these chains to form the chain V_0, p, V_1 , then we have a victory chain for T : every player is present, and every player beat the player immediately after them. Therefore, this arbitrary tournament of $k+1$ players has a victory chain, so $P(k+1)$ is true, completing the induction.

Theorem: Every tournament has a victory chain.

Proof: Let $P(n)$ be the statement “every tournament with n players has a victory chain.” We will prove by induction that $P(n)$ is true for all $n \in \mathbb{N}$, from which the theorem follows.

As a base case, we prove $P(0)$, that any tournament with no players has a victory chain. In a tournament with no players, a list of all 0 of the players vacuously satisfies the claim that every player beat the next player in the list. Therefore, $P(0)$ is true.

For the inductive step, assume that for some $k \in \mathbb{N}$ that $P(0)$, $P(1)$, $P(2)$, ..., and $P(k)$ are all true. In other words, we assume that any tournament with at most k players has a victory chain. We'll prove $P(k+1)$, that any tournament with $k+1$ players has a victory chain.

Consider any tournament T with $k+1$ players. Choose any one player p and form two subtournaments, a subtournament T_0 of all the players who beat p and a subtournament T_1 of all the players who lost to p . Both of these tournaments have between 0 and k players. Therefore, by our inductive hypothesis, these two subtournaments have victory chains, call them V_0 and V_1 . If we then splice together these chains to form the chain V_0, p, V_1 , then we have a victory chain for T : every player is present, and every player beat the player immediately after them. Therefore, this arbitrary tournament of $k+1$ players has a victory chain, so $P(k+1)$ is true, completing the induction. ■

What We Just Did

- In a normal inductive step, we assume that $P(k)$ is true and prove $P(k+1)$.
- In this type of inductive step, we assume $P(0)$, $P(1)$, ..., and $P(k)$ are true before we prove $P(k+1)$.
- That way, when we found *any* kind of smaller tournament, we knew something about its structure.
- This type of proof has a name!

Complete Induction

- If the following are true:
 - $P(0)$ is true, and
 - If $P(0), P(1), P(2), \dots, P(k)$ are true, then $P(k+1)$ is true as well.



then $P(n)$ is true for all $n \in \mathbb{N}$.

- This is called the ***principle of complete induction*** or the ***principle of strong induction***.
 - (A note: this also works starting from a number other than 0; just modify what you're assuming appropriately.)


That's a *lot* of assumptions to make!

Why is this legal?


Review: Induction as a Machine





Review: Induction as a Machine




Review: Induction as a Machine




Review: Induction as a Machine





Review: Induction as a Machine




Review: Induction as a Machine




Review: Induction as a Machine



Review: Induction as a Machine





Review: Induction as a Machine





An Observation

An Observation





$P(0)$



An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation





An Observation




An Observation




Intuiting Complete Induction





Intuiting Complete Induction




Intuiting Complete Induction




Intuiting Complete Induction





Intuiting Complete Induction




Intuiting Complete Induction




Intuiting Complete Induction





Intuiting Complete Induction



Intuiting Complete Induction



Intuiting Complete Induction



More on Complete Induction

- This type of induction can seem too powerful, almost like it's cheating.
- It's actually perfectly safe!
- We'll do more examples when we come back next time.

Next Time

- **More on Complete Induction**
 - Building an intuition for complete induction.
 - More applications!
- **Graphs**
 - Representing relationships between objects.
 - Graph connectivity.