

Investigation of Dust-Ion Acoustic Solitary Structures in a Weakly Relativistic Plasma

S. Ramesh, C. Verma, and S. Chandra

Abstract—In this paper, analytical investigation of dust-ion acoustic waves in a weak relativistically degenerate plasma, with mobile negatively charged dust particles and usual ions and electron components is carried out. The Kortweg-de Vries equation is also derived, to study solitary waves, using relevant stretched coordinates and perturbation expansion.

PACS—52.27.Lw; 52.27.Ny; 52.25.Ya.

Keywords—Dust ion acoustic wave; KdV equation; solitary structures.

Nomenclature—KdV: Korteweg-de Vries

I. INTRODUCTION

DUST ion acoustic waves are an expanding area of research, and with the recognition of quantum and relativistic effects involved in these phenomena, more opportunities come up for investigation of these waves in various regimes. A number of analytical treatments have been carried out already regarding these waves over the past years. More importantly, dust plasmas have been observed in active galactic nuclei [1], laser fields [2], pulsar magnetospheres [3] and polar regions of neutron stars [4], making it a huge region of interest for astrophysics. There is also evidence for presence of these dustplasmas in the early stages of the Universe [5], [6], [7] and the centre of the milky way galaxy [8]. Occurrences of relativistic plasma were found in Van Allen radiation belt [9] and the boundary layer of earths magnetosphere too [10]. This has led to extensive research concerning these waves. Dierent types of dust acoustic waves have been studied by Mamun and Shukla (2002) [11], Rao et al (1990) [12] and others. Rao et al characterized the linear normal modes and nonlinear supersonic solitons and Mamun and Shukla characterized spherical and cylindrical waves as opposed to one dimensional waves. These waves have also been studied with various pseudopotential approaches, such as in Pakzad (2009) [13]. Magnetized plasmas have also been found to have rarefactive solitons in Kundu et al (2013) [14] where soliton width decreases with magnitude of the eld. HF Liu et al (2010) [15] have used a modied KdV-Burgers equation to model cylindrical and spherical dust ion acoustic waves in weakly relativistic plasma. Solitary structures are observed in these models that vary with density of the components, the mass of the constituent particles and temperatures of the components. Chetia [16] studied compressive and rarefactive Ion-Acoustic

solitons. Also, Chandra, Paul, and Ghosh [17] have showed that relativistic degeneracy parameter significantly inuences the conditions of formation and properties of solitary structures. In this paper, we have considered the model consisting of mobile negatively charged dust particles, singly charged ions and electrons to derive the linear dispersion relation and Korteweg-de Vries equations in weakly relativistic regime using standard perturbation techniques. The effects of varying temperatures, dust charge and other parameters are studied. We have divided the paper in five sections (more or less). The second section throws some light on the basic quantum hydrodynamic, and three uid equations; along with a relativistically degenerate pressure scheme. The next section focusses on the derivation of linear dispersion relation. We have also derived the KdV equation, and its solution for the same set of equations in the third section. We discuss some graphical properties of the above derived equations in the fourth section.

II. BASIC FORMULATION

In order to analytically study the propagation of solitary waves in an unmagnetized three component weak relativistically degenerate plasma we start with a set of interpenetrating fluids (three, here) characterized by :

- 1) The equations of continuity and motion of
 - ions,
 - electrons, and
 - dust particles
- 2) The Poisson's equation
- 3) The Pressure Law

A. Governing Equations

- For Dust particles:

$$\frac{\partial n_d}{\partial t} + \frac{\partial(n_d u_d)}{\partial x} = 0 \quad (1)$$

$$\frac{\partial u_d}{\partial t} + u_d \frac{\partial u_d}{\partial x} = \frac{Q_d}{m_d} \frac{\partial \phi}{\partial x} \quad (2)$$

- For Ions:

$$\frac{\partial n_i}{\partial t} + \frac{\partial(n_i u_i)}{\partial x} = 0 \quad (3)$$

$$\frac{\partial u_i}{\partial t} + u_i \frac{\partial u_i}{\partial x} = -\frac{Q_i}{m_i} \frac{\partial \phi}{\partial x} - \frac{1}{m_i n_i} \frac{\partial p_i}{\partial x} \quad (4)$$

- For Electrons:

$$\frac{\partial n_e}{\partial t} + \frac{\partial(n_e u_e)}{\partial x} = 0 \quad (5)$$

Shashank Ramesh is with Physics Department, Kirori Mal College, University of Delhi, India (email - shashankramesh38@hotmail.com)

Chirag Verma is with Physics Department, Kirori Mal College, University of Delhi, India. (email - chirag2000verma@gmail.com)

Swarniv Chandra was the program coordinator for the Summer Workshop on Plasma Physics 2020 (e-mail: swarniv147@gmail.com).

$$\frac{\partial u_e}{\partial t} + u_e \frac{\partial u_e}{\partial x} = -\frac{Q_e}{m_e} \frac{\partial \phi}{\partial x} - \frac{1}{m_e n_e} \frac{\partial p_e}{\partial x} + \frac{\hbar^2}{2m_e^2} \frac{\partial}{\partial x} \left(\frac{1}{\sqrt{n_e}} \frac{\partial^2 \sqrt{n_e}}{\partial x^2} \right) \quad (6)$$

• Poisson's Equation:

$$\frac{\partial^2 \phi}{\partial x^2} = 4\pi(Q_e n_e - Q_i n_i + Q_d n_d) \quad (7)$$

B. Pressure Law

Following from Chandrasekhar (1939), The electron degeneracy pressure in fully degenerate and relativistic form can be expressed as :

$$p_j = (\pi m_e^4 c^5 / 3h^3) [R_j (2R_j^2 - 3) \sqrt{1 + R_j^2} + 3 \sinh^{-1} R_j] \quad (8)$$

$$R_j = p_{Fj} / m_e c = [3h^3 n_j / 8\pi m_e^3 c^3]^{1/3} = R_{j0} n_j^{1/3} \quad (9)$$

$$R_{j0} = (n_{j0} / n_0)^{1/3} = 5.9 \cdot 10^{29} \text{ cm}^{-1}$$

For weak-relativistic degeneracy : $R_j \rightarrow 0$, and, hence

$$p_j = \frac{1}{20} \left(\frac{3}{\pi} \right)^{2/3} \frac{\hbar^2}{m_e} n_j^{5/3}$$

where the suffices d, e, i stand for dust particle, electron, and ion components respectively; u_j and p_j are respectively the fluid velocity and degeneracy pressure of the j^{th} species, i.e. electron or dust particle or ion; ϕ is the electrostatic wave potential; and Q_j is the charge on the j^{th} species.

C. Normalized Equations

Under the following normalized scheme: $n_j \rightarrow n_j / n_{i0}$, $t \rightarrow t \omega_{pd}$, $x \rightarrow x / \lambda_{De}$, $u_j \rightarrow u_j / C_d$, $\phi \rightarrow K_b T_e / m_d$

where $\omega_{pd} = (m_d / 4\pi n_{i0} e^2)^{1/2}$ is the characteristic dust plasma frequency, $\lambda_{De} = (K_b T_e / 4\pi n_{i0} e^2)^{1/2}$ is the Debye length, $C_d = (K_b T_e / m_d)^{1/2}$ is the dust acoustic speed; the above equations are then normalized to:

$$\frac{\partial n_d}{\partial t} + \frac{\partial(n_d u_d)}{\partial x} = 0 \quad (10)$$

$$\frac{\partial u_d}{\partial t} + u_d \frac{\partial u_d}{\partial x} = -Z_d \frac{\partial \phi}{\partial x} \quad (11)$$

$$\frac{\partial n_i}{\partial t} + \frac{\partial(n_i u_i)}{\partial x} = 0 \quad (12)$$

$$\frac{\partial u_i}{\partial t} + u_i \frac{\partial u_i}{\partial x} = -\frac{1}{Q} \left(\frac{\partial \phi}{\partial x} + \frac{\alpha}{n_i} \frac{\partial p_i}{\partial x} \right) \quad (13)$$

$$\frac{\partial n_e}{\partial t} + \frac{\partial(n_e u_e)}{\partial x} = 0 \quad (14)$$

$$\frac{\partial u_e}{\partial t} + u_e \frac{\partial u_e}{\partial x} = -\frac{1}{P} \left(\frac{\partial \phi}{\partial x} + \frac{1}{n_e} \frac{\partial p_e}{\partial x} \right) + \frac{1}{P^2} \frac{H^2}{2} \frac{\partial}{\partial x} \left(\frac{1}{\sqrt{n_e}} \frac{\partial^2 \sqrt{n_e}}{\partial x^2} \right) \quad (15)$$

$$\frac{\partial^2 \phi}{\partial x^2} = (n_e - n_i + Z_d n_d) \quad (16)$$

where $Q = \frac{m_i}{m_d}$, $P = \frac{m_e}{m_d}$, $H = \frac{\hbar \omega_{pd}}{K_b T_e}$, $\alpha = \frac{T_i}{T_e}$

III. ANALYTICAL STUDY

In order to investigate the nonlinear behaviour of electron-acoustic waves we make the following perturbation expansion, about their equilibrium quantities, for the aforementioned field quantities:

$$\begin{bmatrix} n_d \\ n_i \\ n_e \\ u_d \\ u_e \\ u_i \\ \phi \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ u_{0d} \\ u_{0e} \\ u_{0i} \\ \phi_0 \end{bmatrix} + \epsilon \begin{bmatrix} n_d^{(1)} \\ n_e^{(1)} \\ n_i^{(1)} \\ u_d^{(1)} \\ u_e^{(1)} \\ u_i^{(1)} \\ \phi^{(1)} \end{bmatrix} + \epsilon^2 \begin{bmatrix} n_d^{(2)} \\ n_e^{(2)} \\ n_i^{(2)} \\ u_d^{(2)} \\ u_e^{(2)} \\ u_i^{(2)} \\ \phi^{(2)} \end{bmatrix} + \dots \quad (17)$$

A. Dispersion Characteristics

Substituting the expansion (17) in equations (10)-(16), and then linearizing and assuming that all field quantities vary as $e^{i(kx - \omega t)}$, we get for normalized wave frequency ω and wave number k , the following linear dispersion relation:

$$-1 = \frac{1}{P\omega^2 - \frac{5}{3}\alpha F_e k^2 - \frac{H^2 k^4}{4P}} - \frac{1}{Q\omega^2 - \frac{5}{3}\alpha F_i k^2} + \frac{Z_d^2}{\omega^2} \quad (18)$$

where $F_i = F_e = \frac{1}{20} \left(\frac{3}{\pi} \right)^{2/3} \frac{\hbar^2}{m_e}$

In the long wavelength range, we ignore k^4 and equation (18) becomes quadratic in ω^2 as follows:

$$0 = PQ\omega^4 + \left(Q - \frac{5}{3}P\alpha F_i k^2 - \frac{5}{3}QF_e k^2 - P - Z_d^2 PQ \right) \omega^2 + \left(\frac{5}{3}F_e k^2 - \frac{5}{3}\alpha F_i k^2 + \frac{5}{3}Z_d^2 \alpha P F_i k^2 + \frac{5}{3}Z_d^2 Q F_e k^2 \right) \quad (19)$$

The solutions of equation (19) are :

$$\omega_{1,2}^2 = -b \pm \sqrt{b^2 - 4c}$$

where

$$b = \frac{1}{P} - \frac{5}{3} \frac{1}{Q} \alpha F_i k^2 - \frac{5}{3} \frac{1}{P} F_e k^2 - P - Z_d^2$$

$$c = \frac{5}{3} \frac{1}{PQ} F_e k^2 - \frac{5}{3} \frac{1}{PQ} \alpha F_i k^2 + \frac{5}{3} \frac{1}{Q} Z_d^2 \alpha F_i k^2 + \frac{5}{3} \frac{1}{P} Z_d^2 F_e k^2 \quad (13)$$




Fig. 1: ω vs k for different values of H : $F_i = F_e = 1.517, \alpha = 0.01$




Fig. 3: ω vs k for different values of α : $F_i = 1.517, Q = 0.1, H = 2, k = 0.5$




Fig. 2: ω vs k for different values of Q : $F_i = 1.517, \alpha = 0.01, Z_d = 100, H = 2, k = 0.5$

B. KdV Equation

We would be using standard reductive perturbation technique, which is: $\xi = \varepsilon^{1/2}(x - V_0 t)$ and $\tau = \varepsilon^{3/2}t$ where V_0 is the normalized linear long wave phase velocity given by equation (10), and ε is the smallness parameter measuring the dispersion and nonlinear effects. Now writing the equations (10)-(16) in the terms of these stretched coordinates ξ and τ , and applying the perturbation expansion (17), and then solving for the lowest order equation with boundary conditions that $n_j^{(1)}, u_j^{(1)}, \phi^{(1)} \rightarrow 0$ as $|\xi| \rightarrow 0$, the following solutions are obtained :

$$\begin{aligned} u_d^{(1)} &= \frac{Z_d}{M_d} \phi^{(1)}, n_d^{(1)} = \frac{Z_d}{M_d^2} \phi^{(1)} \\ u_i^{(1)} &= \frac{M_i}{A} \phi^{(1)}, n_i^{(1)} = \frac{1}{A} \phi^{(1)} \\ u_e^{(1)} &= \frac{M_e}{B} \phi^{(1)}, n_e^{(1)} = \frac{1}{B} \phi^{(1)} \end{aligned} \quad (20)$$

where

$$M_d = V_0 - u_{0d} \quad (21)$$

$$M_i = V_0 - u_{0i} \quad (22)$$

$$M_e = V_0 - u_{0e} \quad (23)$$

$$A = Q M_i^2 - \frac{5}{3} \alpha F_i \quad (24)$$

$$B = P M_e^2 - \frac{5}{3} F_e \quad (25)$$

Going for the next higher order terms in ε and following the usual method the desired Korteweg-de-Vries equation is obtained :

$$\frac{\partial \phi}{\partial \tau} + N \phi \frac{\partial \phi}{\partial \xi} + D \frac{\partial^3 \phi}{\partial \xi^3} = 0 \quad (26)$$

where

$$N = \frac{\frac{3Z_d^3}{M_d^4} + \frac{3PM_e^2}{B^3} - \frac{3QM_i^2}{A^3} + \frac{5\alpha F_i}{9A^3} - \frac{5F_i}{9B^3}}{\frac{2Z_d^2}{M_d^3} + \frac{2PM_e}{B^2} - \frac{2QM_i}{A^2}}$$

$$D = \frac{1 + \frac{H^2}{2PB^2}}{\frac{2Z_d^2}{M_d^3} + \frac{2PM_e}{B^2} - \frac{2QM_i}{A^2}}$$

C. Solution to KdV Equation

Consider the new variable $\eta = \xi - M\tau$ where M is the normalized constant speed of the wave frame. On applying the boundary conditions as $\eta \rightarrow \pm\infty; \phi, \frac{\partial \phi}{\partial \eta}, \frac{\partial^2 \phi}{\partial \eta^2} \rightarrow 0$, the possible stationary solution of equation (3.8) is :

$$\phi = \phi_m \operatorname{sech}^2\left(\frac{\eta}{\Delta}\right) \quad (27)$$

where the amplitude ϕ_m and the width Δ of the soliton are given by :

$$\phi_m = 3 \frac{M}{N} \text{ and } \Delta = \sqrt{\frac{4D}{M}}$$




Fig. 4: ϕ vs η : $k = 0.8, u_0 = 1, \omega = 0.1892, V = 10, M_j = V - u_0, M_n = 1$

IV. CONCLUSION

A. Results and discussion

Using the model described above and standard perturbative techniques, a linear dispersion relation was found taking into account quantum effects of electrons and relativistic effects.

Figure 1 shows variation of ω with wave number k for different values of H . ω initially has a higher degree variation with respect to wave number k but shows linear variation at larger values of k ($k > 0.1$) for fixed values of other parameters. Initially the k^4 term is significant, but its influence disappears as k increases, validating our assumption to simplify the linear dispersion relation earlier. Also, the wave number has a slight decrease with the increase in quantum diffraction parameter H .


Figure 2 shows variation of ω with k for different values of Q . ω decreases with increase in Q . This is due to dependence of constant A on Q , which is quite large.

Figure 3 shows variation of ω with k for different value of temperature ratio α , where ω increases with increase in α .


The KdV equation was derived to study nonlinear characteristics of the waves, where the coefficients are found to have been modified by the quantum and relativistic effects.

Figure 4 shows the soliton profiles obtained for different values of Q . The amplitude of the soliton decreases with increase in Q .


Figure 5(a) shows variation of amplitude with temperature ratio for different values of Q . The amplitude decreases with increase in α , and as observed earlier there is an overall reduction in amplitude with increase in Q .



(a) ϕ_0 vs α : $Z = 100, P = 1/9000, Q = 1/10, F = 1.517, H = 3, k = 0.4, u_0 = 0.3, V = 10, M_j = V - u_0, M_n = 1$




(b) ϕ_0 vs Q : $Z = 100, P = 1/9000, F_i = 1.517, a = 0.01, H = 3, k = 0.4, u_0 = 0.3, V = 10, M_j = V - u_0, M_n = 1$




(c) ϕ_0 vs Z_d : $P = 1/9000, Q = 1/10, F_i = 1.517, a = 0.01, H = 3, k = 0.4, u_0 = 0.3, V = 10, M_j = V - u_0, M_n = 1$


Fig. 5: Variation of wave's amplitude with different quantities



(a) Δ vs Q



(b) Δ vs Z_d



(c) Δ vs α for different values of Q

Fig. 6: Variation of soliton width with various quantities.

Figure 5(b) shows variation of amplitude with Q . The initial rate of change is small but gets steeper as value of Q increases.

Figure 5(c) shows variation of amplitude with dust charge Z_d . The amplitude decreases with increase in Z_d for fixed values of other parameters.

Figure 6 shows the variation of the soliton width with ion-dust mass ratio, dust charge temperature ratio. Clearly, Δ increases with α but decreases with Z_d and Q .

B. Concluding remarks

The properties and characteristics of dust-ion acoustic waves in weakly relativistic plasma were investigated. The properties of the solitons under consideration depends mainly on the ion-dust mass ratio, the dust charge and temperature ratio. The investigation of these waves may prove helpful in astrophysical studies of interstellar media, astronomical objects such as neutron stars and laser interaction experiments where the additional quantum and relativistic effects may become influential.

V. ACKNOWLEDGEMENTS

We would like to thank our project coordinator, Dr. Swarniv Chandra, for providing us with unending inspiration, motivation, and guidance. We would also like to thank our respective families for their constant support.

REFERENCES

- [1] H. R. Miller and P. J. Wiita, *Active galactic nuclei. Proceedings of a conference held at the Georgia State University, Atlanta, Georgia. 28-30 October, 1987.* Berlin, 1987.
- [2] V. I. Berezhiani, D. D. Tskhakaya, and P. K. Shukla, "Pair production in a strong wake field driven by an intense short laser pulse," *Physical Review A*, vol. 46, no. 10, pp. 6608–6612, 1992.
- [3] F. C. Michel, "Theory of pulsar magnetospheres," *Reviews of Modern Physics*, vol. 54, no. 1, pp. 1–66, 1982.
- [4] F. C. Michel, *Theory of neutron star magnetospheres.* University of Chicago Press, 1991.
- [5] S. Weinberg, *Gravitation and cosmology.* 1972.
- [6] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation.* Freeman, 1973.
- [7] J. Rees, M., *In the Very Early Universe.* Cambridge: Cambridge University Press, 1983.
- [8] M. L. Burns, *Positron-electron pairs in astrophysics.* American Institute of Physics, 1983.
- [9] H. Ikezi, "Experiments on ion-acoustic solitary waves," *Physics of Fluids*, vol. 16, no. 10, p. 1668, 1973.
- [10] J. I. Vette, *Summary of particle population in the magnetosphere.* Goddard Space Flight Center, 1969.
- [11] P. K. Shukla and A. A. Mamun, "Introduction to dusty plasma physics," *Plasma Physics and Controlled Fusion*, vol. 44, no. 3, pp. 395–395, 2002.
- [12] N. Rao, P. Shukla, and M. Yu, "Dust-acoustic waves in dusty plasmas," *Planetary and Space Science*, vol. 38, no. 4, pp. 543–546, 1990.
- [13] H. R. Pakzad and K. Javidan, "Dust acoustic solitary and shock waves in strongly coupled dusty plasmas with nonthermal ions," *Pramana*, vol. 73, no. 5, pp. 913–926, 2009.
- [14] N. R. Kundu, M. M. Masud, K. S. Ashrafi, and A. A. Mamun, "Dust-ion-acoustic solitary waves and their multi-dimensional instability in a magnetized nonthermal dusty electronegative plasma," *Astrophysics and Space Science*, vol. 343, no. 1, pp. 279–287, 2012.
- [15] H. F. Liu, S. Q. Wang, C. Z. Li, Q. Xiang, F. Z. Yang, and Y. Liu, "Cylindrical and spherical dust ion acoustic solitary waves in a relativistic dust plasma," *Physica Scripta*, vol. 82, no. 6, p. 065402, 2010.

- [16] S. Chetia, "Ion acoustic waves in a unidirected dusty plasma," *International Journal Of Mathematics Trends And Technology*, vol. 56(1), pp. 21–27, 2018.
- [17] P. S. . G. B. Chandra, S., "Electron-acoustic solitary waves in a relativistically degenerate quantum plasma with two-temperature electrons," *Astrophysics And Space Science*, vol. 343(1), pp. 213–219, 2012.



Shashank Ramesh Shashank Ramesh is pursuing his B.Sc.(H) Physics at Kirori Mal College, University of Delhi. His academic interests lie in plasma physics and its applications in astrophysics.



Chirag Verma Chirag Verma is pursuing his B.Sc.(H) Physics at Kirori Mal College, University of Delhi. He has varied interests in theoretical and mathematical physics.