

# **Kỷ yếu hội nghị các nhà khoa học trẻ**

**Tiểu ban: Vật lý hạt nhân và ứng dụng**

# Spectroscopic information about a hypothetical tetrahedral configuration in $^{156}\text{Gd}$

Q. T. Doan<sup>i</sup>, O. Stezowski<sup>i,j</sup>, D. Guinet<sup>i,j</sup>, J. Dudek<sup>k</sup>, D. Curien<sup>k</sup>, N. Schunck<sup>l,3</sup>

<sup>a</sup>*Institut de Physique Nucléaire de Lyon, France*

<sup>b</sup>*Université de Lyon, Université Lyon 1, Lyon, France*

<sup>c</sup>*Département de Recherches Subatomiques, Institut Pluridisciplinaire Hubert Curien, Strasbourg, France*

<sup>d</sup>*Institute of Nuclear Physics PAN, PL-31-342 Kraków, Poland*

## Abstract

A detailed  $\gamma$ -ray spectroscopy of the lowest two negative-parity bands in  $^{156}\text{Gd}$  has been performed ...

**Keywords:**  $\gamma$ -ray spectroscopy,  $^{156}\text{Gd}$

## 1. Introduction

Theoretical studies based on the nuclear mean-field approach and group theory considerations suggest [1, 2] that some atomic nuclei may exhibit tetrahedral and/or octahedral symmetries. To the lowest order, tetrahedral symmetry is realized through octupole deformation  $Y3 \pm 2$  of the nuclear surface. Previous research[3] has determined magic numbers for which tetrahedral deformation should be the easiest to observe leading to tetrahedral proton and neutron "magic" numbers  $Z_t/N_t = 32, 40, 56, 64, 70, 90$ , and 112, with extra gaps at  $N_t = 136$  and 142. The authors of Refs. [1, 2, 3] have furthermore demonstrated that nuclei with an exact tetrahedral symmetry have all multipole moments  $Q_\lambda < 7$ ,  $\nu = 0$  except for  $Q_{32}$  - thus, in particular, the corresponding quadrupole moments  $Q_2$  vanish.

.....  
The radiation intensity can be written down using the standard expression in the Equation B.1.

$$I(\theta) = 1 + A_2 P_2(\cos \theta) + A_4 P_4(\cos \theta) \quad (1)$$


\*Corresponding author

Email addresses: qtdoan@gmail.com (Q. T. Doan),  
qtdoan@gmail.com (Q. T. Doan)

<sup>1</sup>Present address: Lawrence Livermore National Laboratory,  
L-414, P.O. Box 808, Livermore, CA 94551, USA

Preprint submitted to Nuclear Physics B

## 2. Experiment setup



Hình 1: Partial level scheme of  $^{156}\text{Gd}$ .

## 3. Spectroscopy information

Transition intensities are presented in the Figure C.2 ...

## 4. Acknowledgements

This work benefited from the TNT2-D cards, developed and financed by CNRS/IN2P3 for the GABRIELA project ...

## Phụ lục A. Experiment data

Experiment data are shown in the Table A.2 ...

Ngày 22 tháng 4 năm 2016

Bảng A.1: Data $\gamma$ ray detection.		
Interaction	Clover	Cluster
1	74.5	79.8
2	22.4	18.3
$\geq 3$	$<3.3$	$<2.8$

## References

- [1] J. Dudek, et al., Phys. Rev. Lett. 97 (072501).
- [2] N. Schunck, et al., Acta Phys. Pol. B 36 (1071).
- [3] J. Dudek, et al., Phys. Rev. Lett. 88 (252502).

# Spectroscopic information about a hypothetical tetrahedral configuration in $^{156}\text{Gd}$

Q. T. Doan<sup>i</sup>, O. Stezowski<sup>i,j</sup>, D. Guinet<sup>i,j</sup>, J. Dudek<sup>k</sup>, D. Curien<sup>k</sup>, N. Schunck<sup>l,3</sup>, Q. T. Doan<sup>i</sup>, O. Stezowski<sup>i,j</sup>, D. Guinet<sup>i,j</sup>, J. Dudek<sup>k</sup>, D. Curien<sup>k</sup>, N. Schunck<sup>l,3</sup>

<sup>e</sup>*Institut de Physique Nucléaire de Lyon, France*

<sup>f</sup>*Université de Lyon, Université Lyon 1, Lyon, France*

<sup>g</sup>*Département de Recherches Subatomiques, Institut Pluridisciplinaire Hubert Curien, Strasbourg, France*

<sup>h</sup>*Institute of Nuclear Physics PAN, PL-31-342 Kraków, Poland*

<sup>i</sup>*Institut de Physique Nucléaire de Lyon, France*

<sup>j</sup>*Université de Lyon, Université Lyon 1, Lyon, France*

<sup>k</sup>*Département de Recherches Subatomiques, Institut Pluridisciplinaire Hubert Curien, Strasbourg, France*

<sup>l</sup>*Institute of Nuclear Physics PAN, PL-31-342 Kraków, Poland*

## Abstract

A detailed  $\gamma$ -ray spectroscopy of the lowest two negative-parity bands in  $^{156}\text{Gd}$  has been performed ...

**Keywords:**  $\gamma$ -ray spectroscopy,  $^{156}\text{Gd}$

## Phụ lụcB. Introduction


Theoretical studies based on the nuclear mean-field approach and group theory considerations suggest [1, 2] that some atomic nuclei may exhibit tetrahedral and/or octahedral symmetries. To the lowest order, tetrahedral symmetry is realized through octupole deformation  $Y3 \pm 2$  of the nuclear surface. Previous research[3] has determined magic numbers for which tetrahedral deformation should be the easiest to observe leading to tetrahedral proton and neutron "magic" numbers  $Zt/Nt = 32, 40, 56, 64, 70, 90$ , and 112, with extra gaps at  $Nt = 136$  and 142. The authors of Refs. [1, 2, 3] have furthermore demonstrated that nuclei with an exact tetrahedral symmetry have all multipole moments  $Q_\lambda < 7$ ,  $\nu = 0$  except for  $Q_{32}$  -

thus, in particular, the corresponding quadrupole moments  $Q_2$  vanish.

.....  
The radiation intensity can be written down using the standard expression in the Equation B.1.

$$I(\theta) = 1 + A_2 P_2(\cos \theta) + A_4 P_4(\cos \theta) \quad (\text{B.1})$$

## Phụ lụcC. Experiment setup



Hình C.2: Partial level scheme of  $^{156}\text{Gd}$ .

\*\*Corresponding author

Corresponding author

Email addresses: qtdoan@gmail.com (Q. T. Doan),  
qtdoan@gmail.com (Q. T. Doan)

<sup>2</sup>Present address: Lawrence Livermore National Laboratory,  
L-414, P.O. Box 808, Livermore, CA 94551, USA

<sup>3</sup>Present address: Lawrence Livermore National Laboratory,  
L-414, P.O. Box 808, Livermore, CA 94551, USA

## Phụ lụcD. Spectroscopy information

Transition intensities are presented in the Figure C.2 ...

## Phụ lụcE. Acknowledgements

This work benefited from the TNT2-D cards, developed and financed by CNRS/IN2P3 for the GABRIELA project ...

## Phụ lụcA. Experiment data

Experiment data are shown in the Table A.2 ...

Bảng A.2: Data  $\gamma$  ray detection.

Interaction	Clover	Cluster
1	74.5	79.8
2	22.4	18.3
$\geq 3$	$<3.3$	$<2.8$

## References

- [1] J. Dudek, et al., Phys. Rev. Lett. 97 (072501).
- [2] N. Schunck, et al., Acta Phys. Pol. B 36 (1071).
- [3] J. Dudek, et al., Phys. Rev. Lett. 88 (252502).