

Early Universe Discoveries with JWST

Javier Álvarez-Márquez

Centro de Astrobiología (CSIC-INTA) - Madrid
e-mail: javier.alvarez@cab.inta-csic.es



MIRI
European
Consortium




CENTRO DE ASTROBIOLOGÍA

CSIC
CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS

INTA Instituto Nacional de
Técnica Aeroespacial



JWST & Primordial Universe: History of the Universe



Begin Dark Ages ($z \sim 1000$, 0.3 Myr)

Begin Reionization ($z \sim 15-10$, 300 Myr)



First stars, galaxies & QSO

Begin Galaxy Assembly ($z \sim 6$, 1 Gyr)



Formation & evolution of
Galaxies & AGNs

Today ($z=0$)



JWST & Primordial Universe: Detection of galaxies in the EoR pre-JWST



HST

Hubble space telescope

Telescope covering the UV, optical, and near-infrared range

$\lambda \sim 200\text{-}1600 \text{ nm}$

Characterization of the UV spectrum of bright galaxies in the EoR



Spitzer

Spitzer

Telescope covering the near and mid infrared range

$\lambda \sim 3.6\text{-}4.5 \mu\text{m}$

Characterization of the optical spectrum of bright galaxies in the EoR



Atacama Large Millimeter Array


Interferómetro cubriendo el rango submilimétrico y milimétrico

$\lambda \sim 0.4\text{-}8.6 \text{ mm}$

Characterization of the neutral, ionized, and cold molecular gas, as well as dust emission in bright galaxies of the EoR








JWST & Primordial Universe: Detection of galaxies in the EoR pre-JWST






JWST & Primordial Universe: Detection of galaxies in the EoR pre-JWST

- **Deep surveys from HST & Spitzer have detected hundreds of galaxies at $z > 7$**
 - Identified as line emitters (e.g., Stark et al. 2013)
- **Spectroscopic confirmation was challenging prior to JWST**
 - Detection of the $\text{Ly}\alpha$ line from ground-based telescopes/HST (Stark et al. 2017)
 - Detection of far-infrared lines ($[\text{CII}]$ 158 μm or $[\text{OIII}]$ 88 μm) in the submillimeter with ALMA/NOEMA (Hashimoto et al. 2018)
- **Only a few dozen galaxies were spectroscopically confirmed in the EoR before JWST**







JWST & Primordial Universe: Redshift frontier (pre-JWST)




HST &
ALMA





JWST & Primordial Universe: The new wavelength coverage of JWST




JWST allows us to observe the rest-frame UV, optical, and near-infrared spectrum of galaxies in the EoR:

- **NIRCam**: Imaging in the UV and optical
- **MIRI imager**: Imaging in the optical and near-infrared
- **NIRSpec IFS & SMA**: Spectroscopy in the UV and optical
- **MIRI MRS & LRS**: Spectroscopy in the optical and near-infrared



JWST & Primordial Universe: JWST discoveries using NIRCam & NIRSpec



NIRSpec is powerful to confirm galaxies at $z > 9-10$ from the detection of the Ly- α break

Bright optical lines (H β , [OIII]4960, 5008, H α) are not covered by NIRSpec

Most of the cases the depths are not sufficient to detect faint UV/optical lines


Expensive in telescope time for faint sources

Analysis of bright or lensed galaxies help to reduce exposure times and detect emission lines

Curtis-Lake+23; Arabal Haro+23; ...



MACS0647-JD ($z \sim 10.2$; 460Myr)




NIRSpec is powerful to confirm galaxies at $z > 9-10$ from the detection of the Ly- α break

Bright optical lines (H β , [OIII]4960, 5008, H α) are not covered by NIRSpec

Most of the cases the depths are not sufficient to detect faint UV/optical lines

GNz11 ($z \sim 10.6$; 440Myr)



Expensive in telescope time for faint sources

Analysis of bright or lensed galaxies help to reduce exposure times and detect emission lines

Bunker+23; Hsiao+24





JWST & Primordial Universe: JWST discoveries using NIRCam & NIRSpec

The highest spectroscopically confirmed galaxies are located ~ 300 Myr after the Big Bang

- Confirmed by the detection of the Ly- α break with NIRSpec R100
- No detection of emission lines in the UV rest-frame spectra
- Bright UV galaxy with Muv ~ 20.8

JWST is detecting a large number of bright galaxies in the primordial Universe in contradiction of cosmological simulations


Rapid evolution of galaxies in the early Universe






Diversity of morphologies and structure in primordial galaxies:


- Clumpy systems as MACS1149-JD1
- Really compact galaxies as GNz11 and GHZ2
- Merging systems as MACS0647-JD
- Extended systems as JADES-GS-z14-0




MACS1149-JD ($z \sim 9.1$)




MACS0647-JD ($z \sim 10.2$)



GNz11 ($z \sim 10.6$)



JADES-GS-z14-0 ($z \sim 14.3$)





JWST & Primordial Universe: The relevance of MIRI spectroscopy


UV and optical lines are key ingredients to investigate the physical properties galaxies:

- Tracers of star-formation, metallicity, dust attenuation, radiation sources, electron temperature and density, ...

For primordial Galaxies ($z > 9-10$):

- Bright optical emission lines ($H\beta$, $[OIII]$, $H\alpha$) enter in the MIRI wavelength domain
- NIRSpec only covers the UV to Balmer Break spectral range with faint emission lines

MIRI has a relevant role to characterize that primordial galaxies





Spatially-resolved H α emission

- Two clumps (S & N)
- High (~15) SNR



Spectrally-resolved H α emission

- Dynamical mass of $2.4 \times 10^9 M_{\odot}$
- Different kinematics in each clump

Larger ionizing photon production efficiency

Large variations of the EW of H α between N and S clumps indicating differences in SFHs:

- Clump N, $EW_0(H\alpha) > 2000 \text{ \AA}$, dominated by a young SF burst (<5 Myrs)
- Clump S, $EW_0(H\alpha) \sim 500 \text{ \AA}$, SF over a larger period of time (~50 Myr)





JWST & Primordial Universe: MACS0647-JD ($z \sim 10.2$)

MRS observed optical [OIII]5008 and H α lines with SNR ~ 10

- Spectrally-resolved (FWHM ~ 160 km/s)




NIRSpec have detected faint UV/optical emission lines

Most distant galaxy with a direct metallicity calculation

- NIRSpec + MRS
- Metallicity of $0.1Z_{\odot}$
- Mass-metallicity relation

High resolution NIRSpec separate [OII] doublet

- High densities ($\sim 1000\text{cm}^{-3}$)



Hsiao, Álvarez-Márquez+24; Abdurro'f+24



MIRI/LRS observations of GHZ2 at $z=12.3$




Most distant galaxy with direct detection of [OIII] and $H\alpha$

Metallicity estimation from a non-direct method

High electron densities

Non-spectrally resolved due to the low resolution of the LRS

- Needs MRS observations for higher SNR and access to kinematics








PRISMS. PRIMordial galaxy Survey with MIRI-LRS Spectroscopy at $z \sim 10$

JWST Cycle 4 program:

- PIs: J. Álvarez-Márquez & L. Colina
- Total of 129 hours
- Observed between 2025 to 2026
- MIRI low resolution spectroscopy
- Survey of 10 galaxies at $z \sim 10$
- Detection of $H\beta + [OIII]$ & $H\alpha$



First statistical study of the ISM properties (SFR, dust, metallicity, ionization, ...) of galaxies 500 Myr after the Big Bang.







JWST & Early Universe: Redshift frontier

JWST &
ALMA

$z=14.4$




$z=14.3$




$z=13.0$



$z=12.3$



$z=10.6$



$z=10.2$



Billions of years ago

13.7

13.5

13.4

13

HUBBLE
2012

HUBBLE
2009

Current JWST + ALMA limits

Big Bang

Neutral
hydrogen
and
helium

Cosmic
'dark ages'

Reionization

HST + ALMA limits

Recombination

First stars


First galaxies

HST &
ALMA


$z=11$




$z=9.1$



$z=8.6$



$z=7.7$



$z=7.1$

