



PROBING THE HALOS OF PRIMORDIAL GALAXIES


Sumukha R Bharadwaj

PhD. Scholar

Indian Institute of Space Science and Technology, Thiruvananthapuram



WHY STUDY THE CIRCUMGALACTIC MEDIUM?




- Baryon Reservoir
- Galaxy Fuelling
- Feedback Engine
- Metal Repository
- Angular Momentum Transfer
- Evolutionary Link

Credits: Jason Tumlinson et al. (2017)



PROBING DIFFUSE UNIVERSE



Absorber

QSO




a

Observer


Redshift

Background
quasar




Metal-bearing
galaxy halos

b



Credit:
Péroux & Howk (2020)

Metal enrichment and evolution in four $z > 6.5$ quasar sightlines observed with JWST/NIRSpec

L. Christensen^{1,2} , P. Jakobsen^{1,2}, C. Willott³, S. Arribas⁴, A. Bunker⁵, S. Charlot⁶, R. Maiolino^{7,8,9},
M. Marshall³, M. Perna⁴, and H. Übler^{7,8}

¹ Cosmic Dawn Center (DAWN), University of Copenhagen, 2200 Copenhagen N, Denmark
e-mail: lichrist@nbi.ku.dk

² Niels Bohr Institute, University of Copenhagen, Jagtvej 128, 2200 Copenhagen N, Denmark

³ National Research Council of Canada, Herzberg Astronomy & Astrophysics Research Centre, 5071 West Saanich Road,
Victoria BC V9E 2E7, Canada

⁴ Centro de Astrobiología (CAB), CSIC-INTA, Ctra. de Ajalvir km 4, Torrejón de Ardoz, 28850 Madrid, Spain

⁵ Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

⁶ Sorbonne Université, CNRS, UMR 7095, Institut d'Astrophysique de Paris, 98 bis bd Arago, 75014 Paris, France

⁷ Kavli Institute for Cosmology, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

⁸ Cavendish Laboratory, University of Cambridge, 19 JJ Thomson Avenue, Cambridge CB3 0HE, UK

⁹ Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK

Received 11 September 2023 / Accepted 11 October 2023

ABSTRACT


We present JWST/NIRSpec $R \approx 2700$ spectra of four high-redshift quasars: VDES J0020–3653 ($z = 6.860$), DELS J0411–0907 ($z = 6.825$), UHS J0439+1634 ($z = 6.519$), and ULAS J1342+0928 ($z = 7.535$). The exquisite data quality, signal-to-noise ratio of 50–200, and large $0.86 \mu\text{m} \leq \lambda \leq 5.5 \mu\text{m}$ spectral coverage allowed us to identify between 13 and 17 intervening and proximate metal absorption line systems in each quasar spectrum, with a total number of 61 absorption-line systems detected at $2.42 < z < 7.48$ including the highest redshift intervening O I $\lambda 1302$ and Mg II systems at $z = 7.37$ and $z = 7.44$. We investigated the evolution of the metal enrichment in the epoch of re-ionisation (EoR) at $z > 6$ and found the following: *i*) a continued increase in the low-ionisation O I, C II, and Si II incidence, *ii*) decreasing high-ionisation C IV and Si IV incidence with a transition from predominantly high- to low-ionisation at $z \approx 6.0$, and *iii*) a constant Mg II incidence across all redshifts. The observations support a change in the ionisation state of the intergalactic medium in the EoR rather than a change in metallicity. The abundance ratio of [Si/O] in five $z > 6$ absorption systems show enrichment signatures produced by low-mass Pop III pair instability supernovae, and possibly Pop III hypernovae. In the Gunn-Peterson troughs, we detected transmission spikes where Ly α photons can escape. From 22 intervening absorption line systems at $z > 5.7$, only a single low-ionisation system out of 13 lies within 2000 km s^{-1} from a spike, while four high-ionisation systems out of nine lie within $\sim 2000 \text{ km s}^{-1}$ from a spike. Most spikes do not have associated metal absorbers close by. This confirms that star-forming galaxies responsible for producing the heavy elements that are transported to the circumgalactic medium via galaxy winds do so in predominantly high-density, neutral environments, while lower density environments are ionised without being polluted by metals at $z \approx 6$ –7.

Key words. cosmology: observations – intergalactic medium – galaxies: high-redshift – quasars: absorption lines – dark ages, reionization, first stars

Christensen et al. (2023), A&A 680, A82



JWST NIRSPEC/SLIT SPECTRUM OF VDES J0020-3653 ($z = 6.855$)



Filters:

F070LP, G140H

F170LP, G235H



Mg II $\lambda\lambda$ 2796, 2803 ABSORPTION LINE SYSTEMS


z_{absorber}	$W_r 2796$	$W_r 2803$
6.4535	0.4590 ± 0.0025	0.2545 ± 0.0027
6.5028	0.2400 ± 0.0026	0.3498 ± 0.0024
6.5625	0.3278 ± 0.0024	0.2862 ± 0.0034
6.6690	0.1855 ± 0.0024	0.1215 ± 0.0023

$$W_r = \int_{\lambda_1}^{\lambda_2} \left(1 - \frac{F_\lambda}{F_c} \right) d\lambda$$

$$\tau_a(v) = \ln \left(\frac{F_c(v)}{F(v)} \right)$$


$$N_a(v) = 3.768 \times 10^{14} \times (f\lambda)^{-1} \times \tau_a(v)$$

... Savage & Sembach(1991)





JWST NIRSPEC/IFU SPECTRUM OF VDES J0020-3653 ($z = 6.855$)



Filters: **F290LP, G395H**



Na I $\lambda\lambda$ 5890, 5895 ABSORPTION LINE SYSTEMS


z_{absorber}	W_r
6.4535	--
6.5028	--
6.5625	--
6.6690	--

$$W_r = \int_{\lambda_1}^{\lambda_2} \left(1 - \frac{F_\lambda}{F_c} \right) d\lambda$$

$$\tau_a(v) = \ln \left(\frac{F_c(v)}{F(v)} \right)$$

$$N_a(v) = 3.768 \times 10^{14} \times (f\lambda)^{-1} \times \tau_a(v)$$

... Savage & Sembach(1991)





Ca II $\lambda\lambda$ 3934, 3969 ABSORPTION LINE SYSTEMS


z_{absorber}	$W_r 3934$	$W_r 3969$
6.4535	< 0.5625	< 0.6446
6.5028	< 0.6466	< 1.2251
6.5625	< 0.4278	< 0.3676
6.6690	< 0.3204	< 0.3027

$$W_r = \int_{\lambda_1}^{\lambda_2} \left(1 - \frac{F_\lambda}{F_c} \right) d\lambda$$

$$\tau_a(v) = \ln \left(\frac{F_c(v)}{F(v)} \right)$$

$$N_a(v) = 3.768 \times 10^{14} \times (f\lambda)^{-1} \times \tau_a(v)$$

... Savage & Sembach(1991)



Thank You