

Spectral Emission Characteristics of NGC 7469



A Hands-on Workshop with JWST and
UVIT



Ruchika Dhaka

PhD Scholar


Department of Physics, IIT Kanpur



Introduction of AGNs


Active Galactic Nuclei (AGNs) are the extremely luminous central regions of some galaxies, powered by **accretion of matter onto a supermassive black hole (SMBH)** at their cores.

Seyfert galaxy NGC 7469



Credit: Adapted from Javier's talk

redshift=0.01627




Strong emission across radio to X-rays Prominent

NGC 7469 with Medium Resolution Spectrograph (MRS)

Total number of Channels and Bands:

Channels: 1, 2, 3, 4

Bands: SHORT, MEDIUM, LONG




Two array detectors of 1024×1024 pixels FASTR1 & SLOWR1 readouts mode


FoV: 3.7"×3.7" (Ch1) – 7.4"×7.9" (Ch4)

Slice width: 0.18" (Ch-1) – 0.64" (Ch-4)


Spectral resolution: $R \approx 1500\text{--}3500$



Characterisation of line profiles of high-excitation and coronal emission lines in the nuclear spectrum of NGC 7469



Emission line maps in the central region of NGC 6552 generated using the MRS channels




Strong high-ionization lines ([Ne V], [Ne VI], [O IV]) indicate **AGN dominance** with some contribution from circumnuclear star formation.

Mid-IR Line Ratios and Seyfert Classification

Comparison based on Pereira-Santaella et al. (2010, A&A 517, A23).

Ratio	NGC 7469	Seyfert 1 median	Seyfert 2 median
[Ne III]/[Ne II]	1.44	≈ 1.1	≈ 0.9
[Ne V]/[Ne II]	0.87	≈ 0.9	≈ 0.7
[Ne V]/[Ne III]	0.60	≈ 0.5	≈ 0.4
[O IV]/[Ne II]	1.52	≈ 1.5	≈ 1.2
[O IV]/[Ne III]	1.06	≈ 1.0	≈ 0.8



Black Hole Mass Estimate — NGC 7469

- Based on [Dasyra et al. \(2008, ApJ 674, L9\)](#) relation using the **[Ne V] 14.32 μm** line width.
- The black hole (BH) mass based on the line FWHM of the high-excitation emission line


$$\log(M_{\text{BH}} / M_{\odot}) = 7.86 + 4.31 \log(\sigma / 200 \text{ km s}^{-1})$$

Results (from [Ne V] narrow component):






- FWHM = **278.2 km s^{-1}**

Estimated black hole mass: $M_{\text{BH}} \approx 8 \times 10^6 M_{\odot}$

- The derived mass agrees with previous reverberation-based estimates ($\sim 10^7 M_{\odot}$). ([Peterson et al. \(2014\)](#))



Highly ionised nuclear outflow



Line	FWHM (km/s)	Velocity offset (km/s)	σ (km/s)	Outflow Velocity (km/s)
[Ne V] 14.32 μm	865.0	351.2	368.1	1087
[O IV] 25.89 μm	659.9	356.0	280.8	918
[Ne VI] 7.65 μm	659.4	195.6	280.6	757
[Ne III] 15.56 μm	819.7	301.3	348.8	999

Outflow maximum velocity:

$$V_{\text{peak}} + 2 \times \sigma_{\text{blue}},$$

Ref: Javier + 2023, A&A

- The **blue-shifted velocity components** identified in all atomic emission lines are interpreted as being due to the **presence of outflowing material close to the AGN**.
- High-velocity outflow with offset velocities of **+195–350 km s⁻¹**.
- All four lines show **high outflow velocities (~750–1100 km s⁻¹)** — typical of **AGN-driven ionized winds**.
- evidence for a highly ionised, AGN-powered, and fast outflowing gas.

Thanks for your attention