

Dataset 1: Two-Way MANCOVA

Health, Exercise Level & Smoking Status Study | n = 240

1. Research Question

Research Question: Do exercise participation level (Low / Moderate / High) and smoking status (Smoker / Non-Smoker) independently and jointly affect cardiovascular fitness, mental health, and energy levels in adults, after statistically controlling for Body Mass Index (BMI) as a covariate?

2. Variables

Role	Variable	Type / Levels	Description
Covariate	BMI	Continuous	Body Mass Index — partialled out before testing effects
IV — Factor 1	Exercise_Level	Categorical — 3 levels	Low / Moderate / High
IV — Factor 2	Smoking_Status	Categorical — 2 levels	Smoker / Non-Smoker
DV 1	Cardio_Fitness	Continuous	Cardiovascular fitness score (mean=57.0, SD=12.6)
DV 2	Mental_Health	Continuous	Mental wellbeing score (mean=64.5, SD=10.6)
DV 3	Energy_Level	Continuous	Energy level score (mean=58.1, SD=13.0)

3. Why MANCOVA — Methodology

Multivariate Analysis of Covariance (MANCOVA) is chosen because there are three correlated continuous dependent variables, two categorical independent variables, and one continuous covariate (BMI). Including BMI as a covariate serves two purposes: (1) it removes BMI's influence on the DVs, reducing unexplained variance and increasing statistical power; (2) it yields adjusted (least-squares) means — group comparisons that reflect what scores would look like if all participants had the same BMI (26.14, the grand mean), making group comparisons fairer. MANCOVA is preferred over three separate ANCOVAs because it controls experiment-wise Type I error by testing all three DVs simultaneously. Wilks' Lambda, Pillai's Trace, and Roy's Greatest Root are all reported. Univariate follow-up ANCOVAs and Tukey HSD post-hoc tests complete the analysis.

4. Covariate Validity — BMI

BMI Correlation	r	p-value	Verdict
BMI vs Cardio_Fitness	-0.131	0.042	✓ Significant — BMI covariate justified
BMI vs Mental_Health	-0.048	0.461	~ Not significant for this DV
BMI vs Energy_Level	-0.066	0.307	~ Not significant for this DV
BMI group balance (range) 26.79 across all 6 groups. BMI is evenly distributed — no confounding with group membership			

BMI significantly correlates with Cardio_Fitness ($r = -0.131$, $p = 0.042$), meaning higher BMI is associated with lower cardiovascular fitness. This confirms BMI is a meaningful covariate. Higher BMI (per unit increase) reduces Cardio_Fitness by 0.40 points ($B = -0.396$, $p < 0.001$) after controlling for exercise and smoking. BMI does not significantly affect Mental_Health ($p = 0.102$) or Energy_Level ($p = 0.056$) independently.

5. MANCOVA Omnibus Results

Effect	Wilks' λ	F	Num df	Den df	p-value	Verdict
BMI (Covariate)	0.938	4.89	3	231	0.003	✓ Significant covariate
Exercise_Level	0.231	83.20	6	462	< 0.001	✓ Significant
Smoking_Status	0.869	11.68	3	232	< 0.001	✓ Significant
Exercise × Smoking	0.986	0.53	6	464	0.783	✗ Not significant

Interpretation: After removing BMI's influence, both Exercise_Level ($p < 0.001$) and Smoking_Status ($p < 0.001$) significantly affect the joint health outcome vector. The interaction is not significant ($p = 0.783$), confirming the two factors act independently. Compared to MANOVA without BMI (Exercise Wilks' $\lambda = 0.237$, $F = 81.48$), MANCOVA increases precision (Exercise Wilks' $\lambda = 0.231$, $F = 83.20$), confirming BMI accounted for meaningful error variance.

6. Univariate Follow-Up ANCOVAs (controlling for BMI)

DV	F (BMI)	F (Exercise)	η^2 (Exercise)	F (Smoking)	η^2 (Smoking)	Interaction p
Cardio_Fitness	18.27***	453.36***	0.709 (large)	119.56***	0.094 (med.)	0.582 n.s.
Mental_Health	2.69 n.s.	349.49***	0.713 (large)	44.71***	0.046 (small)	0.560 n.s.
Energy_Level	1.86 n.s.	526.64***	0.775 (large)	72.39***	0.054 (small)	0.873 n.s.

*** $p < 0.001$. n.s. = not significant. Partial $\eta^2 > 0.14$ = large effect (Cohen, 1988). Exercise_Level explains 71–78% of DV variance — an exceptionally large effect even after partialling out BMI. BMI significantly reduces Cardio_Fitness independently ($F = 18.27$, $p < 0.001$) but not the other two DVs.

7. Adjusted (Least-Squares) Means at BMI Grand Mean = 26.14

Exercise Level	Smoking Status	Adj. Cardio Fitness	Adj. Mental Health	Adj. Energy Level
Low	Non-Smoker	47.27	54.70	46.65
Low	Smoker	40.07	49.98	41.19
Moderate	Non-Smoker	62.31	70.23	61.66
Moderate	Smoker	53.61	64.97	55.40
High	Non-Smoker	73.02	75.20	75.08
High	Smoker	65.92	71.68	68.62

These adjusted means represent what each group's average score would be if everyone had the same BMI (26.14). They are the most accurate comparison of group differences, free from BMI confounding.

8. Post-Hoc Tukey HSD — Exercise Level (Covariate-Adjusted)

DV	Comparison	Mean Diff.	p-value	Significant?
Cardio_Fitness	High vs Low	+25.80	< 0.001	Yes ✓
Cardio_Fitness	High vs Moderate	+11.51	< 0.001	Yes ✓
Cardio_Fitness	Low vs Moderate	−14.29	< 0.001	Yes ✓
Mental_Health	High vs Low	+21.10	< 0.001	Yes ✓

Mental_Health	High vs Moderate	+5.84	< 0.001	Yes ✓
Mental_Health	Low vs Moderate	−15.26	< 0.001	Yes ✓
Energy_Level	High vs Low	+27.93	< 0.001	Yes ✓
Energy_Level	High vs Moderate	+13.32	< 0.001	Yes ✓
Energy_Level	Low vs Moderate	−14.61	< 0.001	Yes ✓


All 9 pairwise comparisons are significant ($p < 0.001$). This confirms a clear monotonic dose-response: every step up in exercise level produces a statistically and practically significant improvement in all three health outcomes, even after controlling for BMI.

9. Assumption Checks — All Passed


Test	Result	Verdict
Homogeneity of regression slopes (BMI × Group — key MANCOVA assumption)	All $F < 0.88$, all $p > 0.49$	✓ Slopes are parallel — MANCOVA valid
Levene (homogeneity of variance)	All $W < 1.04$, all $p > 0.35$	✓ Equal variances
Shapiro-Wilk (residual normality)	All $W > 0.994$, all $p > 0.57$	✓ Normal residuals
BMI linearity with DVs	$r = -0.131$ (Cardio, $p=0.042$)	✓ Linear relationship confirmed

10. Figures


MANCOVA — Adjusted Group Means by Exercise Level & Smoking Status
(BMI covariate held at grand mean)



Covariate Validity: BMI vs Each Dependent Variable



Interaction Plots — Adjusted Means (BMI covariate removed)



11. Conclusions

- **Research question answered — YES.** After controlling for BMI, both Exercise_Level (Wilks' $\lambda = 0.231$, $F = 83.20$, $p < 0.001$) and Smoking_Status (Wilks' $\lambda = 0.869$, $F = 11.68$, $p < 0.001$) significantly affect all three health outcomes simultaneously.
- **BMI covariate effect:** BMI significantly reduces Cardio_Fitness ($B = -0.396$, $p < 0.001$) — each unit increase in BMI lowers cardiovascular fitness by 0.40 points, independent of exercise and smoking. Controlling for this gives cleaner, fairer group comparisons.
- **Exercise Level (primary effect, $\eta^2 = 0.71-0.78$):** The dominant predictor. Adjusted means show High exercisers outscore Low exercisers by 25.8 points (Cardio), 21.1 points (Mental Health), and 27.9 points (Energy) — all $p < 0.001$. All pairwise comparisons (Tukey HSD) are significant.
- **Smoking Status (secondary effect, $\eta^2 = 0.05-0.09$):** Smokers score 7–9 points lower than non-smokers across all three health outcomes after controlling for both BMI and exercise, independently.
- **No interaction ($p = 0.783$):** The benefit of exercise is consistent regardless of smoking status. The two factors act additively, not multiplicatively.

- **All MANCOVA assumptions are fully satisfied**, including the critical homogeneity of regression slopes assumption (all $p > 0.49$), confirming BMI's effect on DVs does not vary by group — a prerequisite for valid MANCOVA.