

Glossary of Metrics for the Maine Aquatic Barrier Prioritization Tool

Published in April 2026

List of fish species in metrics

Common Name	Scientific Name
Alewife	<i>Alosa pseudoharengus</i>
American shad	<i>Alosa sapidissima</i>
Atlantic salmon	<i>Salmo salar</i>
Blueback herring	<i>Alosa aestivalis</i>
Eastern Brook trout	<i>Salvelinus fontinalis</i>
Rainbow smelt	<i>Osmerus mordax</i>

List of entities cited in metrics

Entity Name	Entity Acronym
Maine Department of Inland Fisheries and Wildlife	IFW
Maine Department of Marine Resources	DMR
Maine Geological Survey	MGS
Multi-Resolution Land Characteristics Consortium	MRLC Consortium
National Land Cover Database	NLCD
The Nature Conservancy, Maine	TNC Maine
National Oceanic and Atmospheric Administration	NOAA
U.S. Department of Agriculture – Natural Resources Conservation Service	USDA – NRCS
U.S. Fish and Wildlife Service	USFWS
United State Geologic Survey	USGS

Table of Contents

[Metrics displayed in prioritization tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
Passability	Passability	About the Crossing	N/A	13
BatFuncUS	Upstream Functional Network Length	Network	Coastal Anadromous, Inland Brook Trout, Atlantic Salmon, American shad/Blueback herring	14
batTotUSDS	Total Functional Network Length	Network	Inland Brook Trout	15
batLenUS	Total Upstream Network Length	Network	N/A	16
batCountDS	Count of Downstream Barriers	Network	Atlantic Salmon, Alewife, American shad/Blueback herring	17
DSHydro	Count of downstream hydropower facilities	Network	American shad/Blueback herring	18
DSFishways	Count of downstream fish passage facilities	Network	Atlantic salmon, Alewife, American shad/Blueback herring	19
batAbs	Absolute Gain	Network	Inland Brook Trout	20
dsPassabiliy Product	Downstream Passability Product	Network	Coastal Anadromous, Atlantic Salmon, Alewife, American shad/Blueback herring	21
Tidal	Barrier is located on a tidal stream (current HAT +1m)	Landscape	NA	22
CoastalStreams	Barrier is located on a Coastal Stream	Landscape	NA	23
onConsLand	Barrier is on conservation land	Landscape	NA	24

Table of Contents

[Metrics displayed in prioritization tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
AquiferOr CoarseSed	Aquifer OR coarse sediments in upstream or downstream functional networks or both	Landscape	Coastal Anadromous, Inland brook trout	25
CalcModCalc	Calcareous/moderately calcareous geology in upstream or downstream networks or both	Landscape	Coastal Anadromous, Inland brook trout	26
DA_PercFor	Percent forested landcover in upstream drainage area	Landscape	Coastal Anadromous, Inland Brook Trout, Atlantic salmon, Alewife, American shad/Blueback herring	27
buff100_Tot_Perc_Statewide_forestMerged	Percent forest cover in 100m buffer of total functional network	Landscape	Coastal Anadromous, Inland Brook Trout	28
buff100_US_Perc_Statewide_forestMerged	Percent forest cover in 100m buffer of upstream network	Landscape	NA	29
usAlewifeAcres	Upstream acres of alewife ponds	Fish Habitat - Alewife	Alewife	30
occ_temp7p20	Probability of brook trout occupancy (+2degC)	Fish Habitat - Inland Brook Trout (BKT)	Inland Brook Trout	31
HeritagePondBarrier	Heritage Fish Pond Barrier	Fish Habitat - Inland Brook Trout (BKT)	Inland Brook Trout	32
MilesBT_HVH_Usonly	Number of miles of (high, very high) brook trout habitat in US network	Fish Habitat - Inland Brook Trout (BKT)	Coastal Anadromous, Inland Brook Trout	33

Table of Contents

[Metrics displayed in prioritization tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
MilesBT_HVH	Number of miles of (high, very high) brook trout habitat in US + DS network	Fish Habitat - Inland Brook Trout (BKT)	Coastal Anadromous, Inland Brook Trout	34
EBTJV_Patch	EBTJV Wild Brook Trout Patch	Fish Habitat - Inland Brook Trout (BKT)	NA	35
bkt_p20_occ_gt50	Probability of brook trout occupancy >=50% (+2degC)	Fish Habitat - Inland Brook Trout (BKT)	Inland Brook Trout	36
SeaRunBT	Barrier to Sea-Run Brook Trout	Fish Habitat - Inland Brook Trout (BKT)	Coastal Anadromous	37
HUC10_CritSalmHab	Barrier is in a Salmon Critical Habitat HUC10	Fish Habitat - Atlantic Salmon	Atlantic Salmon	38
SalmonHabQual	Atlantic Salmon Habitat Quality	Fish Habitat - Atlantic Salmon	Atlantic Salmon	39
US_SalmHabUnits	Summed Salmon Habitat Units in US Network	Fish Habitat - Atlantic Salmon	Atlantic Salmon	40
HUC10_Habit_Units	Salmon Habitat Units in HUC10	Fish Habitat - Atlantic Salmon	NA	41
cumulative SalmHabUnits	Cumulative Upstream Salmon Habitat Units	Fish Habitat - Atlantic Salmon	Atlantic Salmon	42
usSmeltSite	Smelt spawning sites upstream of barrier	Fish Habitat - Smelt	Coastal Anadromous	43
InvasiveBarrier	Species of Management Concern on one side of a barrier and not the other	Invasive Species	NA	44

Table of Contents

[Metrics calculated, but not displayed in tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
batFuncDS	Downstream Functional Network Length	Network	NA	46
cumulativeUSNet	Cumulative Upstream Functional Network	Network	NA	47
batCountUS	Upstream Barrier Count	Network	NA	48
batUSDnsty	Density of barriers on upstream network	Network	NA	49
DSFalls	Number of natural barriers on downstream flowpath	Network	NA	50
usNatBar	One or more of the next upstream barriers is a natural barrier	Network	NA	51
usRestPot	Upstream Restoration Potential	Network	NA	52
Aquifer	Aquifers in upstream or downstream functional networks or both	Landscape	NA	53
usCons100	Percent conserved land in 100m buffer of upstream functional network	Landscape	NA	54
dsCons100	Percent conserved land in 100m buffer of downstream functional network	Landscape	NA	55
buff100_DS_Perc_Statewide_forest Merged	Percent forest cover in 100m buffer of downstream network	Landscape	NA	56
buff100_US_Perc_nlcd_2016_developedMedHigh	Percent medium/high intensity developed landcover in 100m buffer of upstream network	Landscape	NA	57

Table of Contents

[Metrics calculated, but not displayed in tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
buff100_DS_Perc_nlcd_2016_developedMedHigh	Percent medium/high intensity developed landcover in 100m buffer of downstream network	Landscape	NA	58
buff100_Tot_Perc_nlcd_2016_developedMedHigh	Percent medium/high intensity developed landcover in 100m buffer of total functional network	Landscape	NA	59
buff100_US_Perc_nlcd_2016_cultivatedCrops	Percent cultivated crops in 100m buffer of upstream network	Landscape	NA	60
buff100_DS_Perc_nlcd_2016_cultivatedCrops	Percent cultivated crops in 100m buffer of downstream network	Landscape	NA	61
buff100_Tot_Perc_nlcd_2016_cultivatedCrops	Percent cultivated crops in 100m buffer of total functional network	Landscape	NA	62
buff100_US_Perc_nlcd_2016_pastureHay	Percent pasture/hay in 100m buffer of upstream network	Landscape	NA	63
buff100_DS_Perc_nlcd_2016_pastureHay	Percent pasture/hay in 100m buffer of downstream network	Landscape	NA	64
buff100_Tot_Perc_nlcd_2016_pastureHay	Percent pasture/hay in 100m buffer of total functional network	Landscape	NA	65

Table of Contents

[Metrics calculated, but not displayed in tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
floodplain_US_Perc_Statewide_forestMerged	Percent forest cover in the floodplain of upstream network	Landscape	NA	66
floodplain_DS_Perc_Statewide_forestMerged	Percent forest cover in the floodplain of downstream network	Landscape	NA	67
floodplain_Tot_Perc_Statewide_forestMerged	Percent forest cover in the floodplain of total functional network	Landscape	NA	68
floodplain_US_Perc_nlcd_2016_developedMedHigh	Percent medium/high intensity developed landcover in the floodplain of the upstream network	Landscape	NA	69
floodplain_DS_Perc_nlcd_2016_developedMedHigh	Percent medium/high intensity developed landcover in the floodplain of the downstream network	Landscape	NA	70
floodplain_Tot_Perc_nlcd_2016_developedMedHigh	Percent medium/high intensity developed landcover in the floodplain of the total functional network	Landscape	NA	71
floodplain_US_Perc_nlcd_2016_cultivatedCrops	Percent cultivated crops in the floodplain of upstream network	Landscape	NA	72
floodplain_DS_Perc_nlcd_2016_cultivatedCrops	Percent cultivated crops in the floodplain of downstream network	Landscape	NA	73

Table of Contents

[Metrics calculated, but not displayed in tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
floodplain_Tot_Perc_nlcd_2016_cultivatedCrops	Percent cultivated crops in the floodplain of total functional network	Landscape	NA	74
floodplain_US_Perc_nlcd_2016_pastureHay	Percent pasture/hay in the floodplain of upstream network	Landscape	NA	75
floodplain_DS_Perc_nlcd_2016_pastureHay	Percent pasture/hay in the floodplain of downstream network	Landscape	NA	76
floodplain_Tot_Perc_nlcd_2016_pastureHay	Percent pasture/hay in the floodplain of total functional network	Landscape	NA	77
NESZCL_Int	River Size Class	Landscape	NA	78
MilesUS_HATcurrent	Miles of tidal habitat (current) upstream of barrier	Landscape/Fish Habitat	NA	79
MilesUS_HATPlus1m	Miles of tidal habitat (current HAT + 1m) upstream of barrier	Landscape/Fish Habitat	NA	80
MilesUS_HATPlus6ft	Miles of tidal habitat (current HAT + 6ft) upstream of barrier	Landscape/Fish Habitat	NA	81

Table of Contents

[Metrics calculated, but not displayed in tool]

GIS Name	Full Name	Category in Tool	Prioritization Scenarios	Slide Num.
bkt_curr_occ_gt50	Probability of brook trout occupancy >=50% (current temps)	Fish Habitat - Inland Brook Trout (BKT)	NA	82
BT_1_4_HVH	High or Very High quality brook trout habitat upstream OR downstream of barrier	Fish Habitat - Inland Brook Trout (BKT)	Inland Brook Trout	83
BT_1_4_VH	Very High quality brook trout habitat upstream OR downstream of barrier	Fish Habitat - Inland Brook Trout (BKT)	NA	84
occ_current	Probability of brook trout occupancy (current temps)	Fish Habitat - Inland Brook Trout (BKT)	NA	85
occ_temp7p40	Probability of brook trout occupancy (+4degC)	Fish Habitat - Inland Brook Trout (BKT)	NA	86
bkt_p40_occ_gt50	Probability of brook trout occupancy >=50% (+4degC)	Fish Habitat - Inland Brook Trout (BKT)	NA	87
US_SalmParrProd	Summed salmon parr productivity in US Functional Network	Fish Habitat - Atlantic Salmon	NA	88
InvasiveBarrierConfirmed	Species of Management Concern on one side of a barrier and not the other - Confirmed only	Invasive Species	NA	89
InvasiveBarrierConfNumSpp	Number of Species of Management Concern that are blocked - Confirmed only	Invasive Species	NA	90

Metrics displayed in the Aquatic Barrier Prioritization Tool

Passability

GIS Field Name: Passability

Category: About the Crossing

A numerical score from 0-1 that indicates how passable a barrier is likely to be, where 0=complete barrier and 1=fully passable. All dams are considered complete barriers, unless there is known fish passage facilities at a dam. Dams with fish passage were reviewed by J. Royte and E. Martin from The Nature Conservancy and assigned a passability score based on the efficacy of passage at the dam. Passability for road-stream crossings is first assigned based on Maine's crossing assessment protocol which splits crossings into barriers, potential barriers, and no barrier. Barriers were assigned a value of 0 and no barriers were assigned a value of 1.

Potential barriers were further refined based on Froude number which is a proxy for the relationship between velocity and depth that can be utilized as a screening metric to identify crossings that have a high probability of being a velocity barrier to fish passage. Passability score of 0.66 correlates directly to a Froude number of 1, so only potential barriers with a score of < 0.66 were prioritized in the final result.

This metric is used in the generation of other metrics.

Unit: Index

Upstream Functional Network Length

GIS Field Name: batFuncUS

Category: Network

Used in Prioritization Scenario(s): Coastal Anadromous, Inland Brook Trout, Atlantic salmon, American shad/Blueback herring

Length of the functional network upstream of a barrier.


“Functional network” is defined by those sections of river that a fish could theoretically access from any other point within that functional network. Terminal ends are barriers, headwaters, and/or river mouth.

Unit: meters (converted to miles in some outputs)

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine



Total Functional Network Length

GIS Field Name: batTotUSDS

Category: Network

Used in Prioritization Scenario(s): Inland Brook Trout


Length of the functional network downstream of a barrier plus the length of the functional network upstream of a barrier. This is the length of the recombined network if the barrier was removed.

Unit: meters (converted to miles in some outputs)

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine



Total Upstream Network Length

GIS Field Name: batLenUS

Category: Network


Total length of river network upstream of a given barrier, regardless of any upstream barriers.

Unit: meters

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine



Count of Downstream Barriers

GIS Field Name: batCountDS

Category: Network

Used in Prioritization Scenario(s): Atlantic salmon, Alewife, American shad/Blueback herring

The number of barriers downstream of a given barrier. Includes natural barriers (waterfalls), which are included in network dataset creation

Does not include barriers excluded from network dataset creation

Unit: #

Sort Order: Ascending

Data Sources:

- Hydrography (2023): USGS, TNC Maine

Count of downstream hydropower facilities

GIS Field Name: DSHydro

Category: Network

Used in Prioritization Scenario(s): American shad/Blueback herring

The number of hydropower facilities downstream of a barrier

Unit: #

Sort Order: Ascending

Data Sources:

- Hydrography (2023): USGS, TNC Maine
- Dam information (2025): TNC Maine compiled

Count of downstream fish passage facilities

GIS Field Name: DSFishways

Category: Network

Used in Prioritization Scenario(s): Atlantic salmon, Alewife, American shad/Blueback herring

The number of fish passage facilities downstream of a barrier

Unit: #

Sort Order: Descending

Data Sources:

- Hydrography (2023), USGS, TNC Maine
- Fishway info: DMR (2024)

Absolute Gain

GIS Field Name: batAbs

Category: Network

Used in Prioritization Scenario(s): Inland Brook Trout

This metric is the minimum of the two functional networks of a barrier. For example, if the upstream functional network is 10 kilometers and downstream functional network is 5 kilometers, then the Absolute Gain is 5 kilometers.

Unit: meters

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine

Downstream Passability Product

GIS Field Name: dsPassabilityProduct

Category: Network

Used in Prioritization Scenario(s): Coastal Anadromous, Atlantic salmon, Alewife, American shad/Blueback herring

The product (multiplication) of all passability scores on the downstream flow path of a barrier.

Unit: Index

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine

Barrier is located on a tidal stream (current HAT +1m)

GIS Field Name: Tidal

Category: Landscape

Barrier is located on a tidal stream, based on the spatial intersection of the barrier and the Highest Annual Tide (HAT2018) plus 1m dataset:

Unit: Boolean

Sort Order: Descending

Data Sources:

- Tidal Waters / Sea-level Rise Scenarios (2018): MGS

Barrier is Located on a Coastal Stream

GIS Field Name: CoastalStreams

Category: Landscape

Barrier is located on a small coastal stream – Small Coastal streams were defined as streams with an upstream drainage area less than 100 km² that drain into a tidal area (directly into the ocean or the tidal portion of a larger river).

Unit: Boolean

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine
- Tidal Waters / Sea-level Rise Scenarios (2018): MGS

Barrier is on conservation land

GIS Name: onConsLand

Category: Landscape

Conservation Lands included in the current analysis included those with gap status 1,2,3, and 39 – based on TNC's modified version of USGS's [Gap Status Codes](#).

Unit: Boolean

Data Sources:

- Conservation Lands / Secured Lands (2025): TNC Maine

Aquifer OR coarse sediments in upstream or downstream functional networks or both

GIS Field Name: AquiferOrCoarseSed

Category: Landscape

Used in Prioritization Scenario(s): Coastal Anadromous, Inland Brook Trout

[Aquifers](#) OR coarse sediments are found in [upstream functional network](#) OR [downstream functional network](#) OR both. This metric evaluates the potential for cold water habitats within a barrier's respective networks.

Unit: Boolean

Sort Order: Descending

Data Sources:

- Aquifers (2020): MGS
- Coarse Sediments (2006): The Nature Conservancy Ecological Land Units

Calcareous / moderately calcareous geology in upstream or downstream networks or both

GIS Field Name: CalcModCalc

Category: Landscape

Used in Prioritization Scenario(s): Coastal Anadromous, Inland Brook Trout

Calcareous or moderately calcareous geology is found in [upstream functional network](#) OR [downstream functional network](#) OR both. This metric evaluates the acid buffering capacity within a barrier's respective networks.

Unit: Boolean

Sort Order: Descending

Data Sources:

- Calcareous/Moderately Calcareous Bedrock (2006): The Nature Conservancy Ecological Land Units

Percent forested landcover in upstream drainage area

GIS Name: DA_PercFor

Category: Landscape

Used in Prioritization Scenario(s): Coastal Anadromous, Inland Brook Trout, Atlantic salmon, Alewife, American shad/Blueback herring

Percent forested landcover in entire upstream watershed.

Forested landcover classes were aggregated separately for northern Maine and southern Maine. In northern Maine, forest cover was defined as deciduous forest, evergreen forest, mixed forest, and woody wetlands. In southern Maine, scrub/shrub was also included as a forest class based on input from Merry Gallagher, Maine DIFW.

Unit: %

Sort Order: Descending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent forest cover in 100m buffer of total functional network

GIS Name: buff100_Tot_Perc_Statewide_forestMerged

Category: Landscape

Used in Prioritization Scenario(s): Coastal Anadromous, Inland Brook Trout

Percent forested landcover in a 100m buffer of a barrier's total functional network (upstream + downstream).

Forested landcover classes were aggregated separately for northern Maine and southern Maine. In northern Maine, forest cover was defined as deciduous forest, evergreen forest, mixed forest, and woody wetlands. In southern Maine, scrub/shrub was also included as a forest class based on input from Merry Gallagher, Maine IFW.

Unit: %

Sort Order: Descending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent forest cover in 100m buffer of upstream network

GIS Name: buff100_US_Perc_Statewide_forestMerged

Category: Landscape

Percent forested landcover in a 100m buffer of a barrier's upstream functional network.

Forested landcover classes were aggregated separately for northern Maine and southern Maine. In northern Maine, forest cover was defined as deciduous forest, evergreen forest, mixed forest, and woody wetlands. In southern Maine, scrub/shrub was also included as a forest class based on input from Merry Gallagher, Maine DIFW.

Unit: %

Sort Order: Descending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Upstream acres of alewife ponds

GIS Field Name: usAlewifeAcres

Category: Fish Habitat – Alewife

Used in Prioritization Scenario(s): Alewife


The summed acreage of documented alewife spawning ponds upstream of the barrier. This metric evaluates potential upstream alewife spawning habitat, thus ignoring any upstream barriers.

Unit: Acres

Sort Order: Descending

Data Sources:

- Alewife Ponds (2024): DMR, TNC Maine



Probability of brook trout occupancy (+2°C)

GIS Field Name: occ_temp7p20

Category: Fish Habitat – Inland Brook Trout (BKT)

Used in Prioritization Scenario(s): Inland Brook Trout

Occupancy probability of Eastern Brook trout for the catchment (NHDHRDV2) that the barrier is located within, given a 2°C increase in July water temperature.

Unit: %

Sort Order: Descending

Data Sources:

- EcoSheds Catchments (2024): USGS/EcoSheds program

Heritage Fish Pond Barrier

GIS Field Name: HeritagePondBarrier

Category: Fish Habitat – Inland Brook Trout (BKT)

Used in Prioritization Scenario(s): Inland Brook Trout


Barrier separates two functional network that both include heritage wild Eastern Brook trout ponds.

Unit: Boolean

Sort Order: Descending

Data Sources:

- Maine Heritage Fish Waters (2025): IFW



Number of miles of (high, very high) brook trout habitat in US network

GIS Field Name: MilesBT_HVH_USonly

Category: Fish Habitat – Inland Brook Trout (BKT)

Used in Prioritization Scenario(s): Coastal Anadromous, Inland Brook Trout

Length of Eastern Brook trout habitat in the Upstream Functional Network that is of high or very high priority for Eastern Brook trout

Unit: Miles

Sort Order: Descending

Data Sources:

- Brook Trout Habitat Rating (2024): IFW

Number of miles of (high, very high) brook trout habitat in US + DS networks

GIS Field Name: MilesBT_HVH

Category: Fish Habitat – Inland Brook Trout (BKT)

Used in Prioritization Scenario(s): Coastal Anadromous, Inland Brook Trout

Length of Eastern Brook trout habitat in the combined [Upstream Functional Network](#) + [Downstream Functional Network](#) that is of high, or very high priority for Eastern Brook trout

Unit: Miles

Sort Order: Descending

Data Sources:

- Brook Trout Habitat Rating (2024): IFW

EBTJV Wild Brook Trout Patch

GIS Field Name: EBTJV_Patch

Category in Tool: Fish Habitat – Inland Brook Trout (BKT)

Unit: Boolean

Sort Order: Descending

Data Sources:

- Eastern Brook Trout Joint Venture Wild Trout Patches (2025): Eastern Brook Trout Joint Venture

Probability of brook trout occupancy $\geq 50\%$ ($+2^{\circ}\text{C}$)

GIS Field Name: bkt_p20_occ_gt50

Category: Fish Habitat - Inland Brook Trout (BKT)

Used in Prioritization Scenario(s): Inland Brook Trout

Occupancy probability of Eastern Brook trout for the catchment (NHDHRDV2) that the barrier is located within is $\geq 50\%$, given a 2°C increase in July water temperature.

Unit: Boolean

Sort Order: Descending

Data Sources:

- EcoSheds Catchments (2024): USGS/EcoSheds program

Barrier to Sea-Run Brook Trout

GIS Field Name: SeaRunBT

Category: Fish Habitat – Inland Brook Trout (BKT)

Used in Prioritization Scenario(s): Coastal Anadromous

Barrier potentially blocks sea-run Eastern Brook trout migrations because it falls on reaches mapped for sea-run Eastern Brook trout use.

Unit: Boolean

Sort Order: Descending

Data Sources:

- Sea Run Brook Trout Streams (2024): TNC Maine, IFW

Barrier is in a Salmon Critical Habitat HUC10

GIS Field Name: HUC10_CritSalmHab

Category: Fish Habitat – Atlantic Salmon

Used in Prioritization Scenario(s): Atlantic salmon

Barrier is located within a designated Atlantic salmon Critical Habitat [10-Digit Hydrologic Unit](#).

Unit: Boolean

Sort Order: Descending

Data Sources:

- Atlantic Salmon Critical Habitat (2023): USDA-NRCS, NOAA

Atlantic Salmon Habitat Quality

GIS Name: SalmonHabQual

Category: Fish Habitat – Atlantic Salmon

Used in Prioritization Scenario(s): Atlantic salmon

The Atlantic salmon habitat quality ranking of the flowline on which the barrier falls.

Unit: Categorical (Very High (4), High (3), Medium(2), Low(1), No (0))

Sort Order: Descending

Data Sources:

- Atlantic Salmon Habitat Quality (2024), DMR, TNC Maine

Summed Salmon Habitat Units in US Network

GIS Field Name: US_SalmHabUnits

Category: Fish Habitat – Atlantic Salmon

The summed value of Atlantic salmon habitat units obtained from a GIS-based habitat model that predicts the amount of accessible Atlantic salmon rearing habitat throughout the Gulf of Maine Atlantic salmon DPS

Units: Habitat units (sq. meters/100)

Sort Order: Descending

Data Sources:

- Atlantic Salmon Modeled Habitat Units (2012): USFWS Gulf of Maine Coastal Program, NOAA

Salmon Habitat Units in HUC10

GIS Field Name: HUC10_Habitat_Units

Category: Fish Habitat – Atlantic Salmon

The sum of Atlantic salmon habitat units within the 10-Digit Hydrologic Unit (HUC10) that the barrier is within

Unit: Index

Sort Order: Descending

Data Sources:

- Atlantic Salmon Modeled Habitat Units (2012): USFWS Gulf of ME Coastal Program, NOAA

Cumulative Upstream Salmon Habitat Units

GIS Field Name: cumulativeSalmHabUnits

Category: Fish Habitat – Atlantic Salmon

Used in Prioritization Scenario(s): Atlantic salmon

The cumulative upstream Atlantic salmon habitat units metric assesses the amount of upstream Atlantic salmon habitat units that would be made available upstream of a barrier, accounting accounts for the cumulative impacts of partial passability at upstream barriers. For a given target barrier, it is calculated as the sum of [the probability weighted salmon habitat units](#) in the barrier’s upstream functional network, plus the weighted Atlantic salmon habitat units in the upstream functional networks of all of its upstream barriers multiplied by the cumulative [passability score](#) between the target barrier and each upstream barrier. Given the sketch below, the cumulative upstream Atlantic salmon habitat units for the target barrier would be:


$$\begin{aligned}
 & (416)+(450*0.5)+(325*0.5*0.75)+(128*0.5*0.75*0.5)+(140*0.5*0)+(50*0.5*0*1) \\
 & = \\
 & 416 + 225 + 121.875 + 24 + 0 + 0 \\
 & = \\
 & \text{786.875}
 \end{aligned}$$

Unit: Index (cumulative salmon habitat * cumulative passability)

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine
- Atlantic Salmon Modeled Habitat Units (2012): USFWS Gulf of ME Coastal Program, NOAA



Smelt spawning sites upstream of barrier

GIS Field Name: usSmeltSite

Category: Fish Habitat – Smelt

Used in Prioritization Scenario(s): Coastal Anadromous

Value of smelt spawning sites priorities located upstream of barrier. Priority classes were assigned the numerical values in the table below. If more than one spawning site is located above the barrier, the maximum value is assigned to the barrier.

Unit: Boolean (Yes/No)

Sort Order: Descending

Data Sources:

- Smelt Reaches (2024): DMR, TNC Maine

Species of Management Concern on one side of a barrier and not the other

GIS Field Name: InvasiveBarrier

Category: Invasive Species

Species of management concern are found on one side of a barrier and not the other. Species evaluated include Black crappie, Largemouth bass, Muskellunge, Northern pike, Smallmouth bass, and Walleye.

***Note!** *This dataset was not updated for the 2025 iteration of the prioritization tool.*

Unit: Boolean

Sort Order: Ascending

Data Sources:

- Fish Species of Management Concern / Invasive Fish Species (2018 approx.): IFW

**Additional metrics
calculated, but not
displayed in the tool**

Downstream Functional Network Length

GIS Field Name: batFuncDS

Category (not displayed in tool): Network

Length of the functional network downstream of a barrier.


“Functional network” is defined by those sections of river that a fish could theoretically access from any other point within that functional network. Terminal ends are barriers, headwaters, and/or the river mouth.

Unit: meters (converted to miles in some outputs)

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine



Cumulative Upstream Functional Network

GIS Field Name: cumulativeUSNet

Category (not displayed in tool): Network

The cumulative upstream functional network metric assesses the amount of river habitat that would be made available upstream of a barrier, accounting accounts for the cumulative impacts of partial passability at upstream barriers. be made available upstream of a barrier, accounting accounts for the cumulative impacts of partial passability at upstream barriers. For a given target barrier, it is calculated as the sum of the barrier's upstream functional network, plus the upstream functional networks of all of its upstream barriers multiplied by the cumulative passability score between the target barrier and each upstream barrier. Given the sketch below, the cumulative upstream functional network for the target barrier would be:


$$\begin{aligned}
 & (4,230) + (3,150 * 0.5) + (2,890 * 0.5 * 0.75) + (1,020 * 0.5 * 0.75 * 0.5) + (2,680 * 0.5 * 0) + (1,785 * 0.5 * 0 * 1) \\
 & = \\
 & 4,230 + 1,575 + 1083.75 + 191.25 + 0 + 0 \\
 & = \\
 & \text{7,080}
 \end{aligned}$$

Unit: Index (cumulative meters * cumulative passability)

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine



Upstream Barrier Count

GIS Field Name: batCountUS

Category (not displayed in tool): Network

The number of barriers upstream of a given barrier.

This metric is used in the generation of other metrics. It is not available for prioritizations. Includes all barriers which are used in network generation (including natural barriers and restricted barriers).

Does not include barriers excluded from network dataset creation

Unit: #

Sort Order: Ascending

Data Sources:

- Hydrography (2023): USGS, TNC Maine

Density of barriers on upstream network

GIS Field Name: batUSDnsty

Category (not displayed in tool): Network

The number of other barriers upstream of a given barrier divided by the length of upstream river (batCountUS / batLenUS).

Includes all barriers which are used in network generation (including natural barriers and restricted barriers)

Unit: #/m

Sort Order: Ascending

Data Sources:

- Hydrography (2023): USGS, TNC Maine

Number of Natural Barriers on Downstream Flowpath

GIS Field Name: DSFalls

Category (not displayed in tool): Network

Count of natural barriers (e.g. waterfalls) on downstream flowpath of a barrier

Unit: #

Sort Order: Ascending

Data Sources:

- Hydrography (2023): USGS, TNC Maine
- Natural Barriers (2025), TNC Maine

One or More of the Next Upstream Barriers is a Natural Barrier

GIS Field Name: usNatBar

Category (not displayed in tool): Network

Value is 'Yes' if one or more of the next upstream barriers is a natural barrier (e.g. waterfall)

Unit: Boolean

Sort Order: Ascending

Data Sources:

- Hydrography (2023): USGS, TNC Maine
- Natural Barriers (2025), TNC Maine

Upstream Restoration Potential

GIS Field Name: usRestPot

Category (not displayed in tool): Network

The upstream restoration potential is an index designed to summarize the restoration potential of a barrier. It is a measure of the total upstream river habitat, divided by the number of upstream barriers, and multiplied by the passability scores of downstream barriers.

Specifically, it is calculated as

$$((((\text{batLenUS} - \text{batFuncUS}) / \text{batCountUS}) + \text{batFunUS}) * \text{dsPasabilityProduct})$$

Unit: Index

Sort Order: Descending

Data Sources:

- Hydrography (2023): USGS, TNC Maine

Aquifers in upstream or downstream functional networks or both

GIS Field Name: Aquifer

Category (not displayed in tool): Landscape

Aquifers are found in upstream functional network OR downstream functional network OR both. This metric evaluates the potential for cold water habitats within a barrier's respective networks.

Unit: Boolean

Sort Order: Descending

Data Sources:

- Aquifers (2020): MGS

Percent conserved land in 100m buffer of upstream functional network

GIS Name: usCons100

Category (not displayed in tool): Landscape

The percent of a 100m buffer of a barrier's upstream functional network that is in conservation. Conservation Lands included in the current analysis included those with gap status 1,2,3, and 39 – based on TNC's modified version of USGS's [Gap Status Codes](#).

Unit: %

Data Sources:

- Conservation Lands / Secured Lands (2025): TNC Maine

Percent conserved land in 100m buffer of downstream functional network

GIS Name: dsCons100

Category (not displayed in tool): Landscape

The percent of a 100m buffer of a barrier's downstream functional network that is in conservation. Conservation Lands included in the current analysis included those with gap status 1,2,3, and 39 – based on TNC's modified version of USGS's [Gap Status Codes](#).

Unit: %

Data Sources:

- Conservation Lands / Secured Lands (2025): TNC Maine

Percent forest cover in 100m buffer of downstream network

GIS Name: buff100_DS_Perc_Statewide_forestMerged

Category (not displayed in tool): Landscape

Percent forested landcover in a 100m buffer of a barrier's downstream functional network.

Forested landcover classes were aggregated separately for northern Maine and southern Maine. In northern Maine, forest cover was defined as deciduous forest, evergreen forest, mixed forest, and woody wetlands. In southern Maine, scrub/shrub was also included as a forest class based on input from Merry Gallagher, Maine IFW.

Unit: %

Sort Order: Descending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent medium/high intensity developed landcover in 100m buffer of upstream network

GIS Name: buff100_US_Perc_nlcd_2016_developedMedHigh

Category (not displayed in tool): Landscape

Percent of medium or high intensity developed landcover in a 100m buffer of a barrier's upstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent medium/high intensity developed landcover in 100m buffer of downstream network

GIS Name: buff100_DS_Perc_nlcd_2016_developedMedHigh

Category (not displayed in tool): Landscape

Percent of medium or high intensity developed landcover in a 100m buffer of a barrier's downstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent medium/high intensity developed landcover in 100m buffer of total functional network

GIS Name: buff100_Tot_Perc_nlcd_2016_developedMedHigh

Category (not displayed in tool): Landscape

Percent of medium or high intensity developed landcover in a 100m buffer of a barrier's total functional network (upstream + downstream).

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent cultivated crops in 100m buffer of upstream network

GIS Name: buff100_US_Perc_nlcd_2016_cultivatedCrops

Category (not displayed in tool): Landscape

Percent of cultivated crop landcover in a 100m buffer of a barrier's upstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent cultivated crops in 100m buffer of downstream network

GIS Name: buff100_DS_Perc_nlcd_2016_cultivatedCrops

Category (not displayed in tool): Landscape

Percent of cultivated crop landcover in a 100m buffer of a barrier's downstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent cultivated crops in 100m buffer of total functional network

GIS Name: buff100_Tot_Perc_nlcd_2016_cultivatedCrops

Category (not displayed in tool): Landscape

Percent of cultivated crop landcover in a 100m buffer of a barrier's total (upstream + downstream) functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent pasture/hay in 100m buffer of upstream network

GIS Name: buff100_US_Perc_nlcd_2016_pastureHay

Category (not displayed in tool): Landscape

Percent of pasture/hay landcover in a 100m buffer of a barrier's upstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent pasture/hay in 100m buffer of downstream network

GIS Name: buff100_DS_Perc_nlcd_2016_pastureHay

Category (not displayed in tool): Landscape

Percent of pasture/hay landcover in a 100m buffer of a barrier's downstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent pasture/hay in a 100m buffer of total functional network

GIS Name: buff100_Tot_Perc_nlcd_2016_pastureHay

Category (not displayed in tool): Landscape

Percent of pasture/hay landcover in a 100m buffer of a barrier's total (upstream + downstream) functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium

Percent forest cover in the floodplain of upstream network

GIS Name: floodplain_US_Perc_Statewide_forestMerged

Category (not displayed in tool): Landscape

Percent forested landcover in the floodplain of a barrier's upstream functional network.

Forested landcover classes were aggregated separately for northern Maine and southern Maine. In northern Maine, forest cover was defined as deciduous forest, evergreen forest, mixed forest, and woody wetlands. In southern Maine, scrub/shrub was also included as a forest class based on input from Merry Gallagher, Maine IF & W.

Unit: %

Sort Order: Descending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent forest cover in the floodplain of downstream network

GIS Name: floodplain_DS_Perc_Statewide_forestMerged

Category (not displayed in tool): Landscape

Percent forested landcover in the floodplain of a barrier's downstream functional network.

Forested landcover classes were aggregated separately for northern Maine and southern Maine. In northern Maine, forest cover was defined as deciduous forest, evergreen forest, mixed forest, and woody wetlands. In southern Maine, scrub/shrub was also included as a forest class based on input from Merry Gallagher, Maine DIFW.

Unit: %

Sort Order: Descending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent forest cover in the floodplain of total functional network

GIS Name: floodplain_Tot_Perc_Statewide_forestMerged

Category (not displayed in tool): Landscape

Percent forested landcover in the floodplain of a barrier's total functional network (upstream + downstream).

Forested landcover classes were aggregated separately for northern Maine and southern Maine. In northern Maine, forest cover was defined as deciduous forest, evergreen forest, mixed forest, and woody wetlands. In southern Maine, scrub/shrub was also included as a forest class based on input from Merry Gallagher, Maine DIFW.

Unit: %

Sort Order: Descending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent medium/high intensity developed landcover in the floodplain of upstream network

GIS Name: floodplain_US_Perc_nlcd_2016_developedMedHigh

Category (not displayed in tool): Landscape

Percent of medium or high intensity developed landcover in the floodplain of a barrier's upstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent medium/high intensity developed landcover in the floodplain of downstream network

GIS Name: floodplain_DS_Perc_nlcd_2016_developedMedHigh

Category (not displayed in tool): Landscape

Percent of medium or high intensity developed landcover in the floodplain of a barrier's downstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent medium/high intensity developed landcover in the floodplain of total functional network

GIS Name: floodplain_Tot_Perc_nlcd_2016_developedMedHigh

Category (not displayed in tool): Landscape

Percent of medium or high intensity developed landcover in the floodplain of a barrier's total functional network (upstream + downstream).

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent cultivated crops in the floodplain of upstream network

GIS Name: floodplain_US_Perc_nlcd_2016_cultivatedCrops

Category (not displayed in tool): Landscape

Percent of cultivated crop landcover in the floodplain of a barrier's upstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent cultivated crops in the floodplain of downstream network

GIS Name: floodplain_DS_Perc_nlcd_2016_cultivatedCrops

Category (not displayed in tool): Landscape

Percent of cultivated crop landcover in the floodplain of a barrier's downstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent cultivated crops in the floodplain of total functional network

GIS Name: floodplain_Tot_Perc_nlcd_2016_cultivatedCrops

Category (not displayed in tool): Landscape

Percent of cultivated crop landcover in the floodplain of a barrier's total (upstream + downstream) functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent pasture/hay in the floodplain of upstream network

GIS Name: floodplain_US_Perc_nlcd_2016_pastureHay

Category (not displayed in tool): Landscape

Percent of pasture/hay landcover in the floodplain of a barrier's upstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent pasture/hay in the floodplain of downstream network

GIS Name: floodplain_DS_Perc_nlcd_2016_pastureHay

Category (not displayed in tool): Landscape

Percent of pasture/hay landcover in the floodplain of a barrier's downstream functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

Percent pasture/hay in the floodplain of total functional network

GIS Name: floodplain_Tot_Perc_nlcd_2016_pastureHay

Category (not displayed in tool): Landscape

Percent of pasture/hay landcover in the floodplain of a barrier's total (upstream + downstream) functional network.

Unit: %

Sort Order: Ascending

Data Sources:

- Land Cover (NLCD 2021): MRLC Consortium
- Floodplains (2019): Fathom, Inc.

River Size Class

GIS Field Name: NESZCL_Int

Category (not displayed in tool): "Land"scape ('Water'scape)

River size class based on [NE Aquatic Habitat Classification](#).

Sort Order: A (used to increase priority of barriers on smaller streams)

Data Sources:

- USGS 10m resolution Digital Elevation Model-derived Flow-direction raster: TNC Maine

- 1a: Headwaters (<3.861 sq.mi.)
- 1b: Creeks ($\geq 3.861 < 38.61$ sq.mi.)
- 2: Small River ($\geq 38.61 < 200$ sq. mi.)
- 3a: Medium Tributary Rivers ($\geq 200 < 1000$ sq.mi.)
- 3b: Medium Mainstem Rivers ($\geq 1000 < 3861$ sq.
- 4: Large Rivers ($\geq 3861 < 9653$ sq.mi.)
- 5: Great Rivers (≥ 9653 sq.mi.)

(measure = upstream drainage area)

Miles of tidal habitat (current) upstream of barrier

GIS Field Name: MilesUS_HATcurrent

Category (not displayed in tool): Landscape/Fish Habitat

Miles of tidal habitat upstream of a barrier where tidal habitat is defined using the Highest Annual Tide Unit: Miles

Sort Order: Descending

Data Sources:

- Tidal Waters / Sea-level Rise Scenarios (2018): MGS

Miles of tidal habitat (current HAT +1m) upstream of barrier

GIS Field Name: MilesUS_HATPlus1m

Category (not displayed in tool): Landscape/Fish Habitat

Miles of tidal habitat upstream of a barrier where tidal habitat is defined using the Highest Annual Tide (HAT2018plus 1m dataset:

Unit: Miles

Sort Order: Descending

Data Sources:

- Tidal Waters / Sea-level Rise Scenarios (2018): MGS

Miles of tidal habitat (current HAT +6ft) upstream of barrier

GIS Field Name: MilesUS_HATPlus6ft

Category (not displayed in tool): Landscape/Fish Habitat

Miles of tidal habitat upstream of a barrier where tidal habitat is defined using the Highest Annual Tide (HAT2018) plus 6ft dataset:

Unit: Miles

Sort Order: Descending

Data Sources:

- Tidal Waters / Sea-level Rise Scenarios (2018): MGS

Probability of brook trout occupancy $\geq 50\%$ (current temps)

GIS Field Name: bkt_curr_occ_gt50

Category (not displayed in tool): Fish Habitat – Inland Brook Trout (BKT)

Occupancy probability of Eastern Brook trout for the catchment (NHDHRDV2) that the barrier is located within is $\geq 50\%$, given current conditions.

Unit: Boolean

Sort Order: Descending

Data Sources:

- EcoSheds Catchments (2024): USGS/EcoSheds program

High or Very High quality brook trout habitat upstream OR downstream of barrier

GIS Field Name: BT_1_4_HVH

Category (not displayed in tool): Fish Habitat – Inland Brook Trout (BKT)

Prioritization Scenario(s): Inland Brook Trout

Presence of ¼ mile or more of high, or very high priority Eastern Brook trout habitat in either the [Upstream Functional Network](#) or [Downstream Functional Network](#)

Unit: Boolean

Sort Order: Descending

Data Sources:

- Brook Trout Habitat Rating (2024): IFW

Very High-quality brook trout habitat upstream OR downstream of barrier

GIS Field Name: BT_1_4_VH

Category (not displayed in tool): Fish Habitat – Inland Brook Trout (BKT)

Presence of ¼ mile or more of very high priority Eastern Brook trout habitat in either the [Upstream Functional Network](#) or [Downstream Functional Network](#)

Unit: Boolean

Sort Order: Descending

Data Sources:

- Brook Trout Habitat Rating (2024): IFW

Probability of brook trout occupancy (current temps)

GIS Field Name: occ_current

Category (not displayed in tool): Fish Habitat – Inland Brook Trout (BKT)

Occupancy probability of Eastern Brook trout for the catchment (NHDHRDV2) that the barrier is located within, given current conditions.

Unit: %

Sort Order: Descending

Data Sources:

- EcoSheds Catchments (2024): USGS/EcoSheds program

Probability of brook trout occupancy (+4°C)

GIS Field Name: occ_temp7p40

Category (not displayed in tool): Fish Habitat – Inland Brook Trout (BKT)

Occupancy probability of Eastern Brook trout for the catchment (NHDHRDV2) that the barrier is located within, given a 4°C increase in July water temperature.

Unit: %

Sort Order: Descending

Data Sources:

- EcoSheds Catchments (2024): USGS/EcoSheds program

Probability of brook trout occupancy $\geq 50\%$ ($+4^{\circ}\text{C}$)

GIS Field Name: bkt_p40_occ_gt50

Category (not displayed in tool): Fish Habitat – Inland Brook Trout (BKT)

Occupancy probability of Eastern Brook trout for the catchment (NHDHRDV2) that the barrier is located within is $\geq 50\%$, given a 4°C increase in July water temperature.

Unit: Boolean

Sort Order: Descending

Data Sources:

- EcoSheds Catchments (2024): USGS/EcoSheds program

Summed salmon parr productivity in US Functional Network

GIS Field Name: US_SalmParrProd

Category (not displayed in tool): Fish Habitat – Atlantic Salmon

The summed Atlantic salmon parr productivity in a barrier's Upstream Functional Network.

Salmon parr productivity obtained from a GIS-based habitat model that predicts the amount of accessible Atlantic salmon rearing habitat throughout the Gulf of Maine Atlantic salmon DPS

Units: # (Predicted parr production based on habitat units and predicted parr density)

Sort Order: Descending

Data Sources:

- Atlantic Salmon Modeled Habitat Units (2012): USFWS Gulf of Maine Coastal Program, NOAA

Species of Management Concern on one side of a barrier and not the other - Confirmed only

GIS Field Name: InvasiveBarrierConfirmed

Category (not displayed in tool): Invasive Species

Species of management concern are confirmed to be found on one side of a barrier and not the other. Species evaluated include Black crappie, Largemouth bass, Muskellunge, Northern pike, Smallmouth bass, and Walleye.

Unit: Boolean

***Note!** *This dataset was not updated for the 2025 iteration of the prioritization tool.*

Sort Order: Ascending

Data Sources:

- Fish Species of Management Concern / Invasive Fish Species (2018 approx.): IFW

Number of Species of Management Concern that are blocked - Confirmed only

GIS Field Name: InvasiveBarrierConfNumSpp

Category (not displayed in tool): Invasive Species

Number of species of management concern are confirmed to be found on one side of a barrier and not the other. Species evaluated include Black crappie, Largemouth bass, Muskellunge, Northern pike, Smallmouth bass, and Walleye.

***Note!** *This dataset was not updated for the 2025 iteration of the prioritization tool.*

Unit: #

Sort Order: Ascending

Data Sources:

- Fish Species of Management Concern / Invasive Fish Species (2018 approx.): IFW