

# Posterior computation for BART

- ▶ In order to generate the posterior for  $f$ , we sample the structure of all the trees  $\mathcal{T}_h$ , for  $h = 1, \dots, H$ ; the values of all leaves  $\mu_{hn}$  for  $n \in \mathcal{L}_h$  within tree  $h$ ; and, the error variance  $\sigma^2$
- ▶ Additionally, with the sparsity prior, there are samples of the vector of splitting variable selection probabilities  $[s_1, \dots, s_P]$  and, when the sparsity parameter is random, samples of  $\theta$
- ▶ The leaf and variance parameters are sampled from the posterior using Gibbs sampling
- ▶ Since the priors on these parameters are conjugate, the Gibbs conditionals are specified analytically
- ▶ The leaves,  $\mu_{hn}$ , are drawn from a Normal conditional density
- ▶ The error variance,  $\sigma^2$ , is drawn from a scaled inverse Chi-square conditional (equivalent to inverse Gamma)

# Posterior computation for BART

- ▶ Drawing a tree from the posterior requires a Metropolis-within-Gibbs sampling scheme, i.e., a Metropolis-Hastings (MH) step within Gibbs sampling
- ▶ For single-tree models, four different proposals are defined in Chipman George 1998 *JASA*
- ▶ The complementary BIRTH/DEATH proposals are essential (the others are CHANGE and SWAP which are optional)
- ▶ This is only one of several schemes that have been proposed; see Pratola 2016 *Bayesian Analysis*
- ▶ For programming simplicity, the **BART/BART3** package only implements BIRTH and DEATH with equal probability
- ▶ BIRTH selects a leaf and turns it into a branch: a new variable and cut-point with two leaves “born” as its descendants
- ▶ DEATH selects a branch leading to two terminal leaves and “kills” the branch by replacing it with a single leaf

# Posterior computation for BART: the BIRTH proposal

- ▶ we present the acceptance probability for a BIRTH proposal
- ▶ a DEATH proposal is the reversible inverse of a BIRTH
- ▶ The algorithm assumes a fixed discrete set of possible split values for each  $x_j$  denoted by  $c_{jk}$
- ▶ the leaf values,  $\mu_{hn}$ , are integrated out so this search in tree space is over a large discrete set of possibilities
- ▶ At the  $m$ th MCMC step, let  $\mathcal{T}_h^m$  denote the current state of tree  $h$  while  $\mathcal{T}_h^*$  denotes the tree proposal
- ▶  $\mathcal{T}_h^*$  are identical  $\mathcal{T}_h^m$  except that one terminal leaf of  $\mathcal{T}_h^m$  is replaced by a branch of  $\mathcal{T}_h^*$  with two terminal leaves

# Posterior computation for BART: the BIRTH proposal

- ▶ The proposed tree is accepted with the following probability:

$$\pi_{\text{BIRTH}} = \min \left( 1, \frac{[\mathcal{T}_h^* | y]}{[\mathcal{T}_h^m | y]} \frac{[\mathcal{T}_h^m | \mathcal{T}_h^*]}{[\mathcal{T}_h^* | \mathcal{T}_h^m]} \right)$$

- ▶  $[\mathcal{T}_h^* | \mathcal{T}_h^m]$  is the probability of proposing  $\mathcal{T}_h^*$  (a BIRTH) given current state  $\mathcal{T}_h^m$
- ▶ while  $[\mathcal{T}_h^m | \mathcal{T}_h^*]$  is the reverse  
the probability of proposing  $\mathcal{T}_h^m$  (a DEATH) given  $\mathcal{T}_h^*$
- ▶ Now we delve into the calculations for  $[\mathcal{T}_h^* | y]$  and  $[\mathcal{T}_h^m | y]$

# Posterior computation for BART: the BIRTH proposal

- ▶ First, we describe the likelihood contribution to the posterior
- ▶ Let  $y_n$  denote the partition of  $y$  corresponding to the leaf node  $n$  given the tree  $\mathcal{T}_h$
- ▶ Because the leaf values are a priori conditionally independent, we have  $[y|\mathcal{T}_h] = \prod_n [y_n|\mathcal{T}_h]$  where  $[y_n|\mathcal{T}_h] = \int [\mu_n] [y_n|\mathcal{T}_h, \mu_n] d\mu_n$  (we will return to this later)
- ▶ After cancellation of terms in the ratio  $\frac{[\mathcal{T}_h^*|y]}{[\mathcal{T}_h^m|y]}$  we have the likelihood contributions

$$\frac{[y_L, y_R|\mathcal{T}_h^*]}{[y_n|\mathcal{T}_h^m]} = \frac{[y_L|\mathcal{T}_h^*] [y_R|\mathcal{T}_h^*]}{[y_n|\mathcal{T}_h^m]}$$

$y_L$ : partition for the newborn left leaf node

$y_R$ : partition for the newborn right leaf node

# Posterior computation for BART

- ▶ Similarly, the terms that the prior contributes to the posterior ratio often cancel since there is only one **area** where the trees differ: either a terminal leaf or a penultimate branch
- ▶ Therefore, the prior contribution to  $\frac{[\mathcal{T}_h^*|y]}{[\mathcal{T}_h^m|y]}$  is

$$\frac{P[\psi_n = 1] P[\psi_l = 0] P[\psi_r = 0] s_j}{P[\psi_n = 0]} = \frac{\alpha(d(n) + 1)^{-\beta} [1 - \alpha(d(n) + 2)^{-\beta}]^2 s_j}{1 - \alpha(d(n) + 1)^{-\beta}}$$

- ▶  $P[\psi_n]$  is the branch regularity prior  
 $s_j$  is the splitting variable selection probability  
 $n$  is the chosen leaf node in tree  $\mathcal{T}_h^m$   
 $l = 2n$  is the newborn left leaf node in tree  $\mathcal{T}_h^*$   
 $r = 2n + 1$  is the newborn right leaf node in tree  $\mathcal{T}_h^*$

# Posterior computation for BART

- Finally, the ratio  $\frac{[\mathcal{T}_h^m | \mathcal{T}_h^*]}{[\mathcal{T}_h^* | \mathcal{T}_h^m]}$  is

$$\frac{[\text{DEATH} | \mathcal{T}_h^*] [n | \mathcal{T}_h^*]}{[\text{BIRTH} | \mathcal{T}_h^m] [n | \mathcal{T}_h^m] s_j}$$

- $[n | \mathcal{T}_h]$  is the probability of choosing node  $n$  given tree  $\mathcal{T}_h$
- N.B.  $s_j$  appears in both the numerator and denominator of the acceptance probability  $\pi_{\text{BIRTH}}$ , therefore, canceling which is mathematically convenient

# Posterior computation for weighted BART

$$y_i = f(x_i) + \epsilon_i \text{ where } f \stackrel{\text{prior}}{\sim} \text{BART}$$

$$\epsilon_i \sim N(0, \mathbf{w}_i^2 \sigma^2)$$

BART variant	Weights	Comment
Homoskedastic	$\mathbf{w}_i = 1$	<i>Chipman, George et al. 2010</i>
Weighted		Non-trivial known weights (circa 2015)
Heteroskedastic	$\mathbf{w}_i = s(x_i)$	$s^2 \stackrel{\text{prior}}{\sim} \mathbf{HBART}$ <i>Pratola, Chipman et al. 2020</i>

- ▶ Until now, we have not dwelled on BART variants
- ▶ From here on, we will focus on weighted BART
- ▶ This development extends the homoskedastic variant and naturally incorporates heteroskedastic BART



## Posterior computation for weighted BART: drawing a tree

- Based on the centered data:  $y_i = \tilde{y}_i - \mu$
- Given that the  $n$ th leaf only depends on  $(y_{i'}, w_{i'})$  where  $i' = 1, \dots, m_n$

$$\mu_{hl} \stackrel{\text{prior}}{\sim} \mathbf{N}(0, \eta_0^{-1})$$

$$\eta_{i'} = w_{i'}^{-2} \sigma^{-2}$$

$$\tilde{\mu} = \eta_n^{-1} \sum_{i'} \eta_{i'} y_{i'}$$

$$\eta_0^{-1} = \frac{\tau^2}{4H\kappa^2}$$
$$\eta_n = \sum_{i'} \eta_{i'}$$

$$\begin{aligned} [y_n | \mathcal{T}_h] &= \int [\mu_n] [y_n | \mathcal{T}_h, \mu_n] d\mu_n \\ &\propto \frac{\prod_{i'} \eta_{i'}^{0.5}}{(2\pi)^{0.5m_n}} \frac{\sqrt{\eta_0}}{\sqrt{\eta_0 + \eta_n}} \exp\left(-0.5 \frac{\eta_0 \eta_n}{\eta_0 + \eta_n} \tilde{\mu}^2\right) \\ &\quad \times \exp\left(-0.5 \sum_{i'} \eta_{i'} (y_{i'} - \tilde{\mu})^2\right) \end{aligned}$$

# Posterior computation for weighted BART: BIRTH

- ▶ Based on the centered data:  $y_i = \tilde{y}_i - \mu$
- ▶ Given that the  $n$ th leaf only depends on  $(y_{i'}, w_{i'})$  where  $i' = 1, \dots, m_n$

$$\eta_n = \eta_L + \eta_R$$

$$\xi_n = \sum_{i'} \eta_{i'} y_{i'} = \xi_L + \xi_R$$

$$\mu_{hl} \stackrel{\text{prior}}{\sim} \text{N}(0, \sigma_0^2)$$

$$\sigma_0^2 = \frac{\tau^2}{4H_K^2}$$

$$[y_n | \mathcal{T}_h] \propto \omega_n \exp(0.5 \sigma_0^2 \xi_n^2 \omega_n^2)$$

$$\omega_n = (\sigma_0^2 \eta_n + 1)^{-0.5}$$

$$[y_L | \mathcal{T}_h] \propto \omega_L \exp(0.5 \sigma_0^2 \xi_L^2 \omega_L^2)$$

$$\omega_L = (\sigma_0^2 \eta_L + 1)^{-0.5}$$

$$[y_R | \mathcal{T}_h] \propto \omega_R \exp(0.5 \sigma_0^2 \xi_R^2 \omega_R^2)$$

$$\omega_R = (\sigma_0^2 \eta_R + 1)^{-0.5}$$

# Posterior computation for weighted BART: BIRTH

- After cancellation of terms in the ratio  $\frac{[\mathcal{T}_h^*|y]}{[\mathcal{T}_h^m|y]}$  we have the likelihood contributions

$$\begin{aligned}\frac{[y_L, y_R | \mathcal{T}_h^*]}{[y_n | \mathcal{T}_h^m]} &= \frac{[y_L | \mathcal{T}_h^*] [y_R | \mathcal{T}_h^*]}{[y_n | \mathcal{T}_h^m]} \\ &= \omega_L \omega_R \omega_n^{-1} \exp 0.5 \sigma_0^2 (\xi_L^2 \omega_L^2 + \xi_R^2 \omega_R^2 - \xi_n^2 \omega_n^2)\end{aligned}$$

## Posterior computation for weighted BART: the leaves

- ▶ Based on the centered data:  $y_i = \tilde{y}_i - \mu$
- ▶ Given  $\mathcal{T}_h$ , suppose that the  $l$ th leaf,  $\mu_{hl}$ , only depends on  $(y_{i'}, w_{i'})$  where  $i' = 1, \dots, m_{hl}$
- ▶ Define the residuals,  $z_{i'}$ , due to everything except  $(\mathcal{T}_h, \mathcal{M}_h)$

$$z_{i'} = y_{i'} - \sum_{h' \neq h} g(x_{i'}; \mathcal{T}_{h'}, \mathcal{M}_{h'})$$
$$\sim N(\mu_{hl}, w_{i'}^2 \sigma^2)$$

$$\bar{z}_{\cdot} = N(\mu_{hl}, \eta_{hl}^{-1})$$

$$\mu_{hl} \stackrel{\text{prior}}{\sim} N(0, \eta_0^{-1})$$

$$\eta_{hl}^{-1} = \frac{\sigma^2}{m_{hl}^2} \sum_{i'} w_{i'}^2$$

$$\eta_0^{-1} = \frac{\tau^2}{4H\kappa^2}$$

$$\mu_{hl} | (\mathcal{T}_h, y) \sim N\left(\frac{\eta_{hl} m_{hl} \bar{z}_{\cdot}}{\eta_0 + \eta_{hl} m_{hl}}, \frac{1}{\eta_0 + \eta_{hl} m_{hl}}\right)$$

# Posterior computation for BART: uncertainty intervals

- ▶ Suppose that we want to estimate  $f$  at some value  $x$  whether from the training or testing
- ▶ The standard Bayesian estimate is  $\hat{f}(x) = M^{-1} \sum_m f_m(x)$
- ▶ Construct a Bayesian  $(1 - \alpha) \times 100\%$  credible interval from the  $\alpha/2$  and the  $1 - \alpha/2$  quantiles of the posterior
- ▶  $(f_{(\alpha/2)}(x), f_{(1-\alpha/2)}(x))$
- ▶ Construct a Bayesian  $(1 - \alpha) \times 100\%$  prediction interval
- ▶ Generate  $\tilde{y}$  from the predictive distribution:  
 $\tilde{y}_m \sim N(f_m(x), \sigma_m^2)$
- ▶ Select the  $\alpha/2$  and the  $1 - \alpha/2$  quantiles of  $\tilde{y}$
- ▶  $(\tilde{y}_{(\alpha/2)}, \tilde{y}_{(1-\alpha/2)})$