


Chapter 1: Linear Systems and Matrices

Section: Inverse of a Matrix

Lecture #4

Lebanese University

Prof Ali WEHBE



- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancelation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Definition

A system of linear equation is said to be **Homogeneous** if the constant terms of each equation is equal to **zero**, i.e. it is of the form;

$$(S) : \begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = 0, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = 0, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = 0, \end{cases}$$

Definition

A system of linear equation is said to be **Homogeneous** if the constant terms of each equation is equal to **zero**, i.e. it is of the form;

$$(S) : \begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = 0, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = 0, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = 0, \end{cases}$$

Theorem

Homogeneous system are always **consistent**. Moreover, if the number of equations are less than the number of unknowns, then the Homogeneous System has an infinite number of solutions.

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Example (Supplementary 1)

Discuss according the value of a for which the consistence of the following system:

$$(S) \begin{cases} x & +2y & +z & = & 2, \\ 2x & -2y & +3z & = & 1, \\ x & +2y & -az & = & a. \end{cases}$$

Solution:

Example (Supplementary 2)

Discuss according the value of a for which the consistence of the following system:

$$(S) \begin{cases} x & +y & +7z & = & -7, \\ 2x & +3y & +17z & = & 11, \\ x & +2y & +(a^2 + 1)z & = & 6a. \end{cases}$$

Solution:

Example (Supplementary 3)

Discuss according the values of a and b for which the consistence of the following system:

$$(S) \begin{cases} ax & & +bz & = & 2, \\ ax & +ay & +4z & = & 4, \\ & ay & +2z & = & b. \end{cases}$$

Solution:

Example (Supplementary 4)

Solve the following system of non-linear equations for the unknowns angles α , β and γ , where $0 \leq \alpha \leq \frac{\pi}{2}$, $0 \leq \beta < 2\pi$ and $0 \leq \gamma \leq \pi$.

$$(S) \begin{cases} 2 \sin \alpha & + \cos \beta & - \tan \gamma & = & 1, \\ -4 \sin \alpha & + \cos \beta & + \tan \gamma & = & 0, \\ -2 \sin \alpha & + 3 \cos \beta & + 2 \tan \gamma & = & 4. \end{cases}$$

Solution:

Example (Supplementary 5)

Find the values of ***a***, ***b*** and ***c*** such that the graph of the polynomial $p(x) = ax^2 + bx + c$ passes through the points **(1, 2)**, **(−1, 6)** and **(2, 3)**.

Solution:

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Definition

Let A be $n \times n$ Matrix.

The matrix A is said to be **nonsingular** or **Invertible** if there exists a matrix B such that:

$$AB = BA = I_n$$

The matrix B is called the inverse of A and we denote it by A^{-1} .

Definition

Let A be $n \times n$ Matrix.

The matrix A is said to be **nonsingular** or **Invertible** if there exists a matrix B such that:

$$AB = BA = I_n$$

The matrix B is called the inverse of A and we denote it by A^{-1} .

Example

a) The matrices $\begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}$ and $\begin{bmatrix} -\frac{1}{10} & \frac{2}{5} \\ \frac{3}{10} & -\frac{1}{5} \end{bmatrix}$ are the inverses of each other.

Definition

Let A be $n \times n$ Matrix.

The matrix A is said to be **nonsingular** or **Invertible** if there exists a matrix B such that:

$$AB = BA = I_n$$

The matrix B is called the inverse of A and we denote it by A^{-1} .

Example

a) The matrices $\begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}$ and $\begin{bmatrix} -\frac{1}{3} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix}$ are the inverses of each other. Since

$$\begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{3} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2$$

Definition

Let A be $n \times n$ Matrix.

The matrix A is said to be **nonsingular** or **Invertible** if there exists a matrix B such that:

$$AB = BA = I_n$$

The matrix B is called the inverse of A and we denote it by A^{-1} .

Example

a) The matrices $\begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}$ and $\begin{bmatrix} -\frac{1}{3} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix}$ are the inverses of each other. Since

$$\begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{3} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2$$

and

Definition

Let A be $n \times n$ Matrix.

The matrix A is said to be **nonsingular** or **Invertible** if there exists a matrix B such that:

$$AB = BA = I_n$$

The matrix B is called the inverse of A and we denote it by A^{-1} .

Example

a) The matrices $\begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}$ and $\begin{bmatrix} -\frac{1}{10} & \frac{2}{5} \\ \frac{3}{10} & -\frac{1}{5} \end{bmatrix}$ are the inverses of each other. Since

$$\begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{10} & \frac{2}{5} \\ \frac{3}{10} & -\frac{1}{5} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2$$

and

$$\begin{bmatrix} -\frac{1}{10} & \frac{2}{5} \\ \frac{3}{10} & -\frac{1}{5} \end{bmatrix} \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} = I_2.$$

Example

b) The 3×3 matrices $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & -2 & 5 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix}$ are the inverses of each other,

Example

b) The 3×3 matrices $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & -2 & 5 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix}$ are the inverses of each other,

since

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 5 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3$$

and

$$\begin{bmatrix} 1 & -2 & 5 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3.$$

Remark:

An $n \times n$ matrix is said to be **Singular** or **Non-Invertible** if it does not have an inverse.

Remark:

An $n \times n$ matrix is said to be **Singular** or **Non-Invertible** if it does not have an inverse.

Example

The matrix $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ is a singular one. Why?

Our aim is to answer the following questions:

Our aim is to answer the following questions:

- 1) How can we know if a matrix is invertible or not ?

Our aim is to answer the following questions:

- 1) How can we know if a matrix is invertible or not ?
- 2) If exist, how to find the inverse of a Matrix?

Our aim is to answer the following questions:

- 1) How can we know if a matrix is invertible or not ?
- 2) If exist, how to find the inverse of a Matrix?
- 3) Can we use the inverse of a matrix to solve a system of Linear equation?

Our aim is to answer the following questions:

- 1) How can we know if a matrix is invertible or not ?
- 2) If exist, how to find the inverse of a Matrix?
- 3) Can we use the inverse of a matrix to solve a system of Linear equation?

Answers:

- 1) YES!!, but in the chapter 2 DETERMINANT.

Our aim is to answer the following questions:

- 1) How can we know if a matrix is invertible or not ?
- 2) If exist, how to find the inverse of a Matrix?
- 3) Can we use the inverse of a matrix to solve a system of Linear equation?

Answers:

- 1) YES!!, but in the chapter 2 DETERMINANT.
- 2) In this lecture, we will see now how to determine the inverse of a matrix using **J.G.R.**

Our aim is to answer the following questions:

- 1) How can we know if a matrix is invertible or not ?
- 2) If exist, how to find the inverse of a Matrix?
- 3) Can we use the inverse of a matrix to solve a system of Linear equation?

Answers:

- 1) YES!!, but in the chapter 2 DETERMINANT.
- 2) In this lecture, we will see now how to determine the inverse of a matrix using **J.G.R.**
- 3) In this lecture, we see how to use the inverse of the matrix to solve a System of Linear Equation.

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$,

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}.$

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}$. $ad - bc = -10 \neq 0$

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}$. $ad - bc = -10 \neq 0$

$$\Rightarrow A^{-1} =$$

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}$. $ad - bc = -10 \neq 0$

$$\Rightarrow A^{-1} = -\frac{1}{10} \begin{bmatrix} -3 & -4 \\ -1 & 2 \end{bmatrix} =$$

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}$. $ad - bc = -10 \neq 0$

$$\Rightarrow A^{-1} = -\frac{1}{10} \begin{bmatrix} -3 & -4 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} \frac{3}{10} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix}.$$

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}$. $ad - bc = -10 \neq 0$

$$\Rightarrow A^{-1} = -\frac{1}{10} \begin{bmatrix} -3 & -4 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} \frac{3}{10} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix}.$$

b) $B = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$.

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}$. $ad - bc = -10 \neq 0$

$$\Rightarrow A^{-1} = -\frac{1}{10} \begin{bmatrix} -3 & -4 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} \frac{3}{10} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix}.$$

b) $B = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$. $ad - bc = 0$

Strategy

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an 2×2 matrix such that $ad - bc \neq 0$, then A is invertible and its inverse is given by

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

a) $A = \begin{bmatrix} 2 & 4 \\ 1 & -3 \end{bmatrix}$. $ad - bc = -10 \neq 0$

$$\Rightarrow A^{-1} = -\frac{1}{10} \begin{bmatrix} -3 & -4 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} \frac{3}{10} & \frac{2}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{bmatrix}.$$

b) $B = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$. $ad - bc = 0 \Rightarrow B$ is a singular matrix.

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Strategy

Let A be an $n \times n$ matrix.

Strategy

Let A be an $n \times n$ matrix.

To determine its inverse, we apply the following strategy:

Strategy

Let A be an $n \times n$ matrix.

To determine its inverse, we apply the following strategy:

- 1) We write the $n \times 2n$ matrix $[A|I_n]$.

Strategy

Let A be an $n \times n$ matrix.

To determine its inverse, we apply the following strategy:

- 1) We write the $n \times 2n$ matrix $[A|I_n]$.
- 2) We apply **J.G.R** method to transform the previous $n \times 2n$ matrix to $[I_n|B]$.

Strategy

Let A be an $n \times n$ matrix.

To determine its inverse, we apply the following strategy:

- 1) We write the $n \times 2n$ matrix $[A|I_n]$.
- 2) We apply **J.G.R** method to transform the previous $n \times 2n$ matrix to $[I_n|B]$.
- 3) The matrix B will be the inverse of A , equivalently, $B = A^{-1}$.

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}.$

Solution:

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}.$

Solution:

$$[A|I_3] =$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \begin{bmatrix} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{bmatrix}$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array}$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \begin{bmatrix} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{bmatrix} \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array} \Rightarrow$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \begin{bmatrix} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{bmatrix} \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array} \Rightarrow$$

$$\begin{bmatrix} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & -6 & -3 & -2 & 0 & 1 \end{bmatrix}$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array} \Rightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & -6 & -3 & -2 & 0 & 1 \end{array} \right] R_3 + 3R_2 \longrightarrow R_3$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array} \Rightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & -6 & -3 & -2 & 0 & 1 \end{array} \right] R_3 + 3R_2 \longrightarrow R_3 \Rightarrow$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array} \Rightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & -6 & -3 & -2 & 0 & 1 \end{array} \right] R_3 + 3R_2 \longrightarrow R_3 \Rightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & 0 & 6 & 1 & 3 & 1 \end{array} \right]$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array} \Rightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & -6 & -3 & -2 & 0 & 1 \end{array} \right] R_3 + 3R_2 \longrightarrow R_3 \Rightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & 0 & 6 & 1 & 3 & 1 \end{array} \right] \begin{array}{l} R_2 - \frac{1}{2}R_3 \longrightarrow R_2 \\ R_1 - \frac{1}{2}R_3 \longrightarrow R_1 \end{array}$$

Example (Find the inverse of the following matrices)

a) $A = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$.

Solution:

$$[A|I_3] = \begin{bmatrix} 1 & 4 & 3 & 1 & 0 & 0 \\ -1 & -2 & 0 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 & 0 & 1 \end{bmatrix} \begin{array}{l} R_2 + R_1 \longrightarrow R_2 \\ R_3 - 2R_1 \longrightarrow R_3 \end{array} \Rightarrow$$

$$\begin{bmatrix} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & -6 & -3 & -2 & 0 & 1 \end{bmatrix} R_3 + 3R_2 \longrightarrow R_3 \Rightarrow$$

$$\begin{bmatrix} 1 & 4 & 3 & 1 & 0 & 0 \\ 0 & 2 & 3 & 1 & 1 & 0 \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \begin{array}{l} R_2 - \frac{1}{2}R_3 \longrightarrow R_2 \\ R_1 - \frac{1}{2}R_3 \longrightarrow R_1 \end{array} \Rightarrow$$

Example

$$\begin{bmatrix} 1 & 4 & 0 & \frac{1}{2} & -\frac{3}{2} & -\frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix}$$

Example

$$\begin{bmatrix} 1 & 4 & 0 & \frac{1}{2} & -\frac{3}{2} & -\frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \quad R_1 - 2R_2 \rightarrow R_1$$

Example

$$\begin{bmatrix} 1 & 4 & 0 & \frac{1}{2} & -\frac{3}{2} & -\frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \quad R_1 - 2R_2 \rightarrow R_1 \Rightarrow$$

Example

$$\begin{bmatrix} 1 & 4 & 0 & \frac{1}{2} & -\frac{3}{2} & -\frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \quad R_1 - 2R_2 \rightarrow R_1 \Rightarrow$$
$$\begin{bmatrix} 1 & 0 & 0 & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix}$$

Example

$$\begin{bmatrix} 1 & 4 & 0 & \frac{1}{2} & -\frac{3}{2} & -\frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \quad R_1 - 2R_2 \longrightarrow R_1 \implies$$

$$\begin{bmatrix} 1 & 0 & 0 & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \quad \begin{array}{l} R_2 \times \frac{1}{2} \longrightarrow R_2 \\ R_3 \times \frac{1}{6} \longrightarrow R_3 \end{array}$$

Example

$$\begin{bmatrix} 1 & 4 & 0 & \frac{1}{2} & -\frac{3}{2} & -\frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \quad R_1 - 2R_2 \longrightarrow R_1 \implies$$

$$\begin{bmatrix} 1 & 0 & 0 & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 2 & 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & 6 & 1 & 3 & 1 \end{bmatrix} \quad \begin{array}{l} R_2 \times \frac{1}{2} \longrightarrow R_2 \\ R_3 \times \frac{1}{6} \longrightarrow R_3 \end{array} \implies$$

Example

$$\left[\begin{array}{cccccc} 1 & 0 & 0 & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 0 & \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ 0 & 0 & 1 & \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{array} \right] = [I_3 | B].$$

Example

$$\begin{bmatrix} 1 & 0 & 0 & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 0 & \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ 0 & 0 & 1 & \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} = [I_3 | B].$$

$$AB = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix} \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3$$

and

Example

$$BA =$$

Example

$$BA = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix} =$$

Example

$$BA = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3.$$

Example

$$BA = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3.$$

Thus, $B = A^{-1}$.

Example

$$BA = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3.$$

Thus, $B = A^{-1}$.

b) $A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ -6 & 2 & 3 \end{bmatrix}.$

Solution:

Example

$$[A|I_3]$$

Example

$$[A|I_3] \Rightarrow \begin{array}{lll} \text{Step1 :} & R_2 - R_1 & \longrightarrow R_2. \\ \text{Step2 :} & R_3 + 6R_1 & \longrightarrow R_3. \\ \text{Step3 :} & R_3 + 4R_2 & \longrightarrow R_3. \\ \text{Step4 :} & -R_3 & \longrightarrow R_3. \\ \text{Step5 :} & R_2 + R_3 & \longrightarrow R_2. \\ \text{Step6 :} & R_1 + R_2 & \longrightarrow R_1. \end{array}$$

Example

$$\begin{array}{llll}
 \text{Step1 :} & R_2 - R_1 & \longrightarrow & R_2. \\
 \text{Step2 :} & R_3 + 6R_1 & \longrightarrow & R_3. \\
 \text{Step3 :} & R_3 + 4R_2 & \longrightarrow & R_3. \\
 [A|I_3] \Rightarrow \text{Step4 :} & -R_3 & \longrightarrow & R_3. \Rightarrow [I_3|A^{-1}] \text{ with} \\
 \text{Step5 :} & R_2 + R_3 & \longrightarrow & R_2. \\
 \text{Step6 :} & R_1 + R_2 & \longrightarrow & R_1.
 \end{array}$$

Example

$$\begin{array}{llll}
 \text{Step1 :} & R_2 - R_1 & \longrightarrow & R_2. \\
 \text{Step2 :} & R_3 + 6R_1 & \longrightarrow & R_3. \\
 \text{Step3 :} & R_3 + 4R_2 & \longrightarrow & R_3. \\
 [A|I_3] \Rightarrow \text{Step4 :} & -R_3 & \longrightarrow & R_3. \Rightarrow [I_3|A^{-1}] \text{ with} \\
 \text{Step5 :} & R_2 + R_3 & \longrightarrow & R_2. \\
 \text{Step6 :} & R_1 + R_2 & \longrightarrow & R_1.
 \end{array}$$

$$A^{-1} = \begin{bmatrix} -2 & -3 & -1 \\ -3 & -3 & -1 \\ -2 & -4 & -1 \end{bmatrix}.$$

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Theorem:

If A is an invertible matrix, k is a positive integer and c is a scalar not equal to zero.

Theorem:

If \mathbf{A} is an invertible matrix, k is a positive integer and c is a scalar not equal to zero. Then \mathbf{A}^{-1} , \mathbf{A}^k , $c\mathbf{A}$ and \mathbf{A}^T are invertible and the following statements are true:

Theorem:

If A is an invertible matrix, k is a positive integer and c is a scalar not equal to zero. Then A^{-1} , A^k , cA and A^T are invertible and the following statements are true:

a) $(A^{-1})^{-1} = A$.

Theorem:

If \mathbf{A} is an invertible matrix, k is a positive integer and c is a scalar not equal to zero. Then \mathbf{A}^{-1} , \mathbf{A}^k , $c\mathbf{A}$ and \mathbf{A}^T are invertible and the following statements are true:

- a) $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$.
- b) $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$.

Theorem:

If \mathbf{A} is an invertible matrix, k is a positive integer and c is a scalar not equal to zero. Then \mathbf{A}^{-1} , \mathbf{A}^k , $c\mathbf{A}$ and \mathbf{A}^T are invertible and the following statements are true:

- a) $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$.
- b) $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$.
- c) $(\mathbf{A}^k)^{-1} = (\mathbf{A}^{-1})^k$.

Theorem:

If \mathbf{A} is an invertible matrix, k is a positive integer and c is a scalar not equal to zero. Then \mathbf{A}^{-1} , \mathbf{A}^k , $c\mathbf{A}$ and \mathbf{A}^T are invertible and the following statements are true:

- a) $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$.
- b) $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$.
- c) $(\mathbf{A}^k)^{-1} = (\mathbf{A}^{-1})^k$.
- d) $(c\mathbf{A})^{-1} = \frac{1}{c}\mathbf{A}^{-1}$.

Theorem:

If \mathbf{A} is an invertible matrix, k is a positive integer and c is a scalar not equal to zero. Then \mathbf{A}^{-1} , \mathbf{A}^k , $c\mathbf{A}$ and \mathbf{A}^T are invertible and the following statements are true:

- a) $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$.
- b) $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$.
- c) $(\mathbf{A}^k)^{-1} = (\mathbf{A}^{-1})^k$.
- d) $(c\mathbf{A})^{-1} = \frac{1}{c}\mathbf{A}^{-1}$.

Example

Compute \mathbf{A}^{-2} in two different ways for $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 2 & 4 \end{bmatrix}$.

Solution:

Theorem:

If \mathbf{A} is an invertible matrix, k is a positive integer and c is a scalar not equal to zero. Then \mathbf{A}^{-1} , \mathbf{A}^k , $c\mathbf{A}$ and \mathbf{A}^T are invertible and the following statements are true:

- a) $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$.
- b) $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$.
- c) $(\mathbf{A}^k)^{-1} = (\mathbf{A}^{-1})^k$.
- d) $(c\mathbf{A})^{-1} = \frac{1}{c}\mathbf{A}^{-1}$.

Example

Compute \mathbf{A}^{-2} in two different ways for $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 2 & 4 \end{bmatrix}$.

Solution: $\mathbf{A}^{-2} = \begin{bmatrix} \frac{9}{2} & -\frac{5}{4} \\ -\frac{5}{2} & \frac{3}{4} \end{bmatrix}$.

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - **The inverse of a product**
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Theorem:

If A and B are invertible matrices of same size then AB is invertible and we have

Theorem:

If A and B are invertible matrices of same size then AB is invertible and we have

$$(AB)^{-1} = B^{-1}A^{-1}.$$

Theorem:

If A and B are invertible matrices of same size then AB is invertible and we have

$$(AB)^{-1} = B^{-1}A^{-1}.$$

Example

Find the inverse of AB with

$$A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 3 \end{bmatrix}.$$

Solution:

Theorem:

If A and B are invertible matrices of same size then AB is invertible and we have

$$(AB)^{-1} = B^{-1}A^{-1}.$$

Example

Find the inverse of AB with

$$A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 3 \end{bmatrix}.$$

Solution: $A^{-1} = \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$ and $B^{-1} = \begin{bmatrix} 1 & -2 & 1 \\ -1 & 1 & 0 \\ \frac{2}{3} & 0 & -\frac{1}{3} \end{bmatrix}.$

Example

$$\text{Hence } (AB)^{-1} = B^{-1}A^{-1} = \begin{bmatrix} 8 & -5 & -2 \\ -8 & 4 & 3 \\ 5 & -2 & -\frac{7}{3} \end{bmatrix}.$$

Example

$$\text{Hence } (AB)^{-1} = B^{-1}A^{-1} = \begin{bmatrix} 8 & -5 & -2 \\ -8 & 4 & 3 \\ 5 & -2 & -\frac{7}{3} \end{bmatrix}.$$

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Theorem:

If C is an invertible matrix, then the following statements hold:

a) If $AC = BC \implies A = B$.

b) If $CA = CB \implies A = B$.

Example

Solve the following matrix equation $AX + 2B = B^T$ where

$$A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 3 \end{bmatrix}$$

Solution

$$X = A^{-1}(B^T - 2B)$$

so

$$X = \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \left(\begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 4 \\ 3 & 3 & 3 \end{bmatrix} - 2 \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 3 \end{bmatrix} \right) = \dots$$

- 1 Homogeneous Systems
- 2 Supplementary Exercises on RREF - GJE
- 3 Inverse of a Matrix
- 4 Find the Inverse of a Matrix
 - 2×2 Matrix
 - $n \times n$ matrix
- 5 Properties of Inverse of a Matrix
 - General Properties
 - The inverse of a product
 - Cancellation Property
- 6 Solving a system of Linear Equation using Inverse of Matrices

Theorem:

The system $AX = b$ of n linear equations in n unknowns has a unique solution if and only if A is **Invertible**. In this case, the solution X is given by $X = A^{-1}b$.

Theorem:

The system $\mathbf{AX} = \mathbf{b}$ of n linear equations in n unknowns has a unique solution if and only if \mathbf{A} is **Invertible**. In this case, the solution \mathbf{X} is given by $\mathbf{X} = \mathbf{A}^{-1}\mathbf{b}$.

Example (Solve the system using the inverse of Matrices)

a)

$$(S) \begin{cases} x_1 & +4x_2 & +3x_3 & = & 12, \\ -x_1 & -2x_2 & & = & -12, \\ 2x_1 & +2x_2 & +3x_3 & = & 8. \end{cases}$$

Solution:

Theorem:

The system $\mathbf{AX} = \mathbf{b}$ of n linear equations in n unknowns has a unique solution if and only if \mathbf{A} is **Invertible**. In this case, the solution \mathbf{X} is given by $\mathbf{X} = \mathbf{A}^{-1}\mathbf{b}$.

Example (Solve the system using the inverse of Matrices)

a)

$$(S) \begin{cases} x_1 & +4x_2 & +3x_3 & = & 12, \\ -x_1 & -2x_2 & & = & -12, \\ 2x_1 & +2x_2 & +3x_3 & = & 8. \end{cases}$$

Solution: System (S) can be written as $\mathbf{AX} = \mathbf{b}$ with

Theorem:

The system $\mathbf{AX} = \mathbf{b}$ of n linear equations in n unknowns has a unique solution if and only if \mathbf{A} is **Invertible**. In this case, the solution \mathbf{X} is given by $\mathbf{X} = \mathbf{A}^{-1}\mathbf{b}$.

Example (Solve the system using the inverse of Matrices)

a)

$$(S) \begin{cases} x_1 + 4x_2 + 3x_3 = 12, \\ -x_1 - 2x_2 = -12, \\ 2x_1 + 2x_2 + 3x_3 = 8. \end{cases}$$

Solution: System (S) can be written as $\mathbf{AX} = \mathbf{b}$ with $\mathbf{A} = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$,

Theorem:

The system $\mathbf{AX} = \mathbf{b}$ of n linear equations in n unknowns has a unique solution if and only if \mathbf{A} is **Invertible**. In this case, the solution \mathbf{X} is given by $\mathbf{X} = \mathbf{A}^{-1}\mathbf{b}$.

Example (Solve the system using the inverse of Matrices)

a)

$$(S) \begin{cases} x_1 + 4x_2 + 3x_3 = 12, \\ -x_1 - 2x_2 = -12, \\ 2x_1 + 2x_2 + 3x_3 = 8. \end{cases}$$

Solution: System (S) can be written as $\mathbf{AX} = \mathbf{b}$ with $\mathbf{A} = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$, $\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$

Theorem:

The system $\mathbf{AX} = \mathbf{b}$ of n linear equations in n unknowns has a unique solution if and only if \mathbf{A} is **Invertible**. In this case, the solution \mathbf{X} is given by $\mathbf{X} = \mathbf{A}^{-1}\mathbf{b}$.

Example (Solve the system using the inverse of Matrices)

a)

$$(S) \begin{cases} x_1 + 4x_2 + 3x_3 = 12, \\ -x_1 - 2x_2 = -12, \\ 2x_1 + 2x_2 + 3x_3 = 8. \end{cases}$$

Solution: System (S) can be written as $\mathbf{AX} = \mathbf{b}$ with $\mathbf{A} = \begin{bmatrix} 1 & 4 & 3 \\ -1 & -2 & 0 \\ 2 & 2 & 3 \end{bmatrix}$, $\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$

and $\mathbf{b} = \begin{bmatrix} 12 \\ -12 \\ 8 \end{bmatrix}$.

Example (Solve the system using the inverse of Matrices)

In the previous example, we have find that the inverse of **A** is given by

$$\mathbf{A}^{-1} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix}.$$

Example (Solve the system using the inverse of Matrices)

In the previous example, we have find that the inverse of **A** is given by

$$\mathbf{A}^{-1} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix}.$$

Then, system (**S**) has a unique solution given by

Example (Solve the system using the inverse of Matrices)

In the previous example, we have find that the inverse of A is given by

$$A^{-1} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix}.$$

Then, system (S) has a unique solution given by

$$X =$$

Example (Solve the system using the inverse of Matrices)

In the previous example, we have find that the inverse of **A** is given by

$$\mathbf{A}^{-1} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix}.$$

Then, system (**S**) has a unique solution given by

$$\mathbf{X} = \mathbf{A}^{-1}\mathbf{b} =$$

Example (Solve the system using the inverse of Matrices)

In the previous example, we have find that the inverse of **A** is given by

$$\mathbf{A}^{-1} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix}.$$

Then, system (**S**) has a unique solution given by

$$\mathbf{X} = \mathbf{A}^{-1}\mathbf{b} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} 12 \\ -12 \\ 8 \end{bmatrix} =$$

Example (Solve the system using the inverse of Matrices)

In the previous example, we have find that the inverse of **A** is given by

$$\mathbf{A}^{-1} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix}.$$

Then, system (**S**) has a unique solution given by

$$\mathbf{X} = \mathbf{A}^{-1}\mathbf{b} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} 12 \\ -12 \\ 8 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ -\frac{8}{3} \end{bmatrix}.$$

Example (Solve the system using the inverse of Matrices)

In the previous example, we have find that the inverse of **A** is given by

$$\mathbf{A}^{-1} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix}.$$

Then, system (**S**) has a unique solution given by

$$\mathbf{X} = \mathbf{A}^{-1}\mathbf{b} = \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} 12 \\ -12 \\ 8 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ -\frac{8}{3} \end{bmatrix}.$$

Example (Solve the system using the inverse of Matrices)

Hence, $x_1 = 4$, $x_2 = 4$ and $x_3 = -\frac{8}{3}$ is the solution of system (S).

Example (Solve the system using the inverse of Matrices)

Hence, $x_1 = 4$, $x_2 = 4$ and $x_3 = -\frac{8}{3}$ is the solution of system (S).

b)

$$(S) \begin{cases} 2x & +3y & +z & = & -1, \\ 3x & +3y & +z & = & 1, \\ 2x & +4y & +z & = & -2. \end{cases}$$

Solution:

Example (Solve the system using the inverse of Matrices)

Hence, $x_1 = 4$, $x_2 = 4$ and $x_3 = -\frac{8}{3}$ is the solution of system (S).

b)

$$(S) \begin{cases} 2x + 3y + z = -1, \\ 3x + 3y + z = 1, \\ 2x + 4y + z = -2. \end{cases}$$

Solution:Homework.

c)

$$(S) \begin{cases} 2x + 3y + z = 4, \\ 3x + 3y + z = 8, \\ 2x + 4y + z = 5. \end{cases}$$

Solution:

Example (Solve the system using the inverse of Matrices)

Hence, $x_1 = 4$, $x_2 = 4$ and $x_3 = -\frac{8}{3}$ is the solution of system (S).

b)

$$(S) \begin{cases} 2x & +3y & +z & = & -1, \\ 3x & +3y & +z & = & 1, \\ 2x & +4y & +z & = & -2. \end{cases}$$

Solution:Homework.

c)

$$(S) \begin{cases} 2x & +3y & +z & = & 4, \\ 3x & +3y & +z & = & 8, \\ 2x & +4y & +z & = & 5. \end{cases}$$

Solution:Homework.