


Chapter 1: Linear Systems and Matrices

Section: Introduction to Linear Systems

Lecture #1

Lebanese University

Prof. Ali WEHBE



1 Systems of Linear Equations

- Linear Equation
- Solutions of a Linear Equation
- System of Linear Equations
- Solution of a System of Linear Equation
- Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

1 Systems of Linear Equations

- Linear Equation
- Solutions of a Linear Equation
- System of Linear Equations
- Solution of a System of Linear Equation
- Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

1 Systems of Linear Equations

- Linear Equation
 - Solutions of a Linear Equation
 - System of Linear Equations
 - Solution of a System of Linear Equation
 - Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

Linear Equation

Definition (Linear Equation)

A **linear equation** in n **unknowns** (or **variables**) is an equation of the form

$$a_1 x_1 + a_2 x_2 + \dots + a_n x_n = b,$$

where

$$\left\{ \begin{array}{ll} x_1, x_2, \dots, x_n : & \text{are the } \mathbf{unknowns} \text{ or the } \mathbf{variables}, \\ a_1, a_2, \dots, a_n : & \text{The } \mathbf{coefficients}, \text{ are real numbers,} \\ b : & \text{The } \mathbf{constant term}, \text{ is a real number,} \\ a_1 : & \text{The } \mathbf{leading coefficient}, \\ x_1 : & \text{The } \mathbf{leading variable}. \end{array} \right.$$

Remark

Linear Equations have no products ($x_1 x_2$, x_1^2), no roots (\sqrt{x} , $\sqrt[3]{x}$), no trigonometric ($\cos(x)$), no exponential (e^x), and no logarithmic functions ($\ln(x)$) for the variables. Variables appear only in first power.

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Variables:

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients:

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

b) $\sin(2)x_1 + \cos(3)x_2 - 2 \cot(3)x_3 + \frac{1}{\sin(1)}x_4 = 3$

Example (Linear Equations and Nonlinear Equations)

- a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

- b) $\sin(2)x_1 + \cos(3)x_2 - 2 \cot(3)x_3 + \frac{1}{\sin(1)}x_4 = 3$ is a Linear Equation.

Variables: x_1 , x_2 , x_3 and x_4 .

Coefficients: $\sin(2)$, $\cos(3)$, $-2 \cot(3)$ and $\frac{1}{\sin(1)}$.

The constant term: 3 .

Example (Linear Equations and Nonlinear Equations)

- a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

- b) $\sin(2)x_1 + \cos(3)x_2 - 2\cot(3)x_3 + \frac{1}{\sin(1)}x_4 = 3$ is a Linear Equation.

Variables: x_1 , x_2 , x_3 and x_4 .

Coefficients: $\sin(2)$, $\cos(3)$, $-2\cot(3)$ and $\frac{1}{\sin(1)}$.

The constant term: 3 .

- c) $2x + 3y - 4e^z + \ln(t) = -1$

Example (Linear Equations and Nonlinear Equations)

- a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

- b) $\sin(2)x_1 + \cos(3)x_2 - 2\cot(3)x_3 + \frac{1}{\sin(1)}x_4 = 3$ is a Linear Equation.

Variables: x_1 , x_2 , x_3 and x_4 .

Coefficients: $\sin(2)$, $\cos(3)$, $-2\cot(3)$ and $\frac{1}{\sin(1)}$.

The constant term: 3 .

- c) $2x + 3y - 4e^z + \ln(t) = -1$ is not a Linear Equation. Its a Nonlinear one.

Example (Linear Equations and Nonlinear Equations)

- a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

- b) $\sin(2)x_1 + \cos(3)x_2 - 2 \cot(3)x_3 + \frac{1}{\sin(1)}x_4 = 3$ is a Linear Equation.

Variables: x_1 , x_2 , x_3 and x_4 .

Coefficients: $\sin(2)$, $\cos(3)$, $-2 \cot(3)$ and $\frac{1}{\sin(1)}$.

The constant term: 3 .

- c) $2x + 3y - 4e^z + \ln(t) = -1$ is not a Linear Equation. Its a Nonlinear one.

Example (Linear Equations and Nonlinear Equations)

a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

b) $\sin(2)x_1 + \cos(3)x_2 - 2\cot(3)x_3 + \frac{1}{\sin(1)}x_4 = 3$ is a Linear Equation.

Variables: x_1, x_2, x_3 and x_4 .

Coefficients: $\sin(2)$, $\cos(3)$, $-2\cot(3)$ and $\frac{1}{\sin(1)}$.

The constant term: 3 .

c) $2x + 3y - 4e^z + \ln(t) = -1$ is not a Linear Equation. Its a Nonlinear one.

d) $xy + 2z \cos(t) = 2$, is not a Linear Equations. Its a Nonlinear one.

Example (Linear Equations and Nonlinear Equations)

- a) $x + 3y - z = 2$ is a Linear Equation.

Variables: x , y and z .

Coefficients: 1 , 3 and -1 .

The constant term: $b = 2$.

- b) $\sin(2)x_1 + \cos(3)x_2 - 2 \cot(3)x_3 + \frac{1}{\sin(1)}x_4 = 3$ is a Linear Equation.

Variables: x_1 , x_2 , x_3 and x_4 .

Coefficients: $\sin(2)$, $\cos(3)$, $-2 \cot(3)$ and $\frac{1}{\sin(1)}$.

The constant term: 3 .

- c) $2x + 3y - 4e^z + \ln(t) = -1$ is not a Linear Equation. Its a Nonlinear one.

- d) $xy + 2z \cos(t) = 2$, is not a Linear Equations. Its a Nonlinear one.

1 Systems of Linear Equations

- Linear Equation
- **Solutions of a Linear Equation**
- System of Linear Equations
- Solution of a System of Linear Equation
- Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

Definition

A solution of a linear equation is the number or set of numbers that when substituted in the equation for the variables, the equality holds.

Definition

A solution of a linear equation is the number or set of numbers that when substituted in the equation for the variables, the equality holds.

Example

Solve the following linear equations:

① $2x + 3 = 4$.

② $2x_1 + 3x_2 = 4$.

③ $2x - 3y + z = 5$.

Definition

A solution of a linear equation is the number or set of numbers that when substituted in the equation for the variables, the equality holds.

Example

Solve the following linear equations:

① $2x + 3 = 4$.

② $2x_1 + 3x_2 = 4$.

③ $2x - 3y + z = 5$.

Solution:

Definition

A solution of a linear equation is the number or set of numbers that when substituted in the equation for the variables, the equality holds.

Example

Solve the following linear equations:

① $2x + 3 = 4$.

② $2x_1 + 3x_2 = 4$.

③ $2x - 3y + z = 5$.

Solution:

① We have $2x + 3 = 4 \Rightarrow 2x = 4 - 3 = 1 \Rightarrow x = 1/2$, so

$$S = \left\{ \frac{1}{2} \right\}.$$

Solution:

② We can rewrite the L. E. as $x_1 = \frac{4-3x_2}{2}$ or $x_2 = \frac{4-2x_1}{3}$.

Solution:

- ② We can rewrite the L. E. as $x_1 = \frac{4-3x_2}{2}$ or $x_2 = \frac{4-2x_1}{3}$.
If we choose the first one.

Solution:

- ② We can rewrite the L. E. as $x_1 = \frac{4-3x_2}{2}$ or $x_2 = \frac{4-2x_1}{3}$.
If we choose the first one. We fix $x_2 = t$ where $t \in \mathbb{R}$.

Solution:

- ② We can rewrite the L. E. as $x_1 = \frac{4-3x_2}{2}$ or $x_2 = \frac{4-2x_1}{3}$.

If we choose the first one. We fix $x_2 = t$ where $t \in \mathbb{R}$. Then the solutions of the L.E is given by

$$\begin{cases} x_1 = \frac{4-3t}{2}, \\ x_2 = t. \end{cases}$$

thus the solution is: is given by

$$S = \left\{ x_1 = \frac{4-3t}{2}, x_2 = t, \text{ where } t \in \mathbb{R} \right\}.$$

Solution:

- ② We can rewrite the L. E. as $x_1 = \frac{4-3x_2}{2}$ or $x_2 = \frac{4-2x_1}{3}$.

If we choose the first one. We fix $x_2 = t$ where $t \in \mathbb{R}$. Then the solutions of the L.E is given by

$$\begin{cases} x_1 = \frac{4-3t}{2}, \\ x_2 = t. \end{cases}$$

thus the solution is: is given by

$$S = \left\{ x_1 = \frac{4-3t}{2}, x_2 = t, \text{ where } t \in \mathbb{R} \right\}.$$

This is called the set of **the parametric solution of the Linear Equation**.

An example for a solution of the L.E., if we take $t = 2$, then $x_1 = -1$ and $x_2 = 2$ is a solution.

Solution:

- ② We can rewrite the L. E. as $x_1 = \frac{4-3x_2}{2}$ or $x_2 = \frac{4-2x_1}{3}$.

If we choose the first one. We fix $x_2 = t$ where $t \in \mathbb{R}$. Then the solutions of the L.E is given by

$$\begin{cases} x_1 = \frac{4-3t}{2}, \\ x_2 = t. \end{cases}$$

thus the solution is: is given by

$$S = \left\{ x_1 = \frac{4-3t}{2}, x_2 = t, \text{ where } t \in \mathbb{R} \right\}.$$

This is called the set of **the parametric solution of the Linear Equation**.

An example for a solution of the L.E., if we take $t = 2$, then $x_1 = -1$ and $x_2 = 2$ is a solution.

Solution:

$$\textcircled{3} \quad 2x - 3y + z = 5.$$

1 Systems of Linear Equations

- Linear Equation
- Solutions of a Linear Equation
- **System of Linear Equations**
- Solution of a System of Linear Equation
- Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

Definition (System of Linear Equations)

Definition (System of Linear Equations)

A **linear system** of m equations in n unknowns (or variables) is given as the following one:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m, \end{cases}$$

where:

- a_{ij} (for $i = 1, \dots, m$ and $j = 1, \dots, n$): Is the **coefficient** of the **variable** x_j in the i^{th} equation number, is a real number,
- b_i (for $i = 1, \dots, m$): Is the **constant term** of the i^{th} equation.

1 Systems of Linear Equations

- Linear Equation
- Solutions of a Linear Equation
- System of Linear Equations
- **Solution of a System of Linear Equation**
- Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

Consistent and Inconsistent systems

Definition

If a system of Linear equations has **no solution**, we say that the system is **inconsistent**. If the system has **at least one solution**, we say that it is **consistent**.

Consistent and Inconsistent systems

Definition

If a system of Linear equations has **no solution**, we say that the system is **inconsistent**. If the system has **at least one solution**, we say that it is **consistent**.

Our aim is to find a solution or the set of solutions of the system of Linear Equation. To this end, we must answer to the following questions:

Consistent and Inconsistent systems

Definition

If a system of Linear equations has **no solution**, we say that the system is **inconsistent**. If the system has **at least one solution**, we say that it is **consistent**.

Our aim is to find a solution or the set of solutions of the system of Linear Equation. To this end, we must answer to the following questions:

- a) Which are the operations that we can applied to obtain the solution of the System of L. E.?

Consistent and Inconsistent systems

Definition

If a system of Linear equations has **no solution**, we say that the system is **inconsistent**. If the system has **at least one solution**, we say that it is **consistent**.

Our aim is to find a solution or the set of solutions of the system of Linear Equation. To this end, we must answer to the following questions:

- a) Which are the operations that we can applied to obtain the solution of the System of L. E.?
- b) When can we say that two systems of Linear Equations are Equivalent?

An example to deduce the operations

Solve the following systems:

a)

$$(S_1) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, \\ x_2 = 3, \\ 2x_3 = 4. \end{cases}$$

b)

$$(S_2) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, \\ -3x_1 - x_2 + x_3 = 5, \\ 3x_1 + 2x_2 + x_3 = 2. \end{cases}$$

An example to deduce the operations

Solution

a)

$$(S_1) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, \\ x_2 = 3, \\ 2x_3 = 4. \end{cases}$$

An example to deduce the operations

Solution

a)

$$(S_1) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, & (1) \\ x_2 = 3, & (2) \\ 2x_3 = 4. & (3) \end{cases}$$

Equation (3) implies that $x_3 = 2$.

An example to deduce the operations

Solution

a)

$$(S_1) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, & (1) \\ x_2 = 3, & (2) \\ 2x_3 = 4. & (3) \end{cases}$$

Equation (3) implies that $x_3 = 2$. Then **substituting** equation (2) and the result of equation (3) in equation (1), we get $x_1 = -2$.

An example to deduce the operations

Solution

a)

$$(S_1) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, & (1) \\ x_2 = 3, & (2) \\ 2x_3 = 4. & (3) \end{cases}$$

Equation (3) implies that $x_3 = 2$. Then **substituting** equation (2) and the result of equation (3) in equation (1), we get $x_1 = -2$. Hence, the solution of system (S_1) is given by

$$\begin{cases} x_1 = -2, \\ x_2 = 3, \\ x_3 = 2. \end{cases}$$

An example to deduce the operations

Solution

a)

$$(S_1) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, & (1) \\ x_2 = 3, & (2) \\ 2x_3 = 4. & (3) \end{cases}$$

Equation (2) implies that $x_3 = 2$. Then **substituting** equation (2) and the result of equation (3) in equation (1), we get $x_1 = -2$. Hence, the solution of system (S_1) is given by

$$\begin{cases} x_1 = -2, \\ x_2 = 3, \\ x_3 = 2. \end{cases}$$

Operation is **Back Substitution**

An example to deduce the operations

b)

$$(S_2) \begin{cases} 3x_1 & +2x_2 & -x_3 & = & -2, \\ -3x_1 & -x_2 & +x_3 & = & 5, \\ 3x_1 & +2x_2 & +x_3 & = & 2. \end{cases}$$

An example to deduce the operations

b)

$$(S_2) \begin{cases} 3x_1 & +2x_2 & -x_3 & = & -2, & (1) \\ -3x_1 & -x_2 & +x_3 & = & 5, & (2) \\ 3x_1 & +2x_2 & +x_3 & = & 2. & (3) \end{cases}$$

Eliminating x_1 from the second equation: $(2) + (1) \Rightarrow (2')$

$$(S'_2) \begin{cases} 3x_1 & +2x_2 & -x_3 & = & -2, & (1) \\ & x_2 & & = & 3, & (2') \\ 3x_1 & +2x_2 & +x_3 & = & 2. & (3) \end{cases}$$

Eliminating x_1 and x_2 from the third equation:

An example to deduce the operations

$$(3) - (1) \Rightarrow (3')$$

$$(S'_2) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, & (1) \\ x_2 = 3, & (2') \\ 2x_3 = 4. & (3') \end{cases}$$

An example to deduce the operations

$$(3) - (1) \Rightarrow (3')$$

$$(S'_2) \begin{cases} 3x_1 + 2x_2 - x_3 = -2, & (1) \\ x_2 = 3, & (2) \\ 2x_3 = 4. & (3') \end{cases}$$

We remark that the system (S'_2) is the system (S_1) . Thus, system (S'_2) has the same solution of (S_1) . Equivalently, system (S_2) has the same solution.

1 Systems of Linear Equations

- Linear Equation
- Solutions of a Linear Equation
- System of Linear Equations
- Solution of a System of Linear Equation
- Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

Summary

Definition (Equivalent Systems)

Two systems of L.E. are said to be **equivalent** if they have the same solution set.

Summary

Definition (Equivalent Systems)

Two systems of L.E. are said to be **equivalent** if they have the same solution set.

Strategy to obtain an Equivalent system

Summary

Definition (Equivalent Systems)

Two systems of L.E. are said to be **equivalent** if they have the same solution set.

Strategy to obtain an Equivalent system

There are three operations that can be used on a system to obtain an **equivalent** system.

Summary

Definition (Equivalent Systems)

Two systems of L.E. are said to be **equivalent** if they have the same solution set.

Strategy to obtain an Equivalent system

There are three operations that can be used on a system to obtain an **equivalent** system. These operations are named by **Elementary operations**:

Summary

Definition (Equivalent Systems)

Two systems of L.E. are said to be **equivalent** if they have the same solution set.

Strategy to obtain an Equivalent system

There are three operations that can be used on a system to obtain an **equivalent** system. These operations are named by **Elementary operations**:

- I- Interchange two equations.

Summary

Definition (Equivalent Systems)

Two systems of L.E. are said to be **equivalent** if they have the same solution set.

Strategy to obtain an Equivalent system

There are three operations that can be used on a system to obtain an **equivalent** system. These operations are named by **Elementary operations**:

- I- Interchange two equations.
- II- Multiply an equation by a nonzero real number.

Summary

Definition (Equivalent Systems)

Two systems of L.E. are said to be **equivalent** if they have the same solution set.

Strategy to obtain an Equivalent system

There are three operations that can be used on a system to obtain an **equivalent** system. These operations are named by **Elementary operations**:

- I- Interchange two equations.
- II- Multiply an equation by a nonzero real number.
- III- Replace an equation by its sum with a multiple of another equation.

1 Systems of Linear Equations

- Linear Equation
- Solutions of a Linear Equation
- System of Linear Equations
- Solution of a System of Linear Equation
- Equivalent systems and Elementary operations

2 Introduction to Matrices

3 Matrices

Solve the following system using the previous operations:

Solve the following system using the previous operations:

$$\left\{ \begin{array}{rrrrrrrrrr} -x_1 & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 1, \\ x_1 & & +x_3 & +42x_4 & +8x_5 & +4x_6 & -23x_7 & +7x_8 & = & 2, \\ & -2x_2 & +4x_3 & +4x_4 & +5x_5 & +6x_6 & -13x_7 & -x_8 & = & 3, \\ -x_1 & -3x_2 & & +4x_4 & +x_5 & -9x_6 & & -x_8 & = & 4, \\ 2x_1 & +9x_2 & +2x_3 & -x_4 & & -7x_6 & -2x_7 & -x_8 & = & 5, \\ 3x_1 & +8x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & & = & 6, \\ -2x_1 & +7x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & -9x_7 & -x_8 & = & 7, \\ & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 8, \\ 8x_1 & +3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +x_7 & -x_8 & = & 9, \\ -11x_1 & -3x_2 & -x_3 & & +5x_5 & & +11x_7 & -x_8 & = & -9. \end{array} \right.$$

Solve the following system using the previous operations:

$$\left\{ \begin{array}{rrrrrrrrrr} -x_1 & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 1, \\ x_1 & & +x_3 & +42x_4 & +8x_5 & +4x_6 & -23x_7 & +7x_8 & = & 2, \\ & -2x_2 & +4x_3 & +4x_4 & +5x_5 & +6x_6 & -13x_7 & -x_8 & = & 3, \\ -x_1 & -3x_2 & & +4x_4 & +x_5 & -9x_6 & & -x_8 & = & 4, \\ 2x_1 & +9x_2 & +2x_3 & -x_4 & & -7x_6 & -2x_7 & -x_8 & = & 5, \\ 3x_1 & +8x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & & = & 6, \\ -2x_1 & +7x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & -9x_7 & -x_8 & = & 7, \\ & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 8, \\ 8x_1 & +3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +x_7 & -x_8 & = & 9, \\ -11x_1 & -3x_2 & -x_3 & & +5x_5 & & +11x_7 & -x_8 & = & -9. \end{array} \right.$$

Solution:

Solve the following system using the previous operations:

$$\left\{ \begin{array}{rrrrrrrrrr} -x_1 & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 1, \\ x_1 & & +x_3 & +42x_4 & +8x_5 & +4x_6 & -23x_7 & +7x_8 & = & 2, \\ & -2x_2 & +4x_3 & +4x_4 & +5x_5 & +6x_6 & -13x_7 & -x_8 & = & 3, \\ -x_1 & -3x_2 & & +4x_4 & +x_5 & -9x_6 & & -x_8 & = & 4, \\ 2x_1 & +9x_2 & +2x_3 & -x_4 & & -7x_6 & -2x_7 & -x_8 & = & 5, \\ 3x_1 & +8x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & & = & 6, \\ -2x_1 & +7x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & -9x_7 & -x_8 & = & 7, \\ & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 8, \\ 8x_1 & +3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +x_7 & -x_8 & = & 9, \\ -11x_1 & -3x_2 & -x_3 & & +5x_5 & & +11x_7 & -x_8 & = & -9. \end{array} \right.$$

Solution:

????????????????

Logically, it is hard to apply the Elementary operations in order to solve the previous problem since it contains 10 equations and 8 variables. We must use another tools. The most important tools to solve these type of systems are the

Matrices

From a system of Linear equations to a matrix

$$\left\{ \begin{array}{rrrrrrrrrr} -x_1 & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 1, \\ x_1 & & +x_3 & +42x_4 & +8x_5 & +4x_6 & -23x_7 & +7x_8 & = & 2, \\ & -2x_2 & +4x_3 & +4x_4 & +5x_5 & +6x_6 & -13x_7 & -x_8 & = & 3, \\ -x_1 & -3x_2 & & +4x_4 & +x_5 & -9x_6 & & -x_8 & = & 4, \\ 2x_1 & +9x_2 & +2x_3 & -x_4 & & -7x_6 & -2x_7 & -x_8 & = & 5, \\ 3x_1 & +8x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & & = & 6, \\ -2x_1 & +7x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & -9x_7 & -x_8 & = & 7, \\ & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 & = & 8, \\ 8x_1 & +3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +x_7 & -x_8 & = & 9, \\ -11x_1 & -3x_2 & -x_3 & & +5x_5 & & +11x_7 & -x_8 & = & -9, \end{array} \right.$$

From a system of Linear equations to a matrix

We see the coefficients of every **variable** in each **equation**

$$\left\{ \begin{array}{cccccccc} -x_1 & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 = & 1, \\ 1x_1 & & +x_3 & +42x_4 & +8x_5 & +4x_6 & -23x_7 & +7x_8 = & 2, \\ 0x_1 & -2x_2 & +4x_3 & +4x_4 & +5x_5 & +6x_6 & -13x_7 & -x_8 = & 3, \\ -x_1 & -3x_2 & & +4x_4 & +x_5 & -9x_6 & & -x_8 = & 4, \\ 2x_1 & +9x_2 & +2x_3 & -x_4 & & -7x_6 & -2x_7 & -x_8 = & 5, \\ 3x_1 & +8x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & = & 6, \\ -2x_1 & +7x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & -9x_7 & -x_8 = & 7, \\ 0x_1 & -3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +11x_7 & -x_8 = & 8, \\ 8x_1 & +3x_2 & +2x_3 & +4x_4 & +5x_5 & +6x_6 & +x_7 & -x_8 = & 9, \\ -11x_1 & -3x_2 & -x_3 & & +5x_5 & & +11x_7 & -x_8 = & -9, \end{array} \right.$$

For example, the **variable** x_1 has the following coefficients: $-1, 1, 0, -1, +2, +3, -2, 0, +8$ and -11 .

From a system of Linear equations to a matrix

We use the coefficients of the variables in each equation and we construct the following object:

$$\begin{bmatrix} -1 & -3 & 2 & 4 & 5 & 6 & 11 & -1 \\ 1 & 0 & 1 & 42 & 8 & 4 & -23 & 7 \\ 0 & -2 & 4 & 4 & 5 & 6 & -13 & -1 \\ -1 & -3 & 0 & 4 & 1 & -9 & 0 & -1 \\ 2 & 9 & 2 & -1 & 0 & -7 & -2 & -1 \\ 3 & 8 & 2 & 4 & 5 & 6 & 11 & 0 \\ -2 & 7 & 2 & 4 & 5 & 6 & -9 & -1 \\ 0 & -3 & 2 & 4 & 5 & 6 & 11 & -1 \\ 8 & 3 & 2 & 4 & 5 & 6 & 1 & -1 \\ -11 & -3 & -1 & 0 & 5 & 0 & 11 & -1 \end{bmatrix}$$

From a system of Linear equations to a matrix

We use the coefficients of the variables in each equation and we construct the following object:

$$\begin{bmatrix} -1 & -3 & 2 & 4 & 5 & 6 & 11 & -1 \\ 1 & 0 & 1 & 42 & 8 & 4 & -23 & 7 \\ 0 & -2 & 4 & 4 & 5 & 6 & -13 & -1 \\ -1 & -3 & 0 & 4 & 1 & -9 & 0 & -1 \\ 2 & 9 & 2 & -1 & 0 & -7 & -2 & -1 \\ 3 & 8 & 2 & 4 & 5 & 6 & 11 & 0 \\ -2 & 7 & 2 & 4 & 5 & 6 & -9 & -1 \\ 0 & -3 & 2 & 4 & 5 & 6 & 11 & -1 \\ 8 & 3 & 2 & 4 & 5 & 6 & 1 & -1 \\ -11 & -3 & -1 & 0 & 5 & 0 & 11 & -1 \end{bmatrix} \Longleftarrow \text{Matrix}$$

How can we characterize a Matrix?

We can characterize a matrix by its number of rows and columns.

How can we characterize a Matrix?

We can characterize a matrix by its number of rows and columns.

$$\begin{bmatrix} -1 & -3 & 2 & 4 & 5 & 6 & 11 & -1 \\ 1 & 0 & 1 & 42 & 8 & 4 & -23 & 7 \\ 0 & -2 & 4 & 4 & 5 & 6 & -13 & -1 \\ -1 & -3 & 0 & 4 & 1 & -9 & 0 & -1 \\ 2 & 9 & 2 & -1 & 0 & -7 & -2 & -1 \\ 3 & 8 & 2 & 4 & 5 & 6 & 11 & 0 \\ -2 & 7 & 2 & 4 & 5 & 6 & -9 & -1 \\ 0 & -3 & 2 & 4 & 5 & 6 & 11 & -1 \\ 8 & 3 & 2 & 4 & 5 & 6 & 1 & -1 \\ -11 & -3 & -1 & 0 & 5 & 0 & 11 & -1 \end{bmatrix}$$

Number of row is **10**=number of equations of the system, and number of columns is **8**=number of variables of the system.

- 1 Systems of Linear Equations
 - Linear Equation
 - Solutions of a Linear Equation
 - System of Linear Equations
 - Solution of a System of Linear Equation
 - Equivalent systems and Elementary operations

- 2 Introduction to Matrices

- 3 Matrices

Definition (Matrix Notation)

A matrix \mathbf{A} of m rows and n columns is given as follows:

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \cdot & \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{bmatrix}$$

where a_{ij} is a real number (constant) that represent the entry of the matrix \mathbf{A} , for $i = 1, \dots, m$ and $j = 1, \dots, n$. In this case, we say that \mathbf{A} is an $m \times n$ matrix and we denote $\mathbf{A} := [a_{ij}]$.

Example

Give all the characterization of the following matrix

$$A = \begin{bmatrix} 1 & 2 & -7 \\ 3 & 0 & 6 \end{bmatrix}$$

Example

Give all the characterization of the following matrix

$$A = \begin{bmatrix} 1 & 2 & -7 \\ 3 & 0 & 6 \end{bmatrix}$$

Solution:

A is 2×3 matrix. $A = [a_{ij}]$, $i = 1, 2$ number of rows, $j = 1, 2, 3$ number of columns.