



# CS1231S

## TUTORIAL #11

---

### Graphs and Trees



# Learning objectives of this tutorial

## Graphs and Trees


- Introducing complement graphs and self-complementary graphs.
- Calculating the number of walks of a certain length in a graph.
- Modeling a problem as a graph problem.
- Understanding pre-order, in-order and post-order traversals.
- Calculating the number of spanning trees in a graph.
- Applying Kruskal's Algorithm and Prim's Algorithm.

# Definitions

If  $G$  is a simple graph, the **complement** of  $G$ , denoted  $\bar{G}$ , is obtained as follows: the vertex set of  $\bar{G}$  is identical to the vertex set of  $G$ . However, two distinct vertices  $v$  and  $w$  of  $\bar{G}$  are connected by an edge iff  $v$  and  $w$  are not connected by an edge in  $G$ .

A **self-complementary graph** is isomorphic with its complement.

A simple circuit (cycle) of length 3 is called a **triangle**.




A graph  $G$  and its complement  $\bar{G}$ .

Q1. Draw all **self-complementary graphs** with


(a) 4 vertices

How about this?




$\cong$



Complement graph




(b) 5 vertices



$\cong$




$\cong$



**Q2.** Show that every simple graph with at least 2 vertices has two vertices of the same degree.

1. Let  $G$  be a simple graph with  $n \geq 2$  vertices.
2. Case 1: If  $G$  has no vertex with degree  $n - 1$ 
  - 2.1 Then all vertices in  $G$  have degree lying in the range 0 to  $n - 2$  inclusive.
3. Case 2: If  $G$  has a vertex  $v$  with degree  $n - 1$ 
  - 3.1 Firstly,  $v$  does not have degree 0 (since  $n - 1 > 0$ ).
  - 3.2 Also, as  $v$  has degree  $n - 1$ , it is connected to every other vertex, hence no other vertex can have degree 0 too.
  - 3.3 Hence, all vertices in  $G$  have degree lying in the range 1 to  $n - 1$  inclusive.
4. In all cases, there are at most  $n - 1$  possible vertex degrees among the  $n$  vertices.
5. Therefore, at least two vertices must have the same degree (by the Pigeonhole Principle).

Q3. Given this graph



(a) Write the adjacency matrix  $A$  for the graph. Let the rows and columns be  $p, q, r, s$ .

$$A = \begin{bmatrix} p & q & r & s \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 2 & 1 & 0 \end{bmatrix} \begin{matrix} p \\ q \\ r \\ s \end{matrix}$$

(b) Find  $A^2$  and  $A^3$ .

$$A^2 = \begin{bmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 2 & 1 & 0 \end{bmatrix} \times \begin{bmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 2 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 2 & 2 & 2 \\ 2 & 6 & 3 & 2 \\ 2 & 3 & 3 & 3 \\ 2 & 2 & 3 & 6 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 2 & 1 & 0 \end{bmatrix} \times \begin{bmatrix} 2 & 2 & 2 & 2 \\ 2 & 6 & 3 & 2 \\ 2 & 3 & 3 & 3 \\ 2 & 2 & 3 & 6 \end{bmatrix} = \begin{bmatrix} 4 & 8 & 6 & 8 \\ 8 & 9 & 11 & 17 \\ 6 & 11 & 9 & 11 \\ 8 & 17 & 11 & 9 \end{bmatrix}$$

(c) How many walks of length 2 are there from  $p$  to  $q$ ? From  $s$  to itself?

2 walks from  $p$  to  $q$ :  $\langle pe_2se_5q \rangle, \langle pe_2se_6q \rangle$ .

6 walks from  $s$  to  $s$ :  $\langle se_2pe_2s \rangle, \langle se_4re_4s \rangle, \langle se_5qe_5s \rangle, \langle se_6qe_6s \rangle, \langle se_5qe_6s \rangle, \langle se_6qe_5s \rangle$ .

(d) How many walks of length 3 are there from  $r$  to  $s$ ? From  $s$  to  $p$ ?

11 walks from  $r$  to  $s$ :

2 ways via  $q$ :  $\langle re_3qe_1pe_2s \rangle, \langle re_3qe_3re_4s \rangle$ .

3 ways via  $r$ :  $\langle re_7re_7re_4s \rangle, \langle re_7re_3qe_5s \rangle, \langle re_7re_3qe_6s \rangle$ .

6 ways via  $s$ :  $\langle re_4se_2pe_2s \rangle, \langle re_4se_4re_4s \rangle, \langle re_4se_5qe_5s \rangle, \langle re_4se_6qe_6s \rangle, \langle re_4se_5qe_6s \rangle, \langle re_4se_6qe_5s \rangle$ .

8 walks from  $s$  to  $p$ :

2 ways via  $p$ :  $\langle se_2pe_2se_2p \rangle, \langle se_2pe_1qe_1p \rangle$ .

4 ways via  $q$ :  $\langle se_5qe_5se_2p \rangle, \langle se_5qe_6se_2p \rangle, \langle se_6qe_5se_2p \rangle, \langle se_6qe_6se_2p \rangle$ .

2 ways via  $r$ :  $\langle se_4re_4se_2p \rangle, \langle se_4re_3qe_1p \rangle$ .

# Q4.

Prove that for any simple graph  $G$  with 6 vertices,  $G$  or its complementary graph  $\bar{G}$  contains a triangle.

1. Take any simple graph  $G$  with 6 vertices (figure 4a).
2. Draw a black edge between adjacent vertices in  $G$  and draw a red edge between non-adjacent vertices in  $G$ .
3. Call this graph  $G'$  (figure 4b). Now  $G'$  is a complete graph with every edge either black or red, and we want to prove that it has a black triangle or a red triangle.
4. Let  $v$  be an arbitrary vertex of  $G'$ .
  - 4.1 There are 5 edges incident to  $v$ , which are either black or red.
  - 4.2 Therefore, (at least) 3 of these 5 edges are of the same colour  $c$  (by **Generalized PHP**).
  - 4.3 For these 3 edges of colour  $c$ , name the vertices at the other end of these edges  $u_1, u_2, u_3$  (figure 4c).
  - 4.4 Case 1: If there is an edge of colour  $c$  between any two of  $u_1, u_2, u_3$ , then that edge forms a triangle of colour  $c$  with the two edges coming from  $v$ . (In figure 4d, the 3 dashed lines are the edges between  $u_1, u_2, u_3$ . (In this example, two of them –  $\{u_1, u_3\}$  and  $\{u_2, u_3\}$  – are of colour  $c$  (red). Let's pick  $\{u_1, u_3\}$ . We have a triangle of colour  $c$  with  $v, u_1, u_3$ .)
  - 4.5 Case 2: If there are no edges of colour  $c$  between any pair of  $u_1, u_2, u_3$ . Then the edges between these 3 vertices form a triangle of colour opposite to  $c$ .
5. In all cases, there is a triangle of the same colour.




Figure 4a: Graph  $G$




Figure 4b: Graph  $G'$




Figure 4c




Figure 4d


Q5.

Given a pile of stones. At each step, you can separate a pile of  $k$  stones into two piles of  $k_1$  and  $k_2$  stones ( $k_1 + k_2 = k$ ). On doing this, you earn  $\$(k_1 \times k_2)$ . What is the maximum amount of money you can earn at the end if you start with a pile of  $n$  stone?



Model this as a graph problem. Let vertices represent the stones. Draw an edge between two vertices if the stones they represent are in the same pile.

In the beginning, we have a  $K_n$  graph.

Eg:  $K_5$



In the end, we have a graph of  $n$  isolated vertices.




Separating a pile of  $k$  stones into two piles of  $k_1$  and  $k_2$  stones corresponds to removing  $k_1 \times k_2$  edges from  $K_k$ .

A  $K_n$  graph has  $\binom{n}{2} = \frac{n(n-1)}{2}$  edges. Therefore, the maximum amount one can earn is  $\$ \frac{n(n-1)}{2}$ .



**Q6.** How many edges are there in a forest with  $v$  vertices and  $k$  components?



1. Let the  $i^{\text{th}}$  component of the forest contains  $v_i$  vertices.
2. Each component is a tree, therefore the  $i^{\text{th}}$  component has  $v_i - 1$  edges.  
(by **Theorem 10.5.2**)
3. Hence, the total number of edges in the forest is

$$\sum_{i=1}^k (v_i - 1) = \sum_{i=1}^k v_i - \sum_{i=1}^k 1 = \mathbf{v - k}.$$


**Theorem 10.5.2**

A tree with  $n$  vertices ( $n > 0$ ) has  $n - 1$  edges.


**Q7.** How many possible binary trees with 4 vertices  $A, B, C$  and  $D$  have this in-order traversal:  $A B C D$ ? Draw them.

**Total: 14**


Rooted at  $A$ : 5




Rooted at  $B$ : 2



Rooted at  $D$ : 5




Rooted at  $C$ : 2




Q8.

(a) The number of spanning trees in a complete graph  $K_n$  is  $n^{n-2}$ .  
Show all the spanning trees in  $K_4$  below.

$$4^{4-2} = 16$$



**Q8.** (b) Following the given steps, verify that the number of spanning trees in  $K_5$  is  $5^{5-3} = 125$ .



Adjacency matrix  $A$  for  $K_5$ :

$$A = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{pmatrix}$$

Diagonal matrix  $D$ :

$$D = \begin{pmatrix} 4 & 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 \\ 0 & 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 0 & 4 \end{pmatrix}$$


$$M = D - A = \begin{pmatrix} 4 & -1 & -1 & -1 & -1 \\ -1 & 4 & -1 & -1 & -1 \\ -1 & -1 & 4 & -1 & -1 \\ -1 & -1 & -1 & 4 & -1 \\ -1 & -1 & -1 & -1 & 4 \end{pmatrix}$$

$$M' = \begin{pmatrix} 4 & -1 & -1 & -1 \\ -1 & 4 & -1 & -1 \\ -1 & -1 & 4 & -1 \\ -1 & -1 & -1 & 4 \end{pmatrix}$$

$$\begin{aligned} \det(M') &= 4 \begin{vmatrix} 4 & -1 & -1 \\ -1 & 4 & -1 \\ -1 & -1 & 4 \end{vmatrix} + \begin{vmatrix} -1 & -1 & -1 \\ -1 & 4 & -1 \\ -1 & -1 & 4 \end{vmatrix} - \begin{vmatrix} -1 & 4 & -1 \\ -1 & -1 & -1 \\ -1 & -1 & 4 \end{vmatrix} + \begin{vmatrix} -1 & 4 & -1 \\ -1 & -1 & 4 \\ -1 & -1 & -1 \end{vmatrix} \\ &= 4(50) + (-25) - (25) + (-25) = \mathbf{125}. \end{aligned}$$

Q9. Find the MST of the graph below.

Kruskal's algorithm




Edges in non-decreasing order:

3  
4  
5  
6  
9  
12  
15  
16  
18  
21  
25  
28

Weight of MST =  $3 + 4 + 5 + 6 + 15 + 18 = 51$ .

Q9. Find the MST of the graph below.



## Prim's algorithm

Start from top vertex.

1.  $\{5, 16, 25\} \rightarrow 5$
2.  $\{25, 16, 15\} \rightarrow 15$
3.  $\{25, 28, 4, 3, 21\} \rightarrow 3$
4.  $\{25, 28, 4, 12, 9\} \rightarrow 4$
5.  $\{25, 28, 18, 6, 9\} \rightarrow 6$
6.  $\{25, 28, 18\} \rightarrow 18$


Weight of MST =  $5 + 15 + 3 + 4 + 6 + 18 = 51$ .

Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: IADJNHBEKOF LGCM

Pre-order: HNAIJD OBKECLFGM




Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: IADJN<sup>H</sup>BEKOF LGCM

Pre-order: <sup>H</sup>N<sub>A</sub> IJ DOBKECLFGM






Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: I A D J N H B E K O F L G C M

Pre-order: H N A I J D O B K E C L F G M




Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: TA D J N H B E K O F L G C M

Pre-order: H N A I J D O B K E C L F G M




Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: I A D J N H B E K O F L G C M

Pre-order: H N A I J D O B K E C L F G M




## Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: **I** **A** **D** **J** **N** **H** **B** **E** **K** **O** **F** **L** **G** **C** **M**

Pre-order: H N A I J D O B K E C L F G M




Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: I A D J N H B E K O F L G C M

Pre-order: H N A I J D O B K E C L F G M




Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: I A D J N H B E K O F L G C M

Pre-order: H N A I J D O B K E C L F G M




Repeat the process...

Q10.

Construct the binary tree give the following in-order and pre-order traversal of the tree:

In-order: I A D J N H B E K O F L G C M

Pre-order: H N A I J D O B K E C L F G M



Final binary tree

You  
I Can  
Do it

The text 'You I Can Do it' is written in a black, cursive, handwritten font. The words are arranged in three lines: 'You' on the top line, 'I Can' on the middle line, and 'Do it' on the bottom line. The text is surrounded by decorative elements: several yellow curved lines and dots, and many small pink dots, creating a celebratory or energetic feel. A faint 'dreamstime' watermark is visible across the middle of the text.



THE END