Looking at the average susceptibility values, Rwandex-Magerwa is overall the sub-catchment that is the most prone to shallow landslide initiation, while Rwabayanga is overall the sub-catchment that is the least prone. However, such average values are meaningless if the distribution patterns are not analysed.

Table 3: **Areas per landslide susceptibility class per site**

Susceptibility Class	Bishenyi		Rwabayanga		Rusizi		Rwandex-Magerwa	
	Area (km^2)	Percentage of total area	Area (km^2)	Percentage of total area	Area (km^2)	Percentage of total area	Area (km^2)	Percentage of total area
<= 0.2	9.87	20.9%	2.54	32.5%	9.50	45.1%	4.45	45.8%
0.2 - 0.45	14.85	31.4%	3.06	39.1%	6.55	31.1%	2.83	29.1%
0.45 - 0.55	12.47	26.4%	1.20	15.3%	3.31	15.7%	1.29	13.3%
0.55 - 0.8	7.45	15.8%	0.67	8.6%	1.45	6.9%	0.73	7.5%
0.8 - 1.0	2.62	5.5%	0.35	4.5%	0.25	1.2%	0.41	4.2%

Table 4: **Sub-catchment average values of landslide susceptibility**

Sub -catchment	Average landslide susceptibility
Bishenyi	0.37
Kamembe-Gihundwe	0.34
Rwandex-Magerwa	0.46
Rwabayanga	0.26

Table 5: **Sub-catchment average values of landslide frequency (mean landslide frequency and mean landslide affected area)**

Sub -catchment	Average frequency LS/year/km^2	Average affected area m^2/year/km^2
Bishenyi	0.023	16.40
Kamembe-Gihundwe	0.016	11.5
Rwandex-Magerwa	0.037	26.9
Rwabayanga	0.009	6.62