

a
Measured at both 12 mo & 24 mo:
• Anthropometry • Stool microbiome
• Plasma lipids, vitamins, amino acids
• Sleep • Perceived stress
Baseline / one-time (or no-timepoint):
• Parent education & economics • Family structure
• Household (roof, water, sanitation) • Media use
• Polygenic risk scores (PRS)
Baseline 12 mo
15 mo
Post-refeed 24 mo
Rapid catchup
Nutritional maintenance
Moderate acute malnutrition (MAM)
Enhanced Refeed arm
n = 80
ERUSF
ESQ-LNS
Local Refeed arm
n = 79
RUSF
SQ-LNS
Recovery = WLZ > -1
Well-nourished controls
n = 70
Covariates: Sex, Delivery mode, Place of birth, Days of exclusive breasfeeding
b
Malnutrition (WLZ)
Rapid catchup
Recovery = -1
MAM
Weeks after baseline (12 mo)
Recovery
No-recovery
Well-nourished
c
Percentage of participants
Recovery status
Sustained recovery
Delayed recovery
Unsustained recovery
No recovery
n = 9
n = 1
n = 25
n = 44
n = 8
n = 2
n = 25
n = 45
d
Time to recovery (days)
Cox
p = 0.724
Local
Enhanced
Refeed arm