

# **Thesis corrections for “Spatiotemporal modelling of all-cause and cause-specific mortality in England”**

Theo Alexander Omar Rashid  
2023-01-12

25th December, 2023

Dear examiners,

Thank you for taking the time to read my thesis “Spatiotemporal modelling of all-cause and cause-specific mortality in England” and for the thoughtful and enjoyable discussion we had during the viva. I attach my response to the corrections.

Kind regards,

Theo Rashid

## **Required corrections**

1. P25 (and P40): While not strictly a paper, a very classical depiction of inequalities in longevity across small areas actually comes from London  
(<https://journals.sagepub.com/doi/abs/10.1068/a45341>)
2. P28: please revise the textual distinction made between areal model using spatial neighbourhood structure and those which use nested hierarchy as they are both a specific case of hierarchical models
3. P31: Please discuss here or elsewhere that in some cases it is not appropriate to have statistical smoothing and how can someone evaluate if that is the case in a specific situation; we discussed using posterior predictive checks and abplots on death rates

4. P32: Here or elsewhere please also discuss potential disadvantages in life expectancy as an outcome measure especially as relevant to the assumptions that you need to make in terms of its calculation
5. P48 the idea of exploring population turnover (and therefore a measure of changing composition) is great. The only thing I'd mention is that turnover itself may be an exposure of ill health (some discussion here <https://www.sciencedirect.com/science/article/abs/pii/S1353829218310700>)
6. P54: An important topic not considered in detail is model adequacy. Please add a table summarizing outcomes of model adequacy and consistency checks performed.
7. P56: Please elaborate on your prior choices and robustness of your analyses.
8. P57: Please reconsider your statement that "INLA scales badly" in light of the discussion during the viva and our elaboration that the number of hyperparameters are probably less than 20.
9. P57: It would be useful to see a deeper discussion on the age-specific random walk priors, in light of the J-shape of all-cause mortality rates, and different age profiles of cause-specific mortality rates, as well as a discussion on independence across gender. For example, another way of modelling age (as compared to random walks) is through the use of linear splines (e.g., TOPALS): <https://www.scielo.br/ijrbepop/a/9szg7XYCXck9dJrKmgBSdrf/?lang=en> and the pros and cons would merit discussion.
10. Figure 5.1: what years are used for the country life expectancies? 2002 or 2019?

As stated in the figure caption, the life expectancy estimates "for other countries [are] from World Bank estimates in 2019".

11. P62: missing from this paper is an evaluation of the uncertainty around life expectancy. This could be measured with a sort of coefficient of variation (SD of all posterior LEs divided over the median LE). The CDC in the US has used 25% as a threshold for poor certainty in some of their papers, though this threshold may be too high.

12. P73: On the discussion about factors driving lower life expectancy in the US (linked to deaths of despair), a recent paper blames the lack of social safety net systems (<https://jamanetwork.com/journals/jamapsychiatry/article-abstract/2788767>) which would fit well with the authors narrative on the erosion of the welfare system in the UK from 2012 onwards
13. P75: please clarify the discussion on spatial analyses of mortality rates at post code level with a greater appreciation of the pitfalls in such.
14. P76: Please specify the mean and variance of the BetaBinomial as expressed in your parameters so that the precise choice of parameterisation is clear.
15. P77: there appear to be a few life expectancies in the high 90s or even 100s (City of London), which seems unfeasible. There must be wide uncertainty around these, right? Alternatively, the author mentions a mismatch of population estimates due to changes in adjacent areas.
16. P81: please clarify how the uncertainty is visualised (for instance in Fig 6.5). Is this the distribution at district level of the average posterior mean at LSOA level?
17. P82: One thing that is not discussed is the potential for stronger spatial effects in a city like London using LSOAs as compared to the whole of England using MSOAs. The other chapter finds similar results, but segregation patterns over larger areas may differ from smaller areas.
18. P88: please add a description on how cause-of-death is assigned in the UK to clarify the nature of the underlying data; similarly has any uncertainty been considered in the estimated of the contributions of death from each cause and would this be important to consider possibly in future work?
19. P88: It is not clear what the "ICD-10 code in the first position of the death record" is for the classification of deaths for neonates without underlying cause of death. Does this refer to contributory causes?
20. P89: It is not entirely clear on where ill-defined deaths go (R chapter mostly). The chapter does not mention anything, and Table D.1 mentions those deaths being included in the residual causes (but are not listed in those). There's also a mention to a redistribution without any details.
21. P91: the descriptions on Arriaga's method are very hard to follow without equations, and please lay out key assumptions also in the main body of the thesis.

22. P91: if the causes of death cover all causes (I don't see anything missing), why is the rescaling of cause specific mortality needed? (there's a mention in P103 to the "total mortality rule" which I assume refers to this, but it is unclear)
23. P97: what may be reasons of the far more common contribution of lung cancer to inequality among women compared to men? (this is somewhat talked about in chapter 8)
24. P109: please better justifying the need for this project. There is not much in the introduction about this.
25. P111: as with our earlier comment above on ill-defined causes, it isn't clear here how they are treated for this analysis.
26. P112: The author indicates there are very few studies looking at geographic patterns and cancer. Spain has been producing reports on that for some time (See here, the newest one is in English:  
<https://www.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAVE/EnfermedadesCronicas/Paginas/Atlas>  
direct link to the newest one here  
<https://www.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAVE/EnfermedadesCronicas/Documents/a>  
)
27. P117: reword the definition of posterior probability of a true decline

I have reworded the sentences to avoid the phrase "true decline" (PXX, LXX).

28. P125: For the discussion on epidemiologic transitions, I suggest  
<https://pubmed.ncbi.nlm.nih.gov/35639549/> and  
<https://academic.oup.com/ije/article/51/4/1054/6525745> for some nuanced discussions. Injuries have definitely been a weak spot of the theory.
29. P145: please provide a full justification of equation A.1, and under what assumptions death probabilities conditional upon survival to age  $x$  can be derived from cross-sectional, empiric mortality rates.

I have extended the derivation to show how equation A.1 can be derived intuitively, and have added that the results assume the mid-year population is used to approximate the number of person-years lived (PX).

30. P145: please provide a worked example to allow readers to build intuition into the approach, we suggest when  $n=1$ .

Equation A.1 is not intuitive, and there is no text in Preston (2001) (where all the life table detail is taken) or other sources that tries to build intuition - it is simply a result of algebraic manipulation. By providing further detail on the intuitive derivation of equation A.1 in #29, the reader is now better-placed to understand how the result has arisen.

31. P146: please explain in words how the equations ensure that the life table is closed in the sense that everyone must die at some point

I have extended the derivation to show how the expression for the number of survivors in the open-ended age interval has been derived intuitively using the fact that the number of deaths in the open-ended interval is equal to the number of people who survived to the final age group (PX).

32. P146: please provide reasonable detail and assumptions behind the Kannisto-Thatcher approach so the thesis is more self-contained.
33. P146 Section A.2: please provide a more detailed justification with greater attention to the underlying assumptions that lead to equation A.6. We did not follow details such as “subtract the probability of surviving”. Further, it seems the construction uses  $D^i_x$  from some year  $t$  to construct a hypothetical cohort that survives to cause  $i$  alone. However there are competing deaths and if these had not occurred,  $D^i_x$  would be larger for old ages  $x$ . It seems the underpinning are strong and these render interpretation of the calculated cause-specific life expectancies challenging.

## Other

1. Add Marta’s paper.
2. Cite latest paper for chapter 8.