

Calculus II Notes

Will Farmer

September 15, 2012

1 Integration By Parts

Formula

$$u \cdot v - \int v \cdot du$$

Example

$$\int x^3 \cdot \sin(x) \cdot dx$$

u	dv
x^3	$\sin(x)$
$3x^2$	$\cos(x)$
$6x$	$-\sin(x)$
6	$-\cos(x)$
0	$\sin(x)$

(1)

$$x^3 \cdot \cos(x) - \int 3x^2 \cdot \cos(x) \cdot dx$$

$$x^3 \cdot \cos(x) + 3x^2 \cdot \sin(x) - \int -6x \cdot \sin(x) \cdot dx$$

$$x^3 \cdot \cos(x) + 3x^2 \cdot \sin(x) + 6x \cdot \cos(x) - \int 6 \cdot \cos(x) \cdot dx$$

$$x^3 \cdot \cos(x) + 3x^2 \cdot \sin(x) + 6x \cdot \cos(x) - 6 \cdot \sin(x)$$

2 Trigonometric Integrals and Substitutions

Trigonometric Identities

$$1. \sec(x) = \frac{1}{\cos(x)}$$

$$2. \csc(x) = \frac{1}{\sin(x)}$$

$$3. \cot(x) = \frac{1}{\tan(x)}$$

$$4. \sin^2(x) + \cos^2(x) = 1$$

$$5. \tan^2(x) + 1 = \sec^2(x)$$

6. Double Angles

$$(a) \sin^2(x) = \frac{1}{2} \cdot (1 - \cos(2x))$$

$$(b) \cos^2(x) = \frac{1}{2} \cdot (1 + \cos(2x))$$

$$7. \frac{d}{dx} \tan(x) = \sec^2(x)$$

$$8. \frac{d}{dx} \sec(x) = \sec(x) \cdot \tan(x)$$

$$9. \int \sec(x) \cdot dx = \ln|\sec(x) + \tan(x)| + C$$

$$10. \int \tan(x) \cdot dx = -\log(\cos(x)) = \ln|\sec(x)| + C$$

11. Substitutions

Integrand	Substitution	Boundaries	Trig Identity
$\sqrt{a^2 - x^2}$	$x = a \cdot \sin(\Theta)$	$\frac{\Pi}{2} \leq \Theta \leq \frac{\Pi}{2}$	$\sin^2(\Theta) + \cos^2(\Theta) = 1$
$\sqrt{a^2 + x^2}$	$x = a \cdot \tan(\Theta)$	$\frac{-\Pi}{2} < \Theta < \frac{\Pi}{2}$	$\tan^2(x) + 1 = \sec^2(x)$
$\sqrt{x^2 - a^2}$	$x = a \cdot \sec(\Theta)$	$0 < \Theta < \frac{\Pi}{2}, \Pi < \Theta < \frac{3\Pi}{2}$	$\tan^2(x) + 1 = \sec^2(x)$

Examples

$$\begin{aligned}
 & \int \frac{x}{\sqrt{1-x^2}} \cdot dx \\
 & x = \sin(\Theta) \\
 & x^2 = \sin^2(\Theta) \\
 & dx = \cos(\Theta) \\
 & \int \frac{\sin(\Theta)}{\sqrt{1-\sin^2(\Theta)}} \cdot d\Theta \\
 & \int \frac{\sin(\Theta)}{\sqrt{\cos^2(\Theta)}} \cdot d\Theta \\
 & \int \frac{\sin(\Theta)}{\cos(\Theta)} \cdot d\Theta \\
 & \int \tan(\Theta) \cdot d\Theta \\
 & \ln|\sec(\Theta)| + C \\
 & \ln|\sec(\arcsin(x))| + C
 \end{aligned} \tag{2}$$

3 Partial Fraction Decomposition

This is meant to simplify integrals of rational functions.

Rational functions are ratios of polynomials in the form $\frac{P(x)}{Q(x)}$ while P(x) and Q(x) are arbitrary polynomials.

PROPER iff (degree of Q(x) > degree of P(x))

Long Division

Suppose $Q(x) \leq P(x)$

After long division you will get $\frac{P(x)}{Q(x)} = S(x) + \frac{R(x)}{Q(x)}$ while $R(x)$ is the remainder ($R(x)$ is ALWAYS less than $Q(x)$).

PFD: Replacing proper fractions by the sum of simpler fractions that we can integrate.

There are two ways to solve for A and B

1. Zero out one or the other (see ex.)
2. Expand and collect terms (see ex.)

Example

$$\begin{aligned}
 & \int \frac{x}{(x+1)(x-2)} \cdot dx \\
 & \frac{A}{x+1} + \frac{B}{x-2} \\
 & \frac{x}{(x+1)(x-2)} = \frac{A}{x+1} + \frac{B}{x-2} \\
 & \frac{(x+1)(x-2) \cdot x}{(x+1)(x-2)} = \frac{(x+1)(x-2) \cdot A}{x+1} + \frac{(x+1)(x-2) \cdot B}{x-2} \\
 & x = A \cdot (x-2) + B \cdot (x+1) \\
 & x = 2 \therefore 2 = 3B \therefore B = \frac{2}{3}, A = \frac{1}{3} \\
 & \text{or} \\
 & x = Ax - 2A + Bx + B \\
 & 1 = A + B \\
 & 0 = B - 2A \\
 & B = 2A \\
 & A = \frac{1}{3}, B = \frac{2}{3}
 \end{aligned} \tag{3}$$

Cases

1. For each 1st order, non-repeated factor, you add to the PFD a term of the form $\frac{A}{ax+b}$

$$\frac{A_0}{a_0x+b_0} + \frac{A_1}{a_1x+b_1} + \dots + \frac{A_n}{a_nx+b_n}$$

2. For each 1st order factor $(ax+b)$ repeated n times, $[(ax+b)^n]$ add it to the PFD n times.

$$\frac{A_0}{a_0x+b_0} + \frac{A_1}{(a_1x+b_1)^2} + \dots + \frac{A_n}{(a_nx+b_n)^n}$$

3. For each irreducible 2nd order, non-repeated factor $[(ax^2+bx+c)$ for $(b^2-4ac) < 0]$ add it to the PFD one term.

$$\frac{Ax+B}{ax^2+bx+c}$$

4. For each irreducible 2^{nd} order, repeated factor $[(ax^2 + bx + c)^n \text{ for } (b^2 - 4ac) < 0]$ add it to the PFD n terms.

$$\frac{A_0x + B_0}{(ax^2 + bx + c)^1} + \frac{A_1x + B_1}{(ax^2 + bx + c)^2} + \dots + \frac{A_nx + B_n}{(ax^2 + bx + c)^n}$$

4 Sequences

A sequence is an ordered, infinite list of numbers.

$\lim_{n \rightarrow \infty} a_1, a_2, a_3, \dots, a_n$
 $\{a_n\}_{n=1}^{\infty}$ indicates a sequence.

We can think of a sequence as a function:

$$n \in \mathbb{N} \text{ and } f(n) = a_n$$

Two types of sequence definition

1. Linearly: $a_n = \frac{n}{n+1}$ so $a_1 = \frac{1}{2}, a_2 = \frac{2}{3}, \text{etc.}$
2. Recursively (Fibonacci): $\{f_n\}_{n=1}^{\infty} f_1 = 1, f_2 = 2, f_n = f_{n-1} + f_{n-2}$

A sequence can also be pictured by graphing.

Squeeze Theorem (Sammich Theorem)

Let $\{a_n\}_{n=1}^{\infty}, \{b_n\}_{n=1}^{\infty}, \{c_n\}_{n=1}^{\infty}$ and $a_n \leq b_n \leq c_n$
 If $\lim_{n \rightarrow \infty} a_n = L$ and $\lim_{n \rightarrow \infty} c_n = L$ then $\lim_{n \rightarrow \infty} b_n = L$

5 Series

A series is a sum of an infinite sequence of terms.

Let $\{a_n\}_{n=1}^{\infty}$, the series with these terms is $\sum_{n=1}^{\infty} a_n$

It is possible for a sum of an infinite number of terms to add up to a finite number. This is called a convergent series.

Consider:

$$\begin{aligned} s_1 &= a_1 \\ s_2 &= a_1 + a_2 \\ s_n &= a_1 + a_2 + \dots + a_n \end{aligned} \tag{4}$$

s_n is called the sequence of partial sums ($\{s_n\}_{n=1}^{\infty}$) and the convergence of the series depends on its convergence.

If $\lim_{n \rightarrow \infty} s_n = L$ then it's convergent.

If $\lim_{n \rightarrow \infty} s_n = (+\infty, -\infty)$ then it's divergent.

If $\lim_{n \rightarrow \infty}$ does not exist, then the test is inconclusive.

Geometric Series

$\sum_{n=1}^{\infty} a \cdot r^{n-1}$ where $a \neq 0$ and r = the ratio of the series

If $-1 < r < 1$ then the series is convergent to $\frac{a}{1-r}$.

If $r \geq 1$ then it is divergent.

If $r \leq -1$ then it is not regular (neither convergent or divergent).

Shifting range of series

Formula:

$$\begin{aligned} & \sum_{n=x}^{\infty} a \cdot r^{n+y} \\ & \sum_{n=x}^{\infty} a \cdot r^{n-x+y+x} \\ & \sum_{n=x}^{\infty} a \cdot r^{n-x} \cdot r^{y+x} \\ & \sum_{n=x}^{\infty} r^{n-1} (a(r^{y+x})) \\ & \frac{a \cdot r^{y+x}}{1-r} \\ & = \frac{a \cdot r^{y+x}}{1-r} \end{aligned} \tag{5}$$

Harmonic Series

$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$ is **DIVERGENT**

$\sum_{n=1}^{\infty} \frac{1}{n^{\alpha}}$ is called the generalized harmonic series. It is convergent if $\alpha > 1$ and divergent if $\alpha \leq 1$.

6 Series Tests

Divergence Test (Test for un-convergence)

If $\sum_{n=1}^{\infty} a_n$ is convergent, then $\lim_{n \rightarrow \infty} a_n = 0$

If $\lim_{n \rightarrow \infty} a_n \neq 0$, then the series may or may not converge...

Integral Test

If $a_n = f(x)$ and the function is continuous, decreasing, and positive on $[1, +\infty)$, then the series is convergent iff the integral of the function is convergent. Iff $\int_1^{\infty} f(x) \cdot dx$ is convergent then $\sum_{n=1}^{\infty} a_n$ is convergent and vice-versa with divergence.

Comparison Test

Let $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ be two series with positive terms. If $a_n \leq b_n$ (for all n , or for all $n \geq N$) and $\sum_{n=1}^{\infty} b_n$ converges, then $\sum_{n=1}^{\infty} a_n$ converges as well. If $a_n \geq b_n$ (for all n , or for all $n \geq N$) and $\sum_{n=1}^{\infty} b_n$ diverges, then $\sum_{n=1}^{\infty} a_n$ diverges.

Limit Comparison Test

Let $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ be two series with positive terms. If $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = C, C \neq [0, \infty)$, then the two series are either both convergent or divergent.

Alternating Series Test (Leibniz' Test)

This ONLY applies to alternating series

$\sum_{n=1}^{\infty} (-1)^n a_n$ or $\sum_{n=1}^{\infty} (-1)^{n-1} a_n$ where $a_n \geq 0$.

if $\lim_{n \rightarrow \infty} a_n = 0$ and a_n is decreasing for all n then the series is convergent.

Absolute Values Test

For any series $\sum_{n=1}^{\infty} a_n$ you must consider the absolute value series $\sum_{n=1}^{\infty} |a_n|$. If the series of absolute values is convergent, it is called absolutely convergent. Any series that is absolutely convergent is also convergent ($-|a_n| \leq a_n \leq |a_n|$). There exist many series that are convergent, but *NOT* absolutely convergent (these are called conditionally convergent). For example, an alternating harmonic series is conditionally convergent.

Ratio Test

$$\sum_{n=1}^{\infty} a_n$$

if:

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L < 1 \text{ then the series is absolutely convergent}$$

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L > 1 \text{ then the series is not absolutely convergent}$$

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \text{ then the test is inconclusive}$$

Root Test

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \lim_{n \rightarrow \infty} (|a_n|)^{\frac{1}{n}} = L$$

If:

$L < 1$ then the series is absolutely convergent $L > 1$ then the series is not absolutely convergent $L = 1$ then the test is inconclusive

7 Power Series

$\sum_{n=1}^{\infty} \frac{1}{n^p}$ is convergent if $p > 1$ and divergent if $p \leq 1$

Representing Functions as Power Series

$f(x) = \frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$ where $|x| < 1$ (geometric series $a = 1$, ratio of x)

This is a power series centered at 0 with a radius of convergence of $R = 1$

If a power series $\sum_{n=0}^{\infty} c_n(x-a)^n$ has a radius of convergence $R > 0$ then the interval of convergence $|x-a| < R$

The function $f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n$ is differentiable inside the interval of convergence.

$$f'(x) = \sum_{n=0}^{\infty} c_n \cdot n \cdot (x-a)^{n-1} \text{ and } \int f(x) \cdot dx = \sum_{n=0}^{\infty} c_n \frac{(x-a)^{n+1}}{n+1} + C$$

Examples

$$\begin{aligned} f(x) &= \frac{1}{1-x^2} \\ &= \frac{1}{1-(-x^2)} \\ u &= (-x^2) \\ &= \frac{1}{1-u} \\ &= \sum_{n=0}^{\infty} u^n \\ &= \sum_{n=0}^{\infty} (-x^2)^n \\ &= \sum_{n=0}^{\infty} (-1)^n \cdot x^{2n} \end{aligned} \tag{6}$$

$$\begin{aligned}
f(x) &= \frac{1}{3+x} \\
&= \frac{1}{3 \cdot (1 + \frac{x}{3})} \\
&= \frac{1}{3} \cdot \frac{1}{1 + \frac{x}{3}} \\
u &= \frac{-x}{3} \\
\frac{1}{3} \sum_{n=0}^{\infty} \left(\frac{-x}{3} \right)^n \\
\frac{1}{3} \sum_{n=0}^{\infty} \left(\frac{-1}{3} \right)^n \cdot x^n \\
\sum_{n=0}^{\infty} \frac{1}{3} \cdot \frac{(-1)^n}{3^n} \cdot x^n \\
\sum_{n=0}^{\infty} \frac{(-1)^n}{3^{n+1}} \cdot x^n
\end{aligned} \tag{7}$$

Interval of convergence = $|x| < 3$

$$\begin{aligned}
f(x) &= \frac{1}{(1-x)^2} \rightarrow 1 - 2x - x^2 \\
&\quad \frac{d}{dx} \frac{1}{1-x} \\
\frac{d}{dx} \sum_{n=0}^{\infty} x^n, |x| < 1 \\
\sum_{n=0}^{\infty} n \cdot x^{n-1}
\end{aligned} \tag{8}$$

8 Taylor and MacLaurin Series

(Taylor series have arbitrary centers while MacLaurin are centered at 0)

Question: How do we know if a function has a power series representation? And for what values of x is it meaningful?

Assume: $\sum_{n=0}^{\infty} c_n(x-a)^n$ for $|x-a| < R$

In other words: $f(x) = c_0 + c_1 \cdot (x-a) + c_2 \cdot (x-a)^2$

Evaluate c_n at $x = a$. $c_n = \frac{f^{(n)}(a)}{n!}$ while $f^{(n)}(x)$ is the n^{th} derivative of $f(x)$

Theorem: If a function has a power series representation (or power series expansion) centered at a , i.e. $\sum_{n=0}^{\infty} c_n(x-a)^n$,

$|x-a| < R$, then the coefficients are given by $c_n = \frac{f^{(n)}(a)}{n!}$.

These are all Taylor series centered at a . If $a = 0$, then it is called a MacLaurin series.

Need:

The function to be infinitely differentiable inside the interval $|x-a| < R$

Take partial sums in the power series ($T_n(x)$)

$$T_n(x) = f(a) + f'(a)(x-a) + \dots + \frac{f^{(n)}(a)(x-a)^n}{n!}$$

$$\lim_{n \rightarrow \infty} T_n(x) = f(x)$$

Consider $f(x) - T_n(x) = R_n(x)$ where $R_n(x)$ is the remainder of order n of the Taylor series.

$$f(x) = \lim_{n \rightarrow \infty} T_n(x) \text{ is equivalent to saying } \lim_{n \rightarrow \infty} R_n(x) = 0$$

Theorem: If $f(x) = T_n(x) + R_n(x)$ where $T_n(x)$ is a Taylor polynomial of degree n of $f(x)$ at a , and if $\lim_{n \rightarrow \infty} R_n(x) =$

$$0 \text{ for all } |x - a| < R, \text{ then } f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)(x-a)^n}{n!}$$

Lagrange's Formula

The tricky bit is to show $\lim_{n \rightarrow \infty} R_n(x) = 0$

In this case it is useful to consider special representations of remainder functions

Formula: If a function has at least $n + 1$ derivatives in some interval I that contains the center, then there exists a number Z such that $x \leq Z \leq a$ and $R_n(x) = \frac{f^{(n+1)}(Z)(x-a)^{n+1}}{(n+1)!}$

If:

$x = 0$, then everything = 0

$x < 0$, then $x < Z < 0$

$x > 0$, then $0 < Z < x$

Application of Taylor Series

Given a function infinitely differentiable around $x = a$, to find its Taylor series centered at a :

1. Computer the Taylor coefficients $c_n = \frac{f^{(n)}(a)}{n!}$ and write down the corresponding Taylor series $\sum_{n=0}^{\infty} c_n (x-a)^n$
2. Find the radius of convergence and interval of convergence $|x - a| < R$
3. Apply Lagrange's formula for the remainder $R_n(x) = \frac{f^{(n+1)}(Z)(x-a)^{n+1}}{(n+1)!}$
4. $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$

Example

Find the MacLaurin series of $f(x) = e^x$ and its radius of convergence.

1

$$f^{(n)}(0) = e^0 = 1$$

$$c_n = \frac{1}{n!}$$

$$\sum_{n=0}^{\infty} \frac{x^n}{n!}$$

2

$$\text{Ratio Test of } \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$\lim_{n \rightarrow \infty} \left| \frac{x^{n+1} \cdot (n!)}{(n+1)! \cdot x^n} \right|$$

(9)

$$\lim_{n \rightarrow \infty} \left| \frac{x}{n+1} \right| = 0 \text{ regardless of } x$$

By the ratio test, the series is convergent for all $x \in \mathbb{R}$

The radius of convergence is $R = \infty$

3

$$0 < Z < x$$

$$R_n(x) = \frac{e^Z \cdot x^{n+1}}{(n+1)!}$$

if $x > 0$, then $0 < Z < x$ and by the Squeeze Theorem, it is 0

if $x < 0$, then $0 < Z < x$ and by the Squeeze Theorem, it is 0