



**GWL (GW + Wannier +Lanczos)**

# **Introduction to GW in QE: the GWL code**

**Margherita Marsili**


Dipartimento di Fisica e Astronomia “Galileo Galilei”,  
Università di Padova, Italy

CNR-NANO S3 center, Modena, Italy

# Summary

- General introduction
- Applications
- GWL
  - analytic continuation
  - no summation over empty states
  - optimal representation of large sets of vectors
  - use of Lanczos chain
  - optimal basis set for the polarizability

# What quantities are we looking for?




Photoemission:  
Experimentally we access **difference of total energies** between the neutral  $N$ -particle ground state and charged  $N+1$  or  $N-1$  particle excited states.

$$h\nu = E_f - E_i = E_{kin} + (E^*(N-1) - E^0(N))$$

$$h\nu = E_i - E_f = E_{kin} + (E^0(N) - E^*(N-1))$$


# Why?

The knowledge of electronic-energy level alignments, of defect levels, etc.. is extremely important for device design in photovoltaics, electronics, optoelectronics, catalysis etc...




In dye-sensitized solar cells the alignment of the electronic levels of the different part of the device is crucial for cells' functioning and performance.

# What kind of systems are we looking at? (1)




## Carbon Nanotubes

P. Umari, O. Petrenko, S. Taioli, and M.M. de Souza, J. Chem. Phys. 136, 181101 (2012).



## Zn-Phthalocyanine


P. Umari and S. Fabris J. Chem. Phys, 136, 174310 (2012)



## Organic dyes on TiO<sub>2</sub> surfaces

P. Umari, L. Giacomazzi, F. De Angelis, M. Pastore, S. Baroni, JCP 139, 014709 (2013)


# What kind of systems are we looking at? (2)



## **H<sub>2</sub>TPP thin films**

(4 porphyrin molecules in the unit cell)


M. Marsili, P. Umari, M. Pedio,  
in progress



## **Organic dyes + solvent on TiO<sub>2</sub> surfaces**

C. Verdi, E. Mosconi, F. De Angelis, M. Marsili, P. Umari  
Submitted to PRB (2014)


# What kind of systems are we looking at? (3)



## **H<sub>2</sub>TPP thin films**

(4 porphyrin molecules in the unit cell)

M. Marsili, P. Umari, M. Pedio,  
in progress



## **Organic dyes + solvent on TiO<sub>2</sub> surfaces**

C. Verdi, E. Mosconi, F. De Angelis, M. Marsili, P. Umari  
Submitted to PRB (2014)

**SYSTEMS WITH MORE THAN 900 ELECTRONS!!**

# GW standard approach

One of the main bottleneck of GW calculations arise from the computation of the electronic dielectric function.

$$\epsilon(r, r'; \omega) = \delta(r - r') - \int dr'' v(r - r'') P(r'', r', \omega)$$

$$P^0(r, r'; \omega) = \sum_{ij} (f_i - f_j) \frac{\Psi_i(r) \Psi_j^*(r) \Psi_j(r') \Psi_i^*(r')}{\omega - (\epsilon_j - \epsilon_i) + i\eta}$$

Computationally demanding for:

- sum over empty states
- dimension and frequency dependence of dielectric matrix



# GWL

GWL main features:

- Perturbative  $G_0W_0$  approach for large and inhomogeneous systems
- No summation on empty states
- Optimal basis set for the polarizability
- Polarizability and self-energy operators are expressed through a set of linear-response equations solved using the Lanczos-chain algorithm
- Full frequency dependence of the screening (no plasmon-pole)
- Mixed openmp and mpi parallelization

# GWL

- GWL is downloaded directly within the QE distribution
- `pw.x make.sys` should work so GWL can be compiled simply by typing:  

```
make gwl
```

in QE main directory.
- [www.gwl-code.org](http://www.gwl-code.org) is the code website where you can find tutorials and the user manual.

GWL Homepage - Mozilla Firefox

GWL Homepage

www.gwl-code.org

MAIN

Introduction

Get a copy

Tutorial

How to install

## GWL (GW + Wannier +Lanczos)

Università degli studi di Padova  
Dipartimento di Fisica e Astronomia  
via Marzolo 8, Padova, Italy

Theory@Elettra Group  
CNR-IOM DEMOCRITOS  
c/o Sincrotrone Trieste - SS14, Km 163,5 Basovizza, I-34012 TRIESTE

Project leader: **Paolo Umari** [paolo.umari<AT>unipd.it]

Authors would appreciate to be cited as followed:

"The results of this work have been obtained using the GWL package [1,2] implemented inside the Quantum-Espresso distribution [3]"

[1] P. Umari, G. Stenuit, S. Baroni, *Phys. Rev. B* **79** (2009) 201104(R)  
[2] P. Umari, G. Stenuit, S. Baroni, *Phys. Rev. B* **81** (2010) 115104  
[3] P. Giannozzi et al., *J. Phys.: Condens. Matt.* **21** (2009) 395502

# GW approximation, $W_c$

QP energies (that can be compared with direct and inverse PE) are computed in first order perturbation theory using  $\Sigma - V_{xc}$  as perturbation to the KS Hamiltonian.

$$\Sigma(1, 2) = iG(1, 2)W(1^+, 2) \quad 1 = (\mathbf{r}_1, t_1)$$

# GW approximation, $W_c$

QP energies (that can be compared with direct and inverse PE) are computed in first order perturbation theory using  $\Sigma - V_{xc}$  as perturbation to the KS Hamiltonian.

$$\Sigma(1, 2) = iG(1, 2)W(1^+, 2) \quad 1 = (\mathbf{r}_1, t_1)$$

$$W(1, 2) = v(1, 2) + \int d(3, 4)v(1, 3)\Pi(3, 4)v(4, 2)$$

# GW approximation, $W_c$

QP energies (that can be compared with direct and inverse PE) are computed in first order perturbation theory using  $\Sigma - V_{xc}$  as perturbation to the KS Hamiltonian.

$$\Sigma(1, 2) = iG(1, 2)W(1^+, 2) \quad 1 = (\mathbf{r}_1, t_1)$$

$$W(1, 2) = v(1, 2) + \int d(3, 4)v(1, 3)\Pi(3, 4)v(4, 2)$$

$$\Sigma(1, 2) = iG(1, 2)v(1^+, 2) + iG(1, 2)W_c(1^+, 2)$$

# GW approximation, $W_c$

QP energies (that can be compared with direct and inverse PE) are computed in first order perturbation theory using  $\Sigma - V_{xc}$  as perturbation to the KS Hamiltonian.

$$\Sigma(1, 2) = iG(1, 2)W(1^+, 2) \quad 1 = (\mathbf{r}_1, t_1)$$

$$W(1, 2) = v(1, 2) + \int d(3, 4)v(1, 3)\Pi(3, 4)v(4, 2)$$

$$\Sigma(1, 2) = iG(1, 2)v(1^+, 2) + iG(1, 2)W_c(1^+, 2)$$

$$\Sigma_c(\mathbf{r}_1, \mathbf{r}_2; \omega)$$

# Frequency dependence, analytic continuation

$$\langle \Psi_n | \Sigma_c(\epsilon_n) | \Psi_n \rangle$$

# Frequency dependence, analytic continuation

$$\langle \Psi_n | \Sigma_c(\epsilon_n) | \Psi_n \rangle$$



Analytic continuation.

$$\langle \Psi_n | \Sigma_c(i\omega) | \Psi_n \rangle$$



# Frequency dependence, analytic continuation

$$\langle \Psi_n | \Sigma_c(\epsilon_n) | \Psi_n \rangle$$

Expectation values computed  
on an imaginary frequency  
grid and then fitted to a  
multipole function.

(Some) input variables:

`n`  
`grid_freq`  
`n_multipoles`  
`n_grid_fit`  
`offset_fit`



Analytic continuation.

$$\langle \Psi_n | \Sigma_c(i\omega) | \Psi_n \rangle$$

# Frequency dependence, analytic continuation

$$\langle \Psi_n | \Sigma_c(\epsilon_n) | \Psi_n \rangle$$



Analytic continuation.

$$\langle \Psi_n | \Sigma_c(i\omega) | \Psi_n \rangle$$



FFT

$$\langle \Psi_n | \Sigma_c(i\tau) | \Psi_n \rangle$$

# From $W_c$ to the irreducible polarizability

$$\langle \Psi_n | \Sigma_c(i\tau) | \Psi_n \rangle = i \int dr dr' \Psi_n(r) G(r, r'; i\tau) W_c(r, r'; i\tau) \Psi_n(r')$$

To compute  $W_c$  we must compute the reducible polarizability  $\Pi(r, r'; i\tau)$


# From $W_c$ to the irreducible polarizability

$$\langle \Psi_n | \Sigma_c(i\tau) | \Psi_n \rangle = i \int dr dr' \Psi_n(r) G(r, r'; i\tau) W_c(r, r'; i\tau) \Psi_n(r')$$

To compute  $W_c$  we must compute the reducible polarizability  $\Pi(r, r'; i\tau)$

$$\Pi(i\omega) = P(i\omega) [1 - vP(i\omega)]^{-1}$$

Irreducible  
polarizability




# From $W_c$ to the irreducible polarizability

$$\langle \Psi_n | \Sigma_c(i\tau) | \Psi_n \rangle = i \int dr dr' \Psi_n(r) G(r, r'; i\tau) W_c(r, r'; i\tau) \Psi_n(r')$$

To compute  $W_c$  we must compute the reducible polarizability  $\Pi(r, r'; i\tau)$

$$\Pi(i\omega) = P(i\omega) [1 - vP(i\omega)]^{-1}$$

Irreducible  
polarizability



$$P(i\omega) = -4Re \sum_{vc} \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega}$$

$$\simeq \sum_{\alpha\beta} \phi_\alpha(r) P_{\alpha\beta}(i\omega) \phi_\beta(r')$$

# From $W_c$ to the irreducible polarizability

$$\langle \Psi_n | \Sigma_c(i\tau) | \Psi_n \rangle = i \int dr dr' \Psi_n(r) G(r, r'; i\tau) W_c(r, r'; i\tau) \Psi_n(r')$$

To compute  $W_c$  we must compute the reducible polarizability  $\Pi(r, r'; i\tau)$

$$\Pi(i\omega) = P(i\omega) [1 - vP(i\omega)]^{-1}$$

Irreducible  
polarizability

$$P(i\omega) = -4Re \sum_{vc} \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega}$$

$$\simeq \sum_{\alpha\beta} \phi_\alpha(r) P_{\alpha\beta}(i\omega) \phi_\beta(r')$$

Optimal basis set.  
How do I get this?

# From $W_c$ to the irreducible polarizability

$$\langle \Psi_n | \Sigma_c(i\tau) | \Psi_n \rangle = i \int dr dr' \Psi_n(r) G(r, r'; i\tau) W_c(r, r'; i\tau) \Psi_n(r')$$

To compute  $W_c$  we must compute the reducible polarizability  $\Pi(r, r'; i\tau)$

$$\Pi(i\omega) = P(i\omega) [1 - vP(i\omega)]^{-1}$$

Irreducible  
polarizability

$$P(i\omega) = -4Re \sum_{vc} \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega}$$

How do we efficiently  
compute the matrix  
elements?

$$\simeq \sum_{\alpha\beta} \phi_\alpha(r) P_{\alpha\beta}(i\omega) \phi_\beta(r')$$

Optimal basis set.  
How do I get this?

# From $W_c$ to the irreducible polarizability

$$\langle \Psi_n | \Sigma_c(i\tau) | \Psi_n \rangle = i \int dr dr' \Psi_n(r) G(r, r'; i\tau) W_c(r, r'; i\tau) \Psi_n(r')$$

To compute  $W_c$  we must compute the reducible polarizability  $\Pi(r, r'; i\tau)$

$$\Pi(i\omega) = P(i\omega) [1 - vP(i\omega)]^{-1}$$

Irreducible  
polarizability

$$P(i\omega) = -4Re \sum_{vc} \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega}$$

How do we efficiently  
compute the matrix  
elements?

$$\simeq \sum_{\alpha\beta} \phi_\alpha(r) P_{\alpha\beta}(i\omega) \phi_\beta(r')$$

Optimal basis set.  
How do I get this?



# GWL approach at work: irreducible polarizability

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_{vc} \int dr dr' \phi_{\alpha}(r) \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega} \phi_{\beta}(r')$$

1) Avoid summation on empty states  $\hat{P}_c = 1 - \sum_v |v\rangle\langle v|$

2)  $\hat{P}|\Psi_v\phi_{\nu}\rangle \simeq \sum_{\alpha=1}^{N_t} T_{\alpha,\mu\nu} |t_{\alpha}\rangle$  ;  $N_t \ll N_v * N_{\phi}$

3) Use lanczos-chain algorithm to compute:  $\langle t_{\alpha} | (H - \epsilon_v + i\omega)^{-1} | t_{\beta} \rangle$

**GWL approach at work:**  
**(1) no summation on empty states**

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_{vc} \int dr dr' \phi_{\alpha}(r) \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega} \phi_{\beta}(r')$$

**GWL approach at work:**  
**(1) no summation on empty states**

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_{vc} \int dr dr' \phi_{\alpha}(r) \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega} \phi_{\beta}(r')$$
$$= -4\text{Re} \sum_{cv} \int dr dr' \frac{\langle \Psi_v \phi_{\alpha} | r \rangle \langle r | \Psi_c \rangle \langle \Psi_c | r' \rangle \langle r' | \phi_{\beta} \Psi_v \rangle}{\epsilon_c - \epsilon_v + i\omega}$$

# **GWL approach at work:**

## **(1) no summation on empty states**

$$\begin{aligned} P_{\alpha\beta}(i\omega) &= -4\text{Re} \sum_{vc} \int dr dr' \phi_{\alpha}(r) \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega} \phi_{\beta}(r') \\ &= -4\text{Re} \sum_{cv} \int dr dr' \frac{\langle \Psi_v \phi_{\alpha} | r \rangle \langle r | \Psi_c \rangle \langle \Psi_c | r' \rangle \langle r' | \phi_{\beta} \Psi_v \rangle}{\epsilon_c - \epsilon_v + i\omega} \\ &= -4\text{Re} \sum_{cv} \frac{\langle \Psi_v \phi_{\alpha} | \Psi_c \rangle \langle \Psi_c | \phi_{\beta} \Psi_v \rangle}{\epsilon_c - \epsilon_v + i\omega} \end{aligned}$$

# GWL approach at work:

## (1) no summation on empty states

$$\begin{aligned}
 P_{\alpha\beta}(i\omega) &= -4\text{Re} \sum_{vc} \int dr dr' \phi_{\alpha}(r) \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega} \phi_{\beta}(r') \\
 &= -4\text{Re} \sum_{cv} \int dr dr' \frac{\langle \Psi_v \phi_{\alpha} | r \rangle \langle r | \Psi_c \rangle \langle \Psi_c | r' \rangle \langle r' | \phi_{\beta} \Psi_v \rangle}{\epsilon_c - \epsilon_v + i\omega} \\
 &= -4\text{Re} \sum_{cv} \frac{\langle \Psi_v \phi_{\alpha} | \Psi_c \rangle \langle \Psi_c | \phi_{\beta} \Psi_v \rangle}{\epsilon_c - \epsilon_v + i\omega} \\
 &= -4\text{Re} \sum_{cvc'} \langle \Psi_v \phi_{\alpha} | \Psi_c \rangle \langle \Psi_c | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | \Psi_{c'} \rangle \langle \Psi_{c'} | \phi_{\beta} \Psi_v \rangle
 \end{aligned}$$

# GWL approach at work:

## (1) no summation on empty states

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_{vc} \int dr dr' \phi_{\alpha}(r) \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega} \phi_{\beta}(r')$$

$$= -4\text{Re} \sum_{cv} \int dr dr' \frac{\langle \Psi_v \phi_{\alpha} | r \rangle \langle r | \Psi_c \rangle \langle \Psi_c | r' \rangle \langle r' | \phi_{\beta} \Psi_v \rangle}{\epsilon_c - \epsilon_v + i\omega}$$

$$= -4\text{Re} \sum_{cv} \frac{\langle \Psi_v \phi_{\alpha} | \Psi_c \rangle \langle \Psi_c | \phi_{\beta} \Psi_v \rangle}{\epsilon_c - \epsilon_v + i\omega}$$

$$= -4\text{Re} \sum_{cvc'} \langle \Psi_v \phi_{\alpha} | \Psi_c \rangle \langle \Psi_c | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | \Psi_{c'} \rangle \langle \Psi_{c'} | \phi_{\beta} \Psi_v \rangle$$

$$= -4\text{Re} \sum_v \langle \Psi_v \phi_{\alpha} | \hat{P}_c \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} \hat{P}_c | \phi_{\beta} \Psi_v \rangle$$

$$\hat{P}_c = 1 - \sum_v |v\rangle \langle v|$$

# **GWL approach at work:**

## **(2) optimal representations**

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_v \langle \Psi_v \phi_\alpha | \hat{P}_c \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} \hat{P}_c | \phi_\beta \Psi_v \rangle$$

Whatever way we choose to compute this expression we have to do it  $N_v \times N_\phi^2$  times!

# **GWL approach at work:**

## **(2) optimal representations**

$$P_{\alpha\beta}(i\omega) = -4Re \sum_v \langle \Psi_v \phi_\alpha | \hat{P}_c \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} \hat{P}_c | \phi_\beta \Psi_v \rangle$$

Whatever way we choose to compute this expression we have to do it  $N_v \times N_\phi^2$  times!

The effort is reduced if we find an optimal set of vectors, in terms of which we decompose the vectors belonging to the original set.

$$\hat{P}_c | \phi_\alpha \Psi_v \rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu,\alpha v} | t_\nu \rangle \quad N_t \ll N_v * N_\phi$$



# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu, \alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

How do we get the  $\{|t_\nu\rangle\}$ ?

# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu, \alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

How do we get the  $\{|t_\nu\rangle\}$ ?

**For every valence state v:**

1)  $\hat{P}_c |\phi_\alpha \Psi_v\rangle \longrightarrow \hat{P}_c |\phi_\alpha w_v\rangle = |\bar{\Psi}_{v\alpha}\rangle$  Rotate valence state into maximally localized Wannier function.

# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu, \alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

How do we get the  $\{|t_\nu\rangle\}$ ?

**For every valence state v:**

1)  $\hat{P}_c |\phi_\alpha \Psi_v\rangle \longrightarrow \hat{P}_c |\phi_\alpha w_v\rangle = |\bar{\Psi}_{v\alpha}\rangle$  Rotate valence state into maximally localized Wannier function.

2)  $O_{\alpha\beta}^v = \langle \bar{\Psi}_{v\alpha} | \bar{\Psi}_{v\beta} \rangle$  Compute the overlap matrix

# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu, \alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

How do we get the  $\{|t_\nu\rangle\}$ ?

**For every valence state v:**

1)  $\hat{P}_c |\phi_\alpha \Psi_v\rangle \longrightarrow \hat{P}_c |\phi_\alpha w_v\rangle = |\bar{\Psi}_{v\alpha}\rangle$  Rotate valence state into maximally localized Wannier function.

2)  $O_{\alpha\beta}^v = \langle \bar{\Psi}_{v\alpha} | \bar{\Psi}_{v\beta} \rangle$  Compute the overlap matrix

3)  $\sum_{\beta} O_{\alpha\beta}^v |o_{\beta}^{\mu}\rangle_v = o_{\mu} |o_{\alpha}^{\mu}\rangle_v$  Solve the eigenvalue problem and **keep the  $N_t$  eigenvectors with the largest eigenvalues.**

# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu, \alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

How do we get the  $\{|t_\nu\rangle\}$ ?

**For every valence state v:**

1)  $\hat{P}_c |\phi_\alpha \Psi_v\rangle \longrightarrow \hat{P}_c |\phi_\alpha w_v\rangle = |\bar{\Psi}_{v\alpha}\rangle$  Rotate valence state into maximally localized Wannier function.

2)  $O_{\alpha\beta}^v = \langle \bar{\Psi}_{v\alpha} | \bar{\Psi}_{v\beta} \rangle$  Compute the overlap matrix

3)  $\sum_{\beta} O_{\alpha\beta}^v |o_{\beta}^{\mu}\rangle_v = o_{\mu} |o_{\alpha}^{\mu}\rangle_v$  Solve the eigenvalue problem and keep the **Nt eigenvectors with the largest eigenvalues.**

This procedure is called singular value decomposition (SVD). Important input variable: `n_pola_lanczos` it defines the number of eigenvectors to be kept.

# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu, \alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

**We are still left with  $N_v$  blocks made of  $N_t$  “local” t states!**

# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu, \alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

**We are still left with  $N_v$  blocks made of  $N_t$  “local” t states!**

**4)** We perform an other SVD among the blocks keeping the eigenvectors with eigenvalue above a threshold value  $s$ .

Important input variable: `s_pola_lanczos`

# GWL approach at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu,\alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

We can now express **P0** in terms of the new set of vectors:

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_{\nu\mu} T_{\nu,\alpha v}^* T_{\mu,\beta v} \langle t_\nu | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | t_\mu \rangle$$

The dimension of the  $|t\rangle$  set is determined by:  
`n_pola_lanczos` and `s_pola_lanczos`.



# GWL tricks at work:

## (2) optimal representations

$$\hat{P}_c |\phi_\alpha \Psi_v\rangle \simeq \sum_{\nu=1}^{N_t} T_{\nu,\alpha v} |t_\nu\rangle \quad N_t \ll N_v * N_\phi$$

We can now express P0 in terms of the new set of vectors:

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_{\nu\mu} T_{\nu,\alpha v}^* T_{\mu,\beta v} \langle t_\nu | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | t_\mu \rangle$$

The dimension of the  $|t\rangle$  set is determined by:  
`n_pola_lanczos` and `s_pola_lanczos`.

# **GWL approach at work:**

## **(3) lanczos chain algorithm**

Efficient computation of  $\langle t_\nu | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | t_\mu \rangle$

# GWL approach at work:

## (3) lanczos chain algorithm

$$\text{Efficient computation of } \langle t_\nu | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | t_\mu \rangle$$

**1)** It is possible (see documentation) to recursively build a set of orthonormal vectors for which  $(\hat{H}_0 - \epsilon_v + i\omega)$  is tridiagonal starting from an arbitrary vector.

$$|t_\mu^0\rangle = |t_\mu\rangle, |t_\mu^1\rangle, \dots, |t_\mu^{N_l}\rangle$$

# GWL approach at work:

## (3) lanczos chain algorithm

$$\text{Efficient computation of } \langle t_\nu | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | t_\mu \rangle$$

**1)** It is possible (see documentation) to recursively build a set of orthonormal vectors for which  $(\hat{H}_0 - \epsilon_v + i\omega)$  is tridiagonal starting from an arbitrary vector.

$$|t_\mu^0\rangle = |t_\mu\rangle, |t_\mu^1\rangle, \dots, |t_\mu^{N_l}\rangle$$

**2)** Compute  $\sum_{i=1}^{N_l} \langle t_\nu | t_\mu^i \rangle \langle t_\mu^i | (\hat{H}_0 - \epsilon_v + i\omega)^{-1} | t_\mu^0 \rangle$

# GWL approach at work:

## (3) lanczos chain algorithm

$$\text{Efficient computation of } \langle t_\nu | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | t_\mu \rangle$$

**1)** It is possible (see documentation) to recursively build a set of orthonormal vectors for which  $(\hat{H}_0 - \epsilon_v + i\omega)$  is tridiagonal starting from an arbitrary vector.

$$|t_\mu^0\rangle = |t_\mu\rangle, |t_\mu^1\rangle, \dots, |t_\mu^{N_l}\rangle$$

**2)** Compute  $\sum_{i=1}^{N_l} \langle t_\nu | t_\mu^i \rangle \langle t_\mu^i | (\hat{H}_0 - \epsilon_v + i\omega)^{-1} | t_\mu^0 \rangle$

**3)** The  $\{|t_\mu^i\rangle\}$  sets are the same for all valence bands and frequencies because the two terms only involve the diagonal part of the operator.

# GWL approach at work:

## (2) lanczos chain algorithm

$$\text{Efficient computation of } \langle t_\nu | \frac{1}{\hat{H}_0 - \epsilon_v + i\omega} | t_\mu \rangle$$

1) It is possible (see documentation) to recursively build a set of orthonormal vectors for which  $(\hat{H}_0 - \epsilon_v + i\omega)$  is tridiagonal starting from an arbitrary vector.

$$|t_\mu^0\rangle = |t_\mu\rangle, |t_\mu^1\rangle, \dots, |t_\mu^{N_l}\rangle$$

2) Compute  $\sum_{i=1}^{N_l} \langle t_\nu | t_\mu^i \rangle \langle t_\mu^i | (\hat{H}_0 - \epsilon_v + i\omega)^{-1} | t_\mu^0 \rangle$

3) The  $\{|t_\mu^i\rangle\}$  sets are the same for all valence bands and frequencies because the two terms only involve the diagonal part of the operator.

This procedure is part of the so called Lanczos-chain algorithm.

Important input variable: `n_steps_lanczos_pola`, it determines the dimension of the vectors chain.

# Optimal representation of the dynamical polarizability (1)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

# Optimal representation of the dynamical polarizability (1)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

We choose the eigenvectors with largest eigenvalues of the time 0-polarizability

$$P_0(t = 0) = \sum_{vc} |\Psi_v \Psi_c\rangle \langle \Psi_c \Psi_v|$$



# Optimal representation of the dynamical polarizability (1)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

We choose the eigenvectors with largest eigenvalues of the time 0-polarizability

$$P_0(t=0) = \sum_{vc} |\Psi_v \Psi_c\rangle \langle \Psi_c \Psi_v|$$

In principle

**1)** SVD on  $Q_{\mu,\nu} = Q_{vc,v'c'} = \langle \Psi_v \Psi_c | \Psi_{v'} \Psi_{c'} \rangle$

namely solve  $\sum_{\nu} Q_{\mu,\nu} |q_{\nu}^{\alpha}\rangle = q_{\alpha} |q_{\mu}^{\alpha}\rangle$

# Optimal representation of the dynamical polarizability (1)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

We choose the eigenvectors with largest eigenvalues of the time 0-polarizability

$$P_0(t=0) = \sum_{vc} |\Psi_v \Psi_c\rangle \langle \Psi_c \Psi_v|$$

In principle

**1)** SVD on  $Q_{\mu,\nu} = Q_{vc,v'c'} = \langle \Psi_v \Psi_c | \Psi_{v'} \Psi_{c'} \rangle$

namely solve  $\sum_{\nu} Q_{\mu,\nu} |q_{\nu}^{\alpha}\rangle = q_{\alpha} |q_{\mu}^{\alpha}\rangle$

**2)**  $\phi_{\alpha} = \sum_{\mu} \frac{1}{\sqrt{q_{\alpha}}} q_{\mu}^{\alpha} |\Psi_{v_{\mu}} \Psi_{c_{\mu}}\rangle$

# Optimal representation of the dynamical polarizability (1)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

We choose the eigenvectors with largest eigenvalues of the time 0-polarizability

$$P_0(t=0) = \sum_{vc} |\Psi_v \Psi_c\rangle \langle \Psi_c \Psi_v|$$

In principle

**1)** SVD on  $Q_{\mu,\nu} = Q_{vc,v'c'} = \langle \Psi_v \Psi_c | \Psi_{v'} \Psi_{c'} \rangle$

namely solve  $\sum_{\nu} Q_{\mu,\nu} |q_{\nu}^{\alpha}\rangle = q_{\alpha} |q_{\mu}^{\alpha}\rangle$

**2)**  $\phi_{\alpha} = \sum_{\mu} \frac{1}{\sqrt{q_{\alpha}}} q_{\mu}^{\alpha} |\Psi_{v_{\mu}} \Psi_{c_{\mu}}\rangle$

**This requires a great number of conduction states to be computed!**

# Optimal representation of the dynamical polarizability (2)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

# Optimal representation of the dynamical polarizability (2)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

1) Consider “fake” conduction states, namely: planewaves below a certain threshold projected onto the conduction manifold.

$$|\tilde{G}\rangle = \hat{P}_c |G\rangle; \quad \frac{1}{2}|G|^2 \leq E^* \Rightarrow P_0(t=0) = \sum_{vG} |\Psi_v \tilde{G}\rangle \langle \tilde{G} \Psi_v|$$

# Optimal representation of the dynamical polarizability (2)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

**1)** Consider “fake” conduction states, namely: planewaves below a certain threshold projected onto the conduction manifold.

$$|\tilde{G}\rangle = \hat{P}_c |G\rangle; \quad \frac{1}{2}|G|^2 \leq E^* \Rightarrow P_0(t=0) = \sum_{vG} |\Psi_v \tilde{G}\rangle \langle \tilde{G} \Psi_v|$$

**2)** Use max. loc. Wannier functions  $|\Psi_v \tilde{G}\rangle \longrightarrow |w_v \tilde{G}\rangle$

# Optimal representation of the dynamical polarizability (2)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

**1)** Consider “fake” conduction states, namely: planewaves below a certain threshold projected onto the conduction manifold.

$$|\tilde{G}\rangle = \hat{P}_c |G\rangle; \quad \frac{1}{2}|G|^2 \leq E^* \Rightarrow P_0(t=0) = \sum_{vG} |\Psi_v \tilde{G}\rangle \langle \tilde{G} \Psi_v|$$

**2)** Use max. loc. Wannier functions  $|\Psi_v \tilde{G}\rangle \longrightarrow |w_v \tilde{G}\rangle$

**3)** Gram-Schmidt orthonormalize  $\{|w_v \tilde{G}\rangle\}$  and keep the vectors  $\{|f_{\mu}\rangle\}$  whose norm is above a certain threshold  $s$ .

$$P_{\mu\nu} = \sum_{vG} \langle f_{\mu} | \Psi_v \tilde{G} \rangle \langle \tilde{G} \Psi_v | f_{\nu} \rangle$$

# Optimal representation of the dynamical polarizability (2)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

**1)** Consider “fake” conduction states, namely: planewaves below a certain threshold projected onto the conduction manifold.

$$|\tilde{G}\rangle = \hat{P}_c |G\rangle; \quad \frac{1}{2}|G|^2 \leq E^* \Rightarrow P_0(t=0) = \sum_{vG} |\Psi_v \tilde{G}\rangle \langle \tilde{G} \Psi_v|$$

**2)** Use max. loc. Wannier functions  $|\Psi_v \tilde{G}\rangle \longrightarrow |w_v \tilde{G}\rangle$

**3)** Gram-Schmidt orthonormalize  $\{|w_v \tilde{G}\rangle\}$  and keep the vectors  $\{|f_{\mu}\rangle\}$  whose norm is above a certain threshold  $s$ .

$$P_{\mu\nu} = \sum_{vG} \langle f_{\mu} | \Psi_v \tilde{G} \rangle \langle \tilde{G} \Psi_v | f_{\nu} \rangle$$

**4)** Diagonalize  $P_{\mu\nu}$  and keep the  $N_p$  eigenvectors with the largest eigenvalues.



# Optimal representation of the dynamical polarizability (2)

$$P(i\omega) \simeq \sum_{\alpha\beta} \phi_{\alpha}(r) P_{\alpha\beta}(i\omega) \phi_{\beta}(r') \quad \{\phi_{\alpha}(r)\}$$

1) Consider “fake” conduction states, namely: planewaves below a certain threshold projected onto the conduction manifold.

$$|\tilde{G}\rangle = \hat{P}_c |G\rangle; \quad \frac{1}{2}|G|^2 \leq E^* \Rightarrow P_0(t=0) = \sum_{vG} |\Psi_v \tilde{G}\rangle \langle \tilde{G} \Psi_v|$$

2) Use max. loc. Wannier functions  $|\Psi_v \tilde{G}\rangle \longrightarrow |w_v \tilde{G}\rangle$

3) Gram-Schmidt orthonormalize  $\{|w_v \tilde{G}\rangle\}$  and keep the vectors  $\{|f_{\mu}\rangle\}$  whose norm is above a certain threshold s.

$$P_{\mu\nu} = \sum_{vG} \langle f_{\mu} | \Psi_v \tilde{G} \rangle \langle \tilde{G} \Psi_v | f_{\nu} \rangle$$

4) Diagonalize  $P_{\mu\nu}$  and keep the  $N_p$  eigenvectors with the largest eigenvalues.

Relevant input variables: `pmat_cutoff`, cutoff for the plane waves; `s_pmat` threshold for the f-states; `numw_prod` dimension of the final basis set.

# Summary of algorithms and relevant input variables

- ✓ Use of optimal representations when expensive operations on large sets of vectors are needed.


```
n_pola_lanczos  
s_pola_lanczos  
n_self_lanczos  
s_self_lanczos
```

- ✓ Use of lanczos chain algorithm for the computation of response functions.

```
n_steps_lanczos_pola, n_steps_lanczos_self
```

- ✓ Optimal basis set for the polarizability

```
pmat_cutoff, s_pmat, numw_prod
```



# Thank you, references, acknowledgments

USER MANUAL: [http://www.gwl-code.org/manual\\_gwl.pdf](http://www.gwl-code.org/manual_gwl.pdf)

## GW APPROXIMATION:

L. Hedin and S. Lundqvist. Solid State Phys., 23:1, 1970.

F. Aryasetiawan, O. Gunnarsson, Rep. Prog. Phys. 61 (3): 237 (1998).

## ANALYTIC CONTINUATION TECHNIQUE:

H. N. Rojas, R. W. Godby, and R. J. Needs. Phys. Rev. Lett., 74, 1827, (1995)

M. M. Rieger, L. Steinbek, I. D. White, H. N. Rojas, and R. W. Godby. Comput. Phys. Commun., 117, 211, (1999)

## GWL NUMERICAL APPROACH:

P. Umari, G. Stenuit, and S. Baroni. Phys. Rev. B, 79:201104(R), 2009.

P. Umari, G. Stenuit, and S. Baroni. Phys. Rev. B, 81:115104, 2010.


**Dott. Paolo Umari,  
Università di Padova  
GWL main developer**




# GWW approach at work:

## (1) no summation on empty states

$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_{vc} \int dr dr' \phi_{\alpha}(r) \frac{\Psi_v(r) \Psi_c(r) \Psi_v(r') \Psi_c(r')}{\epsilon_c - \epsilon_v + i\omega} \phi_{\beta}(r')$$



$$P_{\alpha\beta}(i\omega) = -4\text{Re} \sum_v \langle \Psi_v \phi_{\alpha} | \hat{P}_c (H - \epsilon_v + i\omega)^{-1} \hat{P}_c | \Psi_v(r) \phi_{\beta} \rangle$$

$$\hat{P}_c = 1 - \sum_v |v\rangle \langle v|$$


# GWL approach

GWL is used to compute GW corrections for systems with more than 900 electrons. What makes this possible?

- ✓ Use of optimal representations when expensive operations on large sets of vectors are needed.
- ✓ Use of lanczos chain algorithm for the computation of response functions.
- ✓ Use of maximally localized Wannier functions.
- ✓ Exact removal of summation over empty states. (In traditional approaches the number of empty states to be considered is  $\sim 4$  times the number of valence bands!)