

Quant. Comp. HW - 2

Steven MacCoun

Oct. 18, 2005

1 Is Valid Superposition?

Determine if the state $|\phi\rangle$ is a valid superposition

$$|\phi\rangle = \frac{1}{1+i}|0\rangle + \frac{1}{1-i}|1\rangle$$

The state is valid provided that it is length normalized to one.

$$\left|\frac{1}{1+i}\right|^2 + \left|\frac{1}{1-i}\right|^2$$

First multiply by the complex conjugate/complex conjugate to make it clear what the real and imaginary parts are for each coefficient:

$$\begin{aligned} &= \left|\frac{1}{1+i} \cdot \frac{1-i}{1-i}\right|^2 + \left|\frac{1}{1-i} \cdot \frac{1+i}{1+i}\right|^2 \\ &= \left|\frac{1}{2} - \frac{1}{2}i\right|^2 + \left|\frac{1}{2} + \frac{1}{2}i\right|^2 \\ &= \left(\frac{1}{2}\right)^2 + \left(\frac{-1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 \\ &= (1/4) + (1/4) + (1/4) + (1/4) = 1 \end{aligned}$$

Therefore the state is **valid**.

Note that another, maybe simpler way to do this, would be to exploit the property that $|a/b| = |a|/|b|$ So

$$\begin{aligned} &\left|\frac{1}{1+i}\right|^2 + \left|\frac{1}{1-i}\right|^2 \\ &= \frac{|1|^2}{|1+i|^2} + \frac{|1|^2}{|1-i|^2} \\ &= (1/2) + (1/2) = 1 \end{aligned}$$

2 Find valid superposition

Given $|\phi\rangle = (1/2)|00\rangle + \frac{x}{2\sqrt{2}}|01\rangle + \frac{1}{2\sqrt{2}}|10\rangle + \frac{1}{2}|11\rangle$, what values of x would make this a valid superposition?

Again the normalization condition is applied:

$$|\frac{1}{2}|^2 + |\frac{x}{2\sqrt{2}}|^2 + |\frac{1}{2\sqrt{2}}|^2 + |\frac{1}{2}|^2 = 1$$

$$(1/4) + \frac{|x|^2}{8} + (1/8) + (1/4) = 1$$

$$\frac{(5 + |x|^2)}{8} = 1$$

$$|x|^2 = 3$$

Since x can be imaginary, all we know is that the real part must be equal to $\sqrt{3}$ Therefore:

$$\boxed{x = \sqrt{3} + bi}$$

3

Let $|\psi\rangle = \frac{1}{\sqrt{3}}(|00\rangle + |01\rangle + |10\rangle)$ Find:

$$|\phi\rangle = (H \otimes H)|\psi\rangle$$

$$= \frac{1}{\sqrt{6}}[(|00\rangle + |01\rangle + |10\rangle + |11\rangle)$$

$$+ (|00\rangle - |01\rangle + |10\rangle - |11\rangle)$$

$$+ (|00\rangle + |01\rangle - |10\rangle + |11\rangle)]$$

$$= \frac{1}{\sqrt{6}}[3|00\rangle + |01\rangle + |10\rangle + |11\rangle]$$

$$\boxed{= \frac{1}{2}|0\rangle_2 + \frac{1}{6}|1\rangle_2 + \frac{1}{6}|2\rangle_2 - \frac{1}{6}|3\rangle_2}$$

4 Deutsch Problem

Suppose we take U_f from the Deutsch problem and compute

$$(H \otimes 2)U_f(H \otimes 2)|11 \rangle$$

Recall that for the Deutsch function:

	X=0	X=1
f_0	0	0
f_1	0	1
f_2	1	0
f_3	1	1

First let's apply the first hadamard gate and call this intermediate state $|a \rangle$:

$$\begin{aligned} |a \rangle &= (H \otimes H)|11 \rangle \\ &= \frac{1}{2}(|0 \rangle - |1 \rangle)(|0 \rangle - |1 \rangle) \\ &= \frac{1}{2}(|0 \rangle |0 \rangle - |0 \rangle |1 \rangle - |1 \rangle |0 \rangle + |1 \rangle |1 \rangle) \end{aligned}$$

In computing the Hadamard on this next, a general equation can be set up for the four 'f' functions. Since anything XOR'ed with 0 is itself, and anything XOR'ed with 1 is it's complement, we can write $U_f|a \rangle$ as:

$$U_f|a \rangle = \frac{1}{2}[|0 \rangle |f(0) \rangle - |0 \rangle |\bar{f}(0) \rangle - |1 \rangle |f(1) \rangle + |1 \rangle |\bar{f}(1) \rangle]$$

Now let's consider the two cases $f(0) = f(1)$ and $f(0) \neq f(1)$:

Case 1: $f(0) = f(1)$

This implies we are using either f_0 or f_3 .

f_0 :

$$HU_{f_0}|a \rangle = H\frac{1}{2}[|00 \rangle - |01 \rangle - |10 \rangle + |11 \rangle]$$

$$\begin{aligned}
&= \frac{1}{2} \left[\frac{1}{2} (|00\rangle + |01\rangle + |10\rangle + |11\rangle) \right. \\
&\quad - \frac{1}{2} (|00\rangle - |01\rangle + |10\rangle - |11\rangle) \\
&\quad - \left(\frac{1}{2} (|00\rangle + |01\rangle - |10\rangle + |11\rangle) \right. \\
&\quad \left. \left. - \frac{1}{2} (|00\rangle - |01\rangle - |10\rangle + |11\rangle) \right) \right] \\
&\quad \boxed{= \frac{1}{2} [-|00\rangle + |01\rangle + |10\rangle] \quad (f_0)}
\end{aligned}$$

f_3 :

$$\begin{aligned}
HU_{f_3}|a\rangle &= H \frac{1}{2} [|01\rangle - |00\rangle - |11\rangle - |10\rangle] \\
&= \frac{1}{4} [(|00\rangle - |01\rangle + |10\rangle + |11\rangle) \\
&\quad - (|00\rangle + |01\rangle + |10\rangle + |11\rangle) \\
&\quad - (|00\rangle - |01\rangle - |10\rangle + |11\rangle) \\
&\quad - (|00\rangle + |01\rangle - |10\rangle - |11\rangle)] \\
&= \frac{1}{4} [-2|00\rangle + 2|11\rangle] \\
&\quad \boxed{= \frac{1}{2} [-|00\rangle - |01\rangle + |10\rangle + |11\rangle] \quad (f_3)}
\end{aligned}$$

Now for the case of $f(0) \neq f(1)$, so we look at f_1 and f_2 . Note that applying U is easy here too, because we are always XOR a bit with it's complement, which always yields 1. Since the input gates will be the same for f_1 and f_2 , then each should return the same state:

f_1 :

$$\begin{aligned}
HU_{f_1}|a\rangle &= H \frac{1}{2} [|00\rangle - |01\rangle - |11\rangle - |10\rangle] \\
&= \frac{1}{4} [(|00\rangle + |01\rangle + |10\rangle + |11\rangle)
\end{aligned}$$

$$\begin{aligned}
& -(|00\rangle - |01\rangle + |10\rangle - |11\rangle) \\
& -(|00\rangle - |01\rangle - |10\rangle + |11\rangle) \\
& -(|00\rangle + |01\rangle - |10\rangle - |11\rangle) \\
& = \frac{1}{4}[-2|00\rangle + 2|11\rangle
\end{aligned}$$

$$= \frac{1}{2}[-|00\rangle + |01\rangle + |10\rangle + |11\rangle] \quad (f_1)$$

$f_2 :$

$$\begin{aligned}
HU_{f_3}|a\rangle &= H\frac{1}{2}[|01\rangle - |00\rangle - |11\rangle - |10\rangle] \\
&= \frac{1}{4}[(|00\rangle - |01\rangle + |10\rangle + |11\rangle) \\
&\quad -(|00\rangle + |01\rangle + |10\rangle - |11\rangle) \\
&\quad -(|00\rangle - |01\rangle - |10\rangle + |11\rangle) \\
&\quad -(|00\rangle + |01\rangle - |10\rangle - |11\rangle)]
\end{aligned}$$

$$= \frac{1}{2}[-|00\rangle - |01\rangle + |10\rangle + |11\rangle] \quad (f_1)$$

5 Bernstein-Verizani

Take $f(x)$ from the Bernstein-Verizani problem and compute:

$$U_f(H^{n \otimes 1})(|0\rangle_n |1\rangle_1)$$

$$= U_f\left(\frac{1}{2^{n/2}} \sum_{x=0}^{2^n-1} |x\rangle\right) \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$$

Since $U_f|x\rangle_n \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) = (-1)^{f(x)}|x\rangle_n \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$

$$= \frac{1}{2^{n/2}} \left(\sum_{x=0}^{2^n-1} (-1)^{f(x)} |x\rangle \right) \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$$

$$= \frac{1}{2^{(n+1)/2}} \left(\sum_{x=0}^{2^n-1} (-1)^{f(x)} |x\rangle \right) (|0\rangle - |1\rangle)$$

Recall that the function $f(x)$ in the Bernstein-Verizani is $a \cdot x$. Therefore, $f(x)$ is always either 0 or 1, and so each value in the sum is either $|x\rangle$ or $-|x\rangle$. The possible values of the input register are:

$$\frac{1}{2^{n/2}} \left(\sum_{x=0}^{2^n-1} (-1)^{f(x)} |x\rangle \right)$$

The only possible value of the output register is the superposed state:

$$\frac{1}{2^{(n+1)/2}} (|0\rangle - |1\rangle)$$

6 Derive State

Derive the state from problem 3 using any 1-bit gates and the 2-bit gate cNOT starting from $|00\rangle$. Just applying a hadamard to expand leaves a useless state, since all other gates applied maintain the state. So first apply a NOT gate:

$$|00\rangle \Rightarrow |11\rangle$$

Applying a hadamard to this yields:

$$HX|00\rangle = \frac{1}{2} (|00\rangle - |01\rangle - |10\rangle - |11\rangle)$$

Applying cNOT gates:

$$C_{21}HX|00\rangle = \frac{1}{2} (|00\rangle - |11\rangle - |10\rangle - |01\rangle)$$

And another cNOT:

$$C_{12}C_{21}HX|00\rangle = \frac{1}{2} (|00\rangle - |10\rangle - |11\rangle - |01\rangle)$$

Applying a Hadamard to this:

$$\begin{aligned} HC_{12}C_{21}HX|00\rangle &= \frac{1}{2} (|10\rangle + |11\rangle - (|00\rangle - |01\rangle)) \\ &= \frac{1}{2} (|10\rangle + |11\rangle - |00\rangle + |01\rangle) \end{aligned}$$

Presumably at this point I've accomplished something. After like a billion other permutations/attempts, I did get to having just -00_i , -01_i , -10_i , but could never achieve the correct coefficients.

Note that I also tried using a controlled Hadamard.

This is a controlled hadamard gate, that functions just like a cNOT gate except it applies the hadamard operator instead of the NOT operator to the non-control bit.

$$\begin{aligned}
C_{H_{12}}H|00\rangle &= \frac{1}{2}[|00\rangle + |01\rangle + |1\rangle \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) + |1\rangle \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)] \\
&= \frac{1}{2}[|00\rangle + |01\rangle + \frac{1}{\sqrt{2}}(|10\rangle + \frac{1}{\sqrt{2}}|11\rangle + \frac{1}{\sqrt{2}}|10\rangle - \frac{1}{\sqrt{2}}|11\rangle)] \\
&= \frac{1}{2}[|00\rangle + |01\rangle + \frac{2}{\sqrt{2}}|10\rangle]
\end{aligned}$$

Also close, but not quite there

7 Derive with only one cNOT

Although I did not finish the answer to the previous question, this question is essentially the same, because any cNOT gate can be rewritten as:

$$C_{ij} = \bar{n}_i + X_j n_i$$

Since there is no limit on the number of 1-bit and type of 1 bit gates we can use, it follows that the pretend answer I got for number 6 is equivalent to this question, but each C_{ij} is replaced with the above equation.

8 Numbered formulae

Useful Hadamards:

$$(H \otimes H)|00\rangle = \frac{1}{2}(|00\rangle + |01\rangle + |10\rangle + |11\rangle) \quad (1)$$

$$(H \otimes H)|01\rangle = \frac{1}{2}(|00\rangle - |01\rangle + |10\rangle - |11\rangle) \quad (2)$$

$$(H \otimes H)|10\rangle = \frac{1}{2}(|00\rangle + |01\rangle - |10\rangle - |11\rangle) \quad (3)$$

$$(H \otimes H)|11 \rangle = \frac{1}{2}(|00 \rangle - |01 \rangle - |10 \rangle + |11 \rangle) \quad (4)$$

Use the *equation* environment to get numbered formulae, e.g.,

$$y_{i+1} = x_i^{2n} - \sqrt{5}x_{i-1}^n + \sqrt{x_{i-2}^7} - 1 \quad (5)$$

$$\frac{\partial u}{\partial t} + \nabla^4 u + \nabla^2 u + \frac{1}{2}|\nabla u|^2 = c^2 \quad (6)$$

9 Acknowledgments