

MA 6.101

Probability and Statistics

**Tejas Bodas**

Assistant Professor, IIT Hyderabad

# Topics

Last class we saw

- ▶ Conditioning
- ▶ Law of iterated Expectations

This class ..

- ▶ Sums of random variables & Convolutions
- ▶ Bayes Rule revisited
- ▶ Moment Generating functions

# Sums of independent random variable

- ▶ Consider  $Z = X + Y$ . What is the pdf of  $Z$  when  $X$  and  $Y$ ?
- ▶ What is  $p_Z(z)$  or  $f_Z(z)$ ?
- ▶  $p_Z(z) = \sum_{(x,y):x+y=z} p_{X,Y}(x,y)$
- ▶  $f_Z(z) = \int_{(x,y):x+y=z} f_{X,Y}(x,y) dx dy$ .
- ▶ Integral of a surface over line.
- ▶ [https://en.wikipedia.org/wiki/Line\\_integral](https://en.wikipedia.org/wiki/Line_integral)
- ▶ Since  $X$  and  $Y$  are independent  $p_{X,Y}(x,y) = p_X(x)p_Y(y)$  and  $f_{X,Y}(x,y) = f_X(x)f_Y(y)$ . This gives us

Convolution formula


$$p_Z(z) = \sum_x p_X(x)p_Y(z-x)$$
$$f_Z(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(z-x)dx$$

HW: What if  $X$  and  $Y$  are not independent?

# Examples

- ▶ EX1: Suppose  $X$  and  $Y$  are independent and  $U[0, 1]$ . Find the pdf and CDF of  $Z = X + Y$ .
- ▶ [https://en.m.wikipedia.org/wiki/File:Convolution\\_of\\_box\\_signal\\_with\\_itself2.gif](https://en.m.wikipedia.org/wiki/File:Convolution_of_box_signal_with_itself2.gif)
- ▶ Ex2: Suppose  $X$  and  $Y$  are outcomes of independent roll of dice. Find the pmf of  $Z = X + Y$ .

# Inference problem



- ▶  $X$  is an unobservable random variable with a known distribution.
- ▶ We only observe measurements  $Y$  that takes values according to  $f_{Y|X}(y|x)$ .
- ▶ Objective is to draw inference about  $X$  having seen a realization of  $Y$  i.e., Obtain  $f_{X|Y}(x|y)$  using only  $f_X(x)$  and  $f_{Y|X}(y|x)$ , both of which are known.

## Bayes Rule revisited

$$P(B/A) = \frac{P(A/B)P(B)}{P(A)}$$

For continuous random variables  $X$  and  $Y$

$$f_{X|Y}(x|y) = \frac{f_{Y|X}(y|x)f_X(x)}{f_Y(y)} = \frac{f_{Y|X}(y|x)f_X(x)}{\int_{-\infty}^{\infty} f_{Y|X}(y|t)f_X(t)dt}$$

For discrete random variables  $X$  and  $Y$

$$p_{X|Y}(x|y) = \frac{p_{Y|X}(y|x)p_X(x)}{p_Y(y)} = \frac{p_{Y|X}(y|x)p_X(x)}{\sum_i p_{Y|X}(y|i)p_X(i)}$$

## Example 3.19(Bertsekas)

Lifetime of a Phillips bulb is assumed to be an exponential random variable  $Y$  with parameter  $\Lambda$ .  $\Lambda$  itself is a uniform random variable over  $[1, 1.5]$ . You test a bulb and see that it has a lifetime of  $y$  units. What can you say about randomness of  $\Lambda$  having observed  $Y = y$ ?

► What is  $f_{\Lambda}(\lambda)$ ?

► What is  $f_{Y|\Lambda}(y|\lambda)$ ?

► What is  $f_Y(y)$ ?

►  $f_{\Lambda|Y}(\lambda|y) = \frac{2\lambda e^{-\lambda y}}{\int_1^{1.5} 2te^{-ty} dt}$  for  $\lambda \in [1, 1.5]$ .

# Bayes Rule revisited

For discrete  $N$  and continuous random variable  $Y$

$$P(N = n | Y = y) = \frac{f_{Y|N}(y|n)p_N(n)}{f_Y(y)} = \frac{f_{Y|N}(y|n)p_N(n)}{\sum_i f_{Y|N}(y|i)p_N(i)}$$

Equivalently

$$f_{Y|N}(y|n) = \frac{P(N = n | Y = y)f_Y(y)}{p_N(n)} = \frac{P(N = n | Y = y)f_Y(y)}{\int_{-\infty}^{\infty} P(N = n | Y = t)f_Y(t)dt}$$



## Example 3.20 (Bertsekas)

- ▶ Suppose  $X = 1$  w.p.  $p$  and  $X = -1$  w.p.  $1 - p$ . While transmitting this signal, it is corrupted by a Gaussian noise  $N \sim \mathcal{N}(0, 1)$ . We observe  $Y = X + N$ . Suppose you observe  $Y = y$ , then show that

$$P(X = 1|Y = y) = \frac{pe^y}{pe^y + (1 - p)e^{-y}}$$

- ▶ Intuitively, this probability goes to zero as  $y$  decreases to  $-\infty$  and increases to 1 as  $y$  increases to  $\infty$ .
- ▶  $P(X = 1|Y = y) = \frac{f_{Y|X}(y|1)p_X(1)}{f_Y(y)}$
- ▶ Here  $f_Y(y) = f_{Y|X}(y|1)p_X(1) + f_{Y|X}(y|-1)p_X(-1)$ .
- ▶ Substitute values to obtain answer.