

Triple-Gamma-Regularization

A Flexible Non-Convex Regularization Penalty based on the
Triple-Gamma-Prior

Unterweger Lucas Paul

Vienna University of Economics and Business (WU), Department of Economics

21st of June 2024

Overview

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties
Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

① Motivating Problem

② Existing Concepts

Motivating Duality of Ridge and LASSO

③ Mathematical Derivation

④ Properties of the Triple-Gamma Penalty

Comparison to existing Penalties

Varying the Hyperparameters

⑤ Simulation Study

⑥ Shortcomings and Discussion

⑦ Appendix

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Motivating Problem

[Back to contents](#)

Overfitting I

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical Derivation

Properties of the Triple-Gamma Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and Discussion

References

Appendix

- **Sparse Data Settings:** Scenarios where data is sparse, meaning the number of data points is limited relative to the number of features.
- **High Dimensionality Problem:** Model is at risk of overfitting because the model can fit the noise than the underlying pattern.
- **Ill-posed Problems in Regression:** Solution to the problem becomes sensitive to small changes in the data, resulting in large variances in the estimated parameters \implies model's predictions may generalize poorly.
- **Bias-Variance Tradeoff:** High variance models (*overfitting*) capture noise and fluctuations in training data \implies poor generalization. High bias models (*underfitting*) fail to capture the underlying trend. Hard to find balance!

Overfitting II

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical Derivation

Properties of the Triple-Gamma Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and Discussion

References

Appendix




Figure 1: 1st and 8th order polynomial fit to data (Green & Blue Lines fitted with extra red data point; Orange & Red Lines fitted without).

Some Solutions to Ill-posed Problems

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical Derivation

Properties of the Triple-Gamma Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and Discussion

References

Appendix

- ① **Cross-Validation** Estimate model based on subsets of the data to make estimates more robust. However, often not feasible due to data availability issues.
- ② **Feature Selection** Selecting a subset of relevant features, but often reliant on strict assumptions.
- ③ **Data Augmentation and Acquisition** Gather more data, use stochastic approaches to estimate your models, but similar issue as with *Cross-Validation*.
- ④ **Ensemble Methods** Combining the predictions of multiple models (e.g., bagging, boosting).
- ⑤ **Regularization using Loss Penalty** Induce shrinkage on estimates and penalize too-complex models by altering the Loss-function

Regularization

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical Derivation

Properties of the Triple-Gamma Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and Discussion

References

Appendix

This thesis focuses on the 5th approach to *regularization*, which adds a penalty term to the risk minimization problem.

$$\min_{f \in \mathcal{H}} \left\{ \sum_{i=1}^N L(y_i, f(x_i)) + \lambda J(f) \right\}$$

where $L(\cdot)$ refers to a loss function defined as some function of the true values y_i and the predicted values $f(x_i)$ ¹ and $J(f)$ is a penalty based on the chosen functional from a space of functions \mathcal{H} .

¹In the setting of OLS, this would be the *sum of squared residuals (SSR)*

Existing Concepts

[Back to contents](#)

Proposed Concepts

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Name	Penalty	Reference
Ridge	$\ \beta\ ^2$	(Hoerl & Kennard, 1970)
LASSO	$\ \beta\ $	(Tibshirani, 1996)
Elastic Net	$\lambda_1 \ \beta\ ^2 + \lambda_2 \ \beta\ _1$	(Zou & Hastie, 2005)
Arctan	$\frac{2}{\pi} \arctan(\beta)$	(Y. Wang & Zhu, 2016)
Gaussian	$1 - e^{-\beta^2}$	(John et al., 2022)

Table 1: Established Regularization Penalties

- others are *SCAD*, *MCP*, *SILO*, *Dantzig Selector*, ...
- F. Wang et al. (2020) find no "go-to" method which suits a broad range of problems

Motivating Duality of Ridge and LASSO

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

- An interesting mathematical connection can be found when looking at this problem from a Bayesian point of view.
- In Bayesian regression, the *posterior distribution* (up to a proportionality constant) takes the form

$$p(\beta|X, Y) \propto f(Y|X, \beta) \times p(\beta)$$

with β being a coefficient vector, Y being the target vector and X being a matrix of features.

- Choosing specific prior distributions lead to posterior distributions which have moments that correspond to point estimates from regularization (*Gaussian prior*: Ridge; *double-exponential prior*: LASSO)

The Triple-Gamma-Prior

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

- Cadonna et al. (2020) developed a new prior distribution which has unifying properties and provides a general form for several shrinkage effects. Moreover, it is given by a closed-form solution:

$$p(\sqrt{\beta_j}|\phi^\xi, a^\xi, c^\xi) = \frac{\Gamma(c^\xi + \frac{1}{2})}{\sqrt{2\pi\phi^\xi} \cdot B(a^\xi, c^\xi)} \cdot U\left(c^\xi + \frac{1}{2}, \frac{3}{2} - a^\xi, \frac{\beta_j}{2\phi^\xi}\right)$$

Hypothesis/Aim of this thesis:

Can the closed-form marginal distribution of the Triple-Gamma-Prior be used to derive a new regularization penalty and do its advantages carry over into the frequentist framework?

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

**Mathematical
Derivation**

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Mathematical Derivation

[Back to contents](#)

Likelihood

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Let's assume we have n data points of a response variable \mathbf{y} and a set of features \mathbf{X} . Assuming a standard linear model with a parameter vector β and standard normal i.i.d. errors, the *likelihood* is

$$\begin{aligned}\mathcal{L}(\mathbf{y}|\beta, \sigma^2, \mathbf{X}) &= \prod_i^n p(y_i|\beta, \sigma^2, X_i) \\ &= \prod_i^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2}\left(\frac{x_i - \mu}{\sigma}\right)^2\right) \\ &= \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{1}{2\sigma^2}(\mathbf{y} - \mathbf{X}\beta)^\top(\mathbf{y} - \mathbf{X}\beta)\right)\end{aligned}$$

Prior

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Assuming the individual parameters are independent a priori and using the *Triple-Gamma-Prior* from Cadonna et al. (2020), the *prior* distribution of the parameter vector β is

$$\begin{aligned}
 p(\beta) &= \prod_j^p p(\beta_j) = \prod_j^p p(\beta_j | \phi^\xi, a^\xi, c^\xi) \\
 &\propto \prod_j^p U\left(c^\xi + \frac{1}{2}, \frac{3}{2} - a^\xi, \frac{\beta_j^2}{2\phi^\xi}\right) \\
 &= \prod_j^p \frac{1}{\Gamma(c^\xi + \frac{1}{2})} \int_0^\infty e^{-(\frac{\beta_j^2}{2\phi^\xi})t} t^{c^\xi + \frac{1}{2} - 1} (1+t)^{\frac{3}{2} - a^\xi - c^\xi + \frac{1}{2} - 1} dt
 \end{aligned}$$

Posterior

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Using Bayes' Theorem, the posterior distribution (up to a proportionality constant) $p(\beta|\mathbf{X}, \mathbf{y}, \sigma^2)$ is then given by

$$p(\beta|\mathbf{X}, \mathbf{y}, \sigma^2) \propto \mathcal{L}(\mathbf{y}|\beta, \sigma^2, \mathbf{X}) \times p(\beta)$$

Taking the log yields

$$\begin{aligned} \log(p(\beta|\mathbf{X}, \mathbf{y}, \sigma^2)) &\propto -\frac{1}{2\sigma^2} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \log \left(\prod_j^p U \left(c^\xi + \frac{1}{2}, \frac{3}{2} - a^\xi, \frac{\beta_j^2}{2\phi^\xi} \right) \right) \\ &= -\frac{1}{2\sigma^2} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \sum_j^p \log \left(U \left(c^\xi + \frac{1}{2}, \frac{3}{2} - a^\xi, \frac{\beta_j^2}{2\phi^\xi} \right) \right) \end{aligned}$$

Triple-Gamma-Regularization

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Finally, the Triple-Gamma-Regularization can be retrieved from looking at the *maximum-a-posteriori* estimate, which minimizes the **negative(!)** log-posterior:

$$\hat{\beta}_{MAP} = \arg \min_{\beta \in \mathbb{R}^p} \left(\frac{1}{2\sigma^2} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \sum_j^p -\log \left(U \left(c^\xi + \frac{1}{2}, \frac{3}{2} - a^\xi, \frac{\beta_j^2}{2\phi^\xi} \right) \right) \right)$$

Thus, the **Triple-Gamma-Penalty** is given by:

$$J_{TP}(\beta) = \sum_j^p -\log \left(U \left(c^\xi + \frac{1}{2}, \frac{3}{2} - a^\xi, \frac{\beta_j^2}{2\phi^\xi} \right) \right)$$

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Properties of the Triple-Gamma Penalty

[Back to contents](#)

Comparison to other Penalties

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Comparison of Triple-Gamma-Penalty to other Penalties




Figure 2: Comparison of one form of the Triple-Gamma-Penalty to existing penalties

Varying a^ξ

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters


Simulation Study

Shortcomings and
Discussion

References

Appendix

Penalty Structure for different Values of a^ξ

Given that $c^\xi = 0.1$, $\kappa_B = 2$ & $\lambda = 1$. Normalized to zero.

Varying c^ξ

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties



Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Figure 3: (Left) Varying $c^\xi \rightarrow \infty$, (Right) Varying $c^\xi \rightarrow 0$

Varying κ_B^2

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties


Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Penalty Structure for different Values of κ_B Given that $a^\xi = 0.6$, $c^\xi = 0.1$ & $\lambda = 1$. Normalized to zero.

²Note that per definition $\phi^\xi = (2c^\xi)/(\kappa_2^\xi a^\xi)$

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Simulation Study

[Back to contents](#)

Sparse Data Setting

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix




Figure 4: Comparison of Estimates from several Regularization Approaches
($n = 100, p = 10$)

Simulation Study

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix




Figure 5: Distribution of the Sum of the Absolute Deviations of the Estimates to the True Coefficients (200 runs)

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

**Shortcomings and
Discussion**

References

Appendix

Shortcomings and Discussion

[Back to contents](#)

Summary, Shortcomings & Potential Extensions

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

- Implementation in Python still relatively slow compared to similar approaches
- Cannot reproduce effects of converging non-convex penalites like *Arctan*, *Gaussian*, etc

BUT

- It can replicate results from e.g. *LASSO* regression or induce its own form of shrinkage
- *Triple-Gamma-Penalty* is a flexible regularization penalty that corresponds with the Bayesian *Triple-Gamma-Prior*

References I

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties
Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

- Cadonna, A., Frühwirth-Schnatter, S., & Knaus, P. (2020). Triple the gamma—a unifying shrinkage prior for variance and variable selection in sparse state space and tvp models. *Econometrics*, 8(2), 20.
- Hoerl, A. E., & Kennard, R. W. (1970). Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12(1), 55–67.
- John, M., Vettam, S., & Wu, Y. (2022). A novel nonconvex, smooth-at-origin penalty for statistical learning. *arXiv preprint arXiv:2204.03123*.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1), 267–288.
- Wang, F., Mukherjee, S., Richardson, S., & Hill, S. M. (2020). High-dimensional regression in practice: An empirical study of finite-sample prediction, variable selection and ranking. *Statistics and computing*, 30, 697–719.

References II

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties
Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Wang, Y., & Zhu, L. (2016). Variable selection and parameter estimation with the atan regularization method. *Journal of Probability and Statistics*, 2016.

Zou, H., & Hastie, T. (2005). Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 67(2), 301–320.

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSO

Mathematical
Derivation

Properties of the
Triple-Gamma
Penalty

Comparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Appendix

[Back to contents](#)

Unifying Property of the Triple-Gamma-Prior

Motivating Problem

Existing Concepts

Motivating Duality of Ridge
and LASSOMathematical
DerivationProperties of the
Triple-Gamma
PenaltyComparison to existing
Penalties

Varying the Hyperparameters

Simulation Study

Shortcomings and
Discussion

References

Appendix

Table 1. Priors on $\sqrt{\theta_j}$ which are equivalent to (top) or special cases of (bottom) the triple gamma prior.

Prior for $\sqrt{\theta_j}$		$a^{\tilde{\zeta}}$	$c^{\tilde{\zeta}}$	κ_B^2	$\phi^{\tilde{\zeta}}$
$\mathcal{N}\left(0, \psi_j^2\right), \psi_j^2 \sim \text{GG}\left(a^{\tilde{\zeta}}, c^{\tilde{\zeta}}, \phi^{\tilde{\zeta}}\right)$	normal-gamma-gamma	$a^{\tilde{\zeta}}$	$c^{\tilde{\zeta}}$	$\frac{2c^{\tilde{\zeta}}}{\phi^{\tilde{\zeta}}a^{\tilde{\zeta}}}$	$\phi^{\tilde{\zeta}}$
$\mathcal{N}\left(0, \frac{1}{\kappa_j} - 1\right), \kappa_j \sim \mathcal{TPB}\left(a^{\tilde{\zeta}}, c^{\tilde{\zeta}}, \phi^{\tilde{\zeta}}\right)$	generalized beta mixture	$a^{\tilde{\zeta}}$	$c^{\tilde{\zeta}}$	$\frac{2c^{\tilde{\zeta}}}{\phi^{\tilde{\zeta}}a^{\tilde{\zeta}}}$	$\phi^{\tilde{\zeta}}$
$\mathcal{N}\left(0, \psi_j^2\right), \psi_j^2 \sim \text{SBeta2}\left(a^{\tilde{\zeta}}, c^{\tilde{\zeta}}, \phi^{\tilde{\zeta}}\right)$	hierarchical scaled beta2	$a^{\tilde{\zeta}}$	$c^{\tilde{\zeta}}$	$\frac{2c^{\tilde{\zeta}}}{\phi^{\tilde{\zeta}}a^{\tilde{\zeta}}}$	$\phi^{\tilde{\zeta}}$
$\mathcal{DE}\left(0, \sqrt{2} \psi_j\right), \psi_j^2 \sim \mathcal{G}\left(c^{\tilde{\zeta}}, \frac{1}{\lambda^2}\right)$	normal-exponential-gamma	1	$c^{\tilde{\zeta}}$	$2\lambda^2 c^{\tilde{\zeta}}$	$\frac{1}{\lambda^2}$
$\mathcal{N}\left(0, \tau^2 \psi_j^2\right), \psi_j \sim t_1$	Horseshoe	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{2}{\tau^2}$	τ^2
$\mathcal{N}\left(0, \frac{1}{\kappa_j} - 1\right), \kappa_j \sim \mathcal{B}(1/2, 1)$	Strawderman-Berger	$\frac{1}{2}$	1	4	1
$\mathcal{N}\left(0, \tau^2 \tilde{\xi}_j\right), \tilde{\xi}_j \sim \mathcal{G}\left(a^{\tilde{\zeta}}, a^{\tilde{\zeta}}\right)$	double gamma	$a^{\tilde{\zeta}}$	∞	$\frac{2}{\tau^2}$	-
$\mathcal{N}\left(0, \tau^2 \tilde{\xi}_j\right), \tilde{\xi}_j \sim \mathcal{E}(1)$	Lasso	1	∞	$\frac{2}{\tau^2}$	-
$t_\nu\left(0, \tau^2\right)$	half- t	∞	$\frac{\nu}{2}$	$\frac{2}{\tau^2}$	-
$t_1\left(0, \tau^2\right)$	half-Cauchy	∞	$\frac{1}{2}$	$\frac{2}{\tau^2}$	-
$\mathcal{N}\left(0, B_0\right)$	normal	∞	∞	$\frac{2}{B_0}$	-

Figure 6: Table 1 from Cadonna et al. (2020)