





Ordinary Differential Equations

MATH260

Dr. Ghrist Gonzaga University



EDITED BY

CAMERON WILLIAMSON







Contents

1	Inti	roduction to Ordinary Differential Equations	2
	1.1	Basic Terminology	2
		1.1.1 Ordinary vs. Partial Differential Equations	2
		1.1.2 Independent Variables, Dependent Variables, and Parameters	3
		1.1.3 Order of a Differential Equation	3
		1.1.4 What is a solution	3
	1.2	Systems of Differential Equations	4
	1.3	Families of Solutions, Initial-Value Problems	5
2	First-Order Differential Equations		
	2.1	Separable First-Order Equations	6
	2.2	Solving Linear ODEs	8
	2.3		9
	2.4	Existence and Uniqueness of Initial Value Problems	9
	2.5	Exact Ordinary Differential Equations	10
		2.5.1 Substitution	12
		2.5.2 Bernoulli Ordinary Differential Equations	13
	2.6		15
	2.7	Phase Lines and Equilibrium Solutions	15
		2.7.1 Bifurcations	16
3	Sec	ond-order Differential Equations	18
	3.1	General Theory of Homogeneous Linear Equations	18
	3.2	Homogeneous Linear Equations with Constant Coefficients	21
	3.3	The Spring-Mass Equation	23
	3.4	Non homogeneous Linear Equations	25
		3.4.1 Method of Undetermined Coefficients	25
		3.4.2 Variation of Parameters	27
	3.5	The Forced Spring-Mass System	29
	3.6	Solving Cauchy Euler Ordinary Differential Equations	20

Introduction to Ordinary Differential Equations

Differential equations come from real-world problems and problems in applied mathematics. When mathematics is applied to real-world problems, it is often the case that finding a relation between a function and its rate of change is easier than finding a formula for the function itself; it is this relation between an unknown function and its derivatives that produces a differential equation.

To give a very simple example, a biologist studying the growth of a population with size at time t given by the function P(t), might make the very simple, but logical, assumption that a population grows at a rate directly proportional to its size. In mathematical notation, the equation for P(t) could then be written as:

$$\frac{dp}{dt} = rP(t)$$

Where the constant of proportionality, r would probably be determined experimentally by biologists working in the field. Equations used for modeling population growth can be much more complicated than this, sometimes involving scores of interacting populations with different properties.

§1.1 Basic Terminology

Definition 1.1.1

differential equation is any equation involving an unknown function and one or more if its derivatives.

The following are examples of differential equations:

- 1. P'(t) = rP(t)(1 P(t)/N) H harvested population growth
- 2. $\frac{d^2x}{d\tau^2} + 0.9\frac{dx}{d\tau} + 2x = 0$ spring mass equation
- 3. $I''(t) + 4I(t) = sin(\omega t)$ RCL circuit showing beats
- 4. $y''(t) + \mu(y^2(t) 1)y'(t) + y(t)$ ban der Pol equation
- 5. $\frac{\partial^2}{\partial x^2}u(x,y)+\frac{\partial^2}{\partial y^2}u(x,y)=0$ Laplace's equation

§1.1.1 Ordinary vs. Partial Differential Equations

Differential equations fall into two very broad categories, called ordinary differential equations and partial differential equations. If the unknown function in the equation is a function of only one variable, the equation is called an ordinary differential equation. If

the unknown function in the equation depends on more than one independent variable, the equation is called a partial differential equation, and in this case, the derivatives appearing in the equation will be partial derivatives.

§1.1.2 Independent Variables, Dependent Variables, and Parameters

Three different types of quantities can appear in a differential equation. The unknown function, for which the equation is to be solved, is called the dependent variable, and when considering ordinary differential equations, the dependent variable is a function of a single independent variable. In addition to the independent and dependent variables, a third type of variable, called a parameter, may appear in the equation. A parameter is a quantity that remains fixed in any specification of the problem, but can very from problem to problem.

§1.1.3 Order of a Differential Equation

Another important way in which differential equations are classified is in terms of their order.

Definition 1.1.2

he order of a differential equation is the order of the highest derivative of the unknown function that appears in the equation.

The differential equation 1 is a first-order equation and the others are all secondorder. Even though equation 5 is a partial differential equation, it is still said to be of second order since no derivatives of order higher than two appear in the equation.

§1.1.4 What is a solution

Given a differential equation, what is a solution? We must realize that we are looking for a function, and therefore it needs to be defined on some interval of its independent variable.

Definition 1.1.3

n analytic solution of a differential equation is a sufficiently differentiable function that, if substituted into the equation, together with the necessary derivatives, makes the equation an identity (a true statement for all values of the independent variable) over some interval of the independent variables.

Problem 1. Show that the function $p(t) = e^{-2t}$ is a solution to the differential equation:

$$x'' + 3x' + 3x = 0$$

Solution. To show that it is a solution, compute the first and second derivatives of p(t):

$$p'(t) = -2e^{-2t}$$
$$p''(t) = 4e^{-2t}$$

When the three functions p(t), p'(t), and p''(t) are substituted into the differential equation in place of x, x', and x'', it becomes:

$$(4e^{-2t}) + 3(-2e^{-2t}) + 2(e^{-2t}) \equiv 0$$
$$(4 - 6 + 2)(e^{-2t}) \equiv 0$$
$$(0)(e^{-2t}) \equiv 0$$

which is an identity (in the independent variable t for all real values of t). When showing that both sides of an equation are identical for all values of the variables, we will use the equivalence $sign \equiv$.

Problem 2. Show that the function $\phi(t) = (1 - t^2)^{1/2} \equiv \sqrt{1 - t^2}$ is a solution of the differential equation x' = -t/x.

Solution. First, notice that $\phi(t)$ is not even defined outside the interval $-1 \le t \le 1$. In the interval -1 < t < 1, $\phi(t)$ can be differentiated by the chain rule (for powers of functions):

$$\phi'(t) = \frac{1}{2}(1 - t^2)^{-\frac{1}{2}}(-2t) = -\frac{t}{(1 - t^2)^{\frac{1}{2}}}$$

The right-hand side of the equation x' = -t/x, with phi(x) substituted for x, is

$$-\frac{t}{\phi(t)} = -\frac{t}{(1-t^2)^{\frac{1}{2}}}$$

which is identically equal to $\phi(t)$ wherever ϕ and ϕ' are both defined. Therefore, $\phi(t)$ is a solution to the differential equation x' = -t/x on the interval (-1, 1).

§1.2 Systems of Differential Equations

Problem 3. Show that the functions $x(t) = e^{-t}$, $y(t) = -4e^{-t}$ form a solution of the system of differential equations

$$x'(t) = 3x + yy'(t) = -4x - 2y$$

Solution. The derivatives that we need are $x'(t) = -e^-t$ and $y'(t) = -(-4e^{-t}) = 4e^{-t}$. Then substitution into the second equation gives:

$$3x + y = (2e^{-t}) + (-4e^{-t}) = (3 - 4)e^{-t} = -e^{-t} \equiv x'(t),$$

$$-4x - 2y = -4(e^{-t}) - 2(-4e^{-t}) = (-4 + 8)e^{-t} = 4e^{-t} \equiv y'(t);$$

therefore, the given functions of x and y form a solution for the system.

§1.3 Families of Solutions, Initial-Value Problems

In this section the solutions of some very simple differential equations will be examined in order to give us an understanding of the terms n-parameter family of solutions and general solution of a differential equation. We will also be shown how to use certain types of information to pick one particular solution out of a set of solutions.

While we do not yet have any formal methods for solving differential equations, there are some very simple equations that can be solved by inspection. One of these is:

$$x' = x$$

This first-order differential equation asks you to find a function x(t) which is equal to its own derivative at every value of t.

Definition 1.3.1

first-order differential equation with one initial condition specified is called an initial-value problem, usually abbreviated as an IVP. The solution of an IVP will be called a particular solution of the differential equation.

Problem 4. Solve the IVP $x' = x, x(0) = \frac{1}{2}$.

Solution. Since we just found that the general solution of x' = x is $x(t) = Ce^t$, we only need to use the initial condition to determine the value of C. This will pick out one particular curve in the family. Substituting t = 0 and $x(0) = \frac{1}{2}$ into the general solution,

$$x(0) = Ce^0 = C = \frac{1}{2}.$$

With $C = \frac{1}{2}$, the solution of the IVP is $x(t) = \frac{1}{2}e^t$. This particular solution is the dotted curve shown in Figure 1.1, with the initial point $(0, \frac{1}{2})$ circled.

First-Order Differential Equations

In this chapter, methods will be given for solving first- order differential equations. First-order means that the first derivative of the unknown function is the highest derivative appearing in the equation. This implied that the most general first-order differential equation has the form F(t,x,t')=0 for some function F, in this chapter, we will assume that the equation can be solved explicitly for x'. This means that our first-order differential equations can always be put in the form:

$$x' = f(t, x)$$

where f denotes an arbitrary function of two variables. To see why such an assumption makes sense, supposed the differential equation is:

$$(x'(x))^2 + 4x'(t) + 3x(t) = t.$$

It would be messy, but not impossible to use the quadratic formula to extract two differential equations of the form x' = f(t, x) from this quadratic equation. However, one could also imagine equations where solving for x'(t) is not even possible, and in such a case, some of our methods might not be applicable.

The material in this chapter will cover several analytic methods for solving first-order differential equations, each requiring the function f to have a special form. Two different graphical methods are also described; one for the general equation depending only on x. Numerical methods for first-order equations are introduced and theoretical issues of existence and uniqueness of solutions are discussed.

§2.1 Separable First-Order Equations

The first analytic method we will consider applies to first-order equations that can be written in the form

$$\frac{dx}{dt} = g(t)h(t);$$

that is, when the function f(t, x) can be factored into a product of a function of t times a function of x. Such a differential equation is called separable.

Problem 5. Determine which of the following first-order differential equations are separable. Hint: try to factor the right- hand side if the equation does not initially appear to be separable.

$$x' = xt + 2x \rightarrow x' = x(t+2) \rightarrow g(t) = t+2, h(x) = x$$

$$x' = x + \cos(t)$$

$$x' = xt^2 + t^2 - tx \rightarrow x' = (t^2 - t)(x + 1) \rightarrow g(t) = (t^2 - t), h(x) = (x + 1)$$

$$x' = x^2 + x + 3 \rightarrow x' = (1)(x^2 + x + 3) \rightarrow g(t) = 1, h(x) = x^2 + x + 3$$

1. If h(x) = 1, the separable equation x' = g(t) is just an integrating problem and the solution is

$$x = \int g(t)dt;$$

that is, x is just the **indefinite integral** of the function g(t). Remember that this means that x can be *any* function G(t) such that G'(t) = g(t), and this introduces an arbitrary constant into the solution. As an example, the solution of x' = t + 1 is

$$x(t) = \int (t+1)dt = \frac{t^2}{2} + t + c$$

Even in this simple case the solution is an infinite one-parameter family of functions.

2. If g(t) = 1, the separable equation x' = h(x) is called an **autonomous** first-order differential equation. Unless h(x) is a constant, it is no longer possible to solve the equation by simple integration, and the method given below must be used. Autonomous first-order differential equations are important and will be investigated more thoroughly in section 2.7. In the above examples, only the last equation is autonomous. The other three contain functions of t (other than the unknown function x(t)) on the right-and side.

Problem 6. Solve the differential Equation $\frac{dx}{dt} = -tx^2$.

Solution. Split dx/dt into two pieces, dx and dt, and do a bit of algebra to write:

$$-\frac{dx}{x^2} = tdt$$

Integrate each side with respect to its own variable to obtain:

$$\int \left(-\frac{1}{x^2}\right) = \int t dt \to \frac{1}{x} = \frac{t^2}{x} + C.$$

where the arbitrary constants on each side have been collected on the right. Solve this equation for x to obtain the one parameter family of solutions

$$x = \frac{1}{(t^2/2) + C}.$$

We should check that the function x(t) does satisfy the differential equation for any value of the constant C. It appears that this method works, but splitting dx/dt into

two pieces is not a mathematically condoned operation; therefore, a justification of the method needs to be given.

If an equation is separable, and x'(t) is written as dx/dt, both sides of the equation dx/dt = g(t)h(x) can be divided by h(x), and the equation becomes

$$\frac{1}{h(x(t))} \left(\frac{dx}{dt} \right) dt = \int g(t) dt + C.$$

The method of simple substitution can be applied to the integral on the left. If we substitute u = x(t), then du = (dx/dt)dt, and the equation becomes

$$\int \frac{1}{h(u)} du = \int g(t) dt + C.$$

Now let H(u) be any function such that H'(u) = 1/h(u) and G(t) any function with G'9t) = g(t). Then the equation above implies that

$$H(u) + C_1 = G(t) + C_2 \rightarrow H(u) = G(t) + C,$$

Where C is the constant $C_2 - C_1$ Replacing u again by x(t):

$$H(x(t)) = G(t) + C$$

Check carefully that the expression H(x) = G(t) + C is exactly the same as the solution obtained above. It is an **implicit solution** of $-dx/x^2 = tdt$; that is, it defines a relationship between the unknown function x and its independent variable t. If it can be solved explicitly for x as a function of t, the result is called an **explicit solution of the differential equation. As expected, the integration produces an infinite on-parameter family of solutions

Theorem 2.1.1

To solve a separable first-order differential equation, x'(t) = q(t)h(x):

- Write the equation in the form dx/dt = g(t)h(x).
- Multiply both sides by dt, divide by h(x), and integrate, to put the equation in the form

$$\int \frac{1}{h(x)} dx = \int g(t) dt.$$

- Find any function H(x) such that H'(x) = 1/h(x) and any function G(t) such that
- Write the solution as H(x) = G(t) + C
- \bullet If possible, solve the equation from the previous step explicitly for x, as a function of t.

§2.2 Solving Linear ODEs

Definition 2.2.1

An ODE is linear for the dependent variable y if it is homogeneous when g(x) = 0 and otherwise it is non-homogeneous.

$$a_1(x)\frac{dy}{dx} + a_0(x)y = g(x).$$

The standard form of a linear ODE is

$$\frac{dy}{dx} + P(x)y = f(x).$$

If the homogeneous ODE is separable: $y = e^{-\int P(x)dx}$, and $y_c = cy_1(x)$ where $y_1 = e^{-\int P(x)dx}$. For a nonhomogeneous ODE, we need to use the process of "variation of parameters". First we need to find the function u so that $y_p = u(x)y_1(x) = u(x)e^{-\int P(x)dx}$, and then we follow the following steps

- 1. Put in standard form
- 2. Determine P(x) and integrating factor $e^{\int P(x)dx}$.
- 3. Multiply std form by integrating factor.
- 4. Write

$$\frac{d}{dx}\left[e^{\int P(x)dx}y\right] = e^{\int P(x)dx}f(x).$$

5. Integrate both sides.

$\S 2.3$

§2.4 Existence and Uniqueness of Initial Value Problems

Given an initial value problem, how can I know that a solution exists, and if so, the uniqueness of that solution.

Problem 7. $\frac{dx}{dt} = \sqrt{x}$, x(0) = 0 has two solutions. These solutions are $x(t) = \frac{t^2}{4}$ and 0. We can change the initial condition to be x(0) = 5. We can separate our values to get:

$$x(t) = \left(\frac{t}{2} + \sqrt{5}\right)^2$$

If we change the initial condition to x(5) = -5 we get no solutions. We can also change the initial condition to x(5) = 0 which gives us two solutions:

$$x(t) = \left(\frac{t-5}{2}\right)^2$$
$$x(t) = 0$$

§2.5 Exact Ordinary Differential Equations

This highlights an analytic technique to find the solutions to an ordinary differential equation. This technique depends on if we are able to write an ODE in the form $\frac{d}{dx}(F(x,y(x))) = 0$. How can we tell if this is possible?

We can determine this by using the multi-variable chain rule from calculus 3:

$$\frac{d}{dx}(F(x,y(x))) = \frac{\partial F}{\partial x} + \frac{\partial F}{\partial y}\frac{dy}{dx} = 0$$

Now in order to actually solve the ordinary differential equation, we will turn the equation above into something that we can work with:

$$M(x,y) + N(x,y)\frac{dy}{dx} = 0$$

Now we ask, does f(x,y) exist such that $\frac{\partial F}{\partial x} = M(x,y)$ and $\frac{\partial F}{\partial y} = N(x,y)$? We can test this by checking to see if we can do the following:

$$\frac{\partial^2 F}{\partial x \partial y} = \frac{\partial^2 F}{\partial y \partial x} \rightarrow \boxed{\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}}$$

If it passed the test, we use the equations to find F(x,y), and then we can use F(x,y(x)) = C to find the solution to the problem.

Problem 8. Calculus 3 example. Consider a field $\vec{v} = \langle 2y + 4x, 2x - 5y \rangle$. Is this a gradient field? If so, what is its potential function. In order to find this, there must be a function f(x,y) such that $\nabla f = \vec{v}$.

$$\partial f \partial y \rangle = <6xy, 2x - 5y >$$

$$\frac{\partial f}{\partial x} = 6xy, \qquad \qquad \frac{\partial f}{\partial y} = 2x - 5y$$

Therefore, we know that $f_{xy} = f_{yx}$ Now we just need to set the two equations equal to each other and solve:

$$\frac{\partial}{\partial y}(2y+4x) \stackrel{?}{=} \frac{\partial}{\partial x}(2x-5y)$$

And we can solve that equation algebraically and get 2 = 2, therefore this is the solution to the ordinary differential equation

Problem 9. Solve $(4x + 2y) + (2x - 5y) \frac{dy}{dx} = 0$. We should set M(x, y) = 4x + 2y and set N(x, y) = (2x - 5y).

1. Testing to see if $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$

$$\frac{\partial}{\partial y}(4x+2y) \stackrel{?}{=} \frac{\partial}{\partial x}(2x-5y)$$

2. Now we need to find F(x,y):

$$\frac{\partial F}{\partial x} = 4x + 2y$$

$$\frac{\partial F}{\partial y} = 2x - 5y$$

$$F(x,y) = \int (4x + 2y)dx = 2x^2 + 2xy + c_y, \quad F(x,y) = \int (2x - 5y)dy = 2xy - \frac{5}{2}y^2 + c_x$$

We can determine that $c_y = \frac{5}{2}y^2$ and $c_x = 2x^2$ by looking at the two equations next to each other. Therefore, we know that $F(x,y) = 2x^2 + 2xy - \frac{5}{2}y^2$.

Our resulting solution is:

$$2xy - \frac{5}{2}y^2 + 2x^2 = C$$

This was an implicitly defined solution, so $F(x,y) = 2xy - \frac{5}{2}y^2 + 2x$, and our ode is:

$$\frac{d}{dx}\left[2xy(y) - \frac{5}{2}(y(x))^2 + 2x^2\right] = \frac{d}{dx}[C]$$
$$(2y + 2xy') - \frac{5}{2}2(y) + 4x = 0$$

Example 2 Solve the differential equation,

$$2x^{2}y + e^{y} + (x^{3} + xe^{y} - 2y)\frac{dy}{dx} = 0.$$

We can see that $3x^2y + e^y = M(x,y)$ and $(x^3 + xe^y - 2y)\frac{dy}{dx} = 0$. We can test this by checking to see if:

$$\frac{\partial M}{\partial u} \stackrel{?}{=} \frac{\partial N}{\partial x}$$

and by taking the partial derivatives of each of the equations, we can see that $3x^2 + e^y = 3x^2 + e^y$. Now that we know that this is an exact ordinary differential equation, we need to solve for F(x,y). In order to find F(x,y), we must do

$$\frac{\partial F}{\partial x} = M \qquad \qquad \frac{\partial F}{\partial y} = N.$$

This means that

$$F(x,y) = \int (3x^2y + e^y)dx \qquad F(x,y) = \int (x^3 + xe^y - 2y)dy$$
$$= x^3y + xe^y + C_y \qquad = x^3y + xe^y - y^2 + C_x$$

So, $F(x,y) = x^3yxe^y - y^2$, and our solution is:

$$x^3y + xe^y - y^2 = C$$

Now we need to find a solution that satisfies the initial value problem, y(-2) = 1.

$$(-2)^{3}(1) + (-2)e^{1} - (1)^{2} = C$$
$$-8 - 2e - 1 = C$$
$$-9 - 2e = C$$

Which results in our particular solution being:

$$x^3y + xe^y - y^2 = -9 - 2e$$

§2.5.1 Substitution

Sometimes we can make a substitution that turns one ordinary differential equation into one that we can solve.

Problem 10. Let's take the equation $\frac{dy}{dx} = \frac{1-x-y}{x+y}$. In order to do this you need to trade out y(x) for u(x):

$$\frac{dy}{dx} = \frac{du}{dx} - 1$$

$$\frac{du}{dx} - 1 = \frac{1-u}{u} = \frac{1}{u} - 1$$

After we do those steps, we can see that this is a separable ordinary differential equation, and we can use the normal method to solve a separable equation:

$$\int udu = \int 1dx$$

$$\frac{u^2}{2} = x + C$$

$$u^2 = 2x + D \leftarrow D = 2C$$

$$(x+y)^2 = 2x + D \rightarrow x + y = \pm\sqrt{2x + D}$$

$$y = -x \pm\sqrt{2x + D}$$

Where the initial condition determines the last equation above. Now we need to solve for the initial condition of y(6) = 4, and we can do this by just plugging in the values, and determining the sign on the square root:

$$4 - 6 \pm \sqrt{12 + D}$$

$$10 = \pm \sqrt{12 + D}$$

$$D = 88 \to y(x) = -x + \sqrt{2x + 88}$$

§2.5.2 Bernoulli Ordinary Differential Equations

We use Bernoulli's solution to an ordinary differential equation if the equation can be written in the form $\frac{dx}{dt} + p(t)x = q(t)x^n$, where if n = 0, the equation is linear and where if n = 1, the equation is both linear and separable. We can do a substitution for Bernoulli ordinary differential equations where we let $v = x^{1-n}$, and we trade out $x(t) \to v(t)$ to turn the differential into a linear differential. Another formula we can use for plugging in to find the separable ordinary differential equation is $\frac{du}{dx} + (1-n)p(x)u = (1-n)q(x)$, but we do not have an example to show for that method.

Problem 11. Consider the Ordinary differential equation $t\frac{dy}{dt} + y = \frac{1}{y^2}$, and solve. Our first step for this problem is to divide both sides by t to rearrange it into the Bernoulli form:

$$\frac{dy}{dt} + \frac{1}{t}y = \frac{1}{t}y^{-2}$$

Because y is our dependent variable, we will be looking for our n value, and we will find it on the y on the right-hand side. This gives us n = -2. Now we need to let $v(t) = y^{1-(-2)} = y^3$ and this means that $y(t) = (v(t))^{1/3}$. Now wee need to take the differential of this equation in order to find a substitution for $\frac{dy}{dt}$:

$$\frac{dy}{dt} = \frac{1}{3y^2} \frac{dv}{dt}$$
$$\frac{dv}{dt} = 3y^{-2} \frac{dy}{dt}$$

Once we have found a substitution for $\frac{dy}{dt}$, we can simply plug in our equation and solve for v(t):

$$\frac{1}{3y^2}\frac{dv}{dt} + y = \frac{1}{y^2}$$
$$\frac{t}{3}\frac{dv}{dt} + y^3 = 1$$

We solve for v(t), and then we can use the integrating factors method to solve the ordinary differential equation:

$$\frac{dv}{dt} + \frac{3}{t}v = \frac{3}{t}$$

$$\mu(t) = e^{\int \frac{3}{t}dt} = e^{3ln(t)} = t^3$$

$$t^3 \frac{dv}{dt} + 3t^2v = 3t^2$$

$$\frac{d}{dt}(t^3v) = 3t^2$$

$$\int t^3v = \int 3t^2dt = t^3 + C$$

$$v(t) = 1 + \frac{C}{t^3} \to y(t) = (v(t))^{\frac{1}{3}}$$

Now we just need to let $v(t) = y^3$ and plug this back into the equation to get $y(t) = (1 + \frac{C}{t^3})^{\frac{1}{3}}$ as our final solution.

Homogenous ordinary differential equations can be written in the form $\frac{dx}{dt} = f\left(\frac{x}{t}\right)$. This means that if we just substitute v for $\frac{x}{t}$, we will turn the Homogenous equation into a separable ordinary differential equation.

Problem 12. Consider the homogenous ODE, $(5y - 2x)\frac{dy}{dx} = 4x + 2y$. We can divide the 5y - 2x, so we will end up with the equation

$$\frac{dy}{dx} = \frac{4x + 2y}{5y - 2x} = \frac{4 + \frac{2y}{x}}{5\frac{y}{x} - 2}$$

The reason we can make this change is that we are dividing everything on the left side by x to get it in the form $\frac{y}{x}$. Our next step is to trade out y(x) for v(x):

$$\frac{4+2v}{5v+2} = v + x\frac{dv}{dx}$$

Which we can further simplify into $\frac{dv}{dx} = \frac{1}{x} \left(\frac{4+2v}{5v-2} - v \right)$. Now, we are able to solve this as a separable ordinary differential equation:

$$\frac{dv}{dx} = \frac{1}{x} \left(\frac{4+4v-5v^2}{5v-2} \right)$$
$$\frac{5v-2}{4+4v-5v^2} dv = \frac{1}{x} dx$$

Now we need to let $u-4+4v-5v^2$, and let du=(4-10v)dv=-2(5v-2)dv. And then we integrate with respect to u to end up with

$$-\frac{1}{2} \int \frac{1}{u} du = \int \frac{1}{x} dx$$

$$-\frac{1}{2} ln|u| = ln|x| + C$$

$$ln|4 + 4v - 5v^2| = -2ln|x| + D$$

$$4 + 4v + 5v^2 = e^{ln|x|^{-2} + D}$$

$$= e^{ln|x|^{-2}} e^D$$

$$4 + 4\left(\frac{y}{x}\right) - 5\left(\frac{y^2}{x}\right) = e^{ln|x|^{-2}} A$$

$$4x^2 + 4xy - 5y^2 = A$$

$\S 2.6$


§2.7 Phase Lines and Equilibrium Solutions

This is a qualitative technique for autonomous ordinary differential equations. This is used for finding the long term behavior of solutions for various initial conditions. An equilibrium solution is a constant function satisfying the ODE:

$$y(t) = k \rightarrow \frac{dy}{dt} = 0$$

and we need to look for the roots of f(y).

Problem 13. Consider the ODE: $\frac{dy}{dx} = 4 - y^2$, we know that the equilibrium solutions (stationary points / fixed points / critical points) are $0 = 4 - y^2$ and $y = \pm 2$. If we were to plot this as a slope field, we would see an image like the following:



We need to classify each of the lines here as stable (a sink), unstable (a source), or semi-stable (a node). From this slope diagram we can make a phase diagram.

The arrows in this diagram indicate whether $\frac{dy}{dx}$ is above or below 0. We can see that the long term behavior of the solution:

$$y_0 = y(t_0) > 2, \rightarrow y(t) \rightarrow 2$$

 $-2 < 2_0, \rightarrow y(t) \rightarrow 2$
 $y_0 < -2, y(t) \rightarrow -\infty$

Let's take a look at another example problem:


Problem 14. Compare the pivots of the two ordinary differential equations, $\frac{dy}{dt} = y^2(1-y)^2$, and $\frac{dy}{dx} = y^2(1-y^2)$.

$$\frac{dy}{dt} = y^{2}(1-y)^{2} \qquad \frac{dy}{dt} = y^{2}(1-y^{2})$$

$$0 = y^{2}(1-y)^{2} \qquad 0 = y^{2}(1-y^{2})$$

$$y = 0, \pm 1$$

From finding the equilibrium points of the differential equations, we can now craft a pivot diagram:



From this we can see that in the long term, the equations are going to look like this:

$$y_0 < 0, y(t) \rightarrow 0$$

$$0 < y_0 < 1, y(t) \rightarrow 1$$

$$1 < y_0, y(t) \rightarrow \infty$$

§2.7.1 Bifurcations

For ordinary differential equations with a parameter, there are times that a small change in the value of the parameter results in a huge change in the behavior of the solutions. It causes a change in the number or stability of the equilibrium solutions.

Problem 15. Find the bifurcation value for

$$\frac{dp}{dt} = 0.5p \left(1 - \frac{p}{100} \right) - H$$


The first step for finding a bifurcation for an equation is finding the equilibrium solutions. In order to do that, we set equation (2.69) equal to 0 and then rearrange it, so it is in the form

$$p^2 - 100p + 250H = 0$$

In this case, we are going to need to use the quadratic formula to find the equilibrium points:

$$p = \frac{100 \pm \sqrt{10000 - 1000H}}{2}$$
$$= 50 \pm 5\sqrt{100 - 10H}$$

We can determine that our bifurcation value is H=10, because if we take what's under the square root, and set it equal to zero we can find those values. Now we can plot our bifurcations:



Second-order Differential Equations

§3.1 General Theory of Homogeneous Linear Equations

The standard form for a general second order ordinary differential equation is

$$x''(t) + p(t)x'(t) + q(t)x(t) = f(t).$$

For us, p(t) and q(t) are typically going to be constants. Also we know that if f(t) = 0, the equation is homogeneous.

Lemma 3.1.1

Given an equation in the form

$$x''(t) + p(t)x'(t) + q(t)x(t) = 0,$$

if $x_1(t)$ and $x_2(t)$ are solutions to equation (3.2), then

$$x(t) = C_1 x_1(t) + C_2 x_2(t)$$

is also a solution to equation (3.1) for all C_1, C_2 's. Equation (3.3) is the general solution to a homogeneous ODE

Let's take a look at an application of lemma (3.1.1).

Problem 16. Consider the equation

$$t^2x''' + 2tx'' - 6x' = 0.$$

This equation is a third order, linear and homogeneous, but not autonomous ordinary differential equation. Let's put this differential equation into standard form:

$$x'' + \frac{2}{t}x'' + \frac{6}{t^2}x' = 0$$

As long as $t_0 \neq 0$, we have a unique solution if given 3 initial conditions. The domain of this ordinary differential equation is t > 0, t < 0. Earlier, in section 3.1, we showed

that $1, t^3, \frac{1}{t^2}$ were all solutions. Because this ordinary differential equation is linear and homogeneous, we know that

$$x(t) = C_1 1 + C_2 t^3 + C_3 \frac{1}{t^2},$$

is also a solution for all C_1, C_2, C_3 . Just note, equation (3.6) is a linear combination of $1, t^3, \frac{1}{t^2}$. So, this is a <u>general solution</u> as long as the functions, $1, t^3, \frac{1}{t^2}$ are all "different" from each other.

What is "different"? Different means no one function in a set can be written as a linear combination of other functions in the set. Different means that all of the functions are linearly independent. More generally, if

$$C_1 f_1(t) + C_2 f_2(t) + \dots + C_n f_n(t) = 0$$

Theorem 3.1.1

We can instead test for linear independence by using the wronskian,

$$w(f_1, f_2)(t) = det \begin{vmatrix} f_1(t) & f_2(t) \\ f'_1(t) & f'_2(t) \end{vmatrix}$$

- If w(t) = 0 for all t's, then $\{f_1, f_2\}$ are linearly dependent.
- If w(t) = 0 for all t's, then $\{f_1, f_2\}$ are linearly independent.

Now, if we try to use the wronskian on problem 16, we do it like so:

$$w(t) = \begin{vmatrix} 1 & t^3 & \frac{1}{t^2} \\ 0 & 3t^2 & -\frac{2}{t^2} \\ 0 & 6t & \frac{6}{t^4} \end{vmatrix}$$
$$\rightarrow 1 \times \begin{vmatrix} 3t^2 & -\frac{2}{t^2} \\ 6t & \frac{6}{t^4} \end{vmatrix}$$
$$= \frac{18}{t^2} + \frac{12}{t^2} = \frac{30}{t^2} \neq 0$$

So we know that $\{1, t^3, \frac{1}{t^3}\}$ are linearly independent.

Let's consider another problem where we need to show that the solutions are linearly dependent.

Problem 17. Let's consider the equation y''(x) = -9y, which has the solutions $\{\cos(3x), \sin(3x)\}$ and let's show that the two solutions are linearly independent:

$$w(x) = \begin{bmatrix} \cos(3t) & \sin(3t) \\ -3\sin(3t) & 3\cos(3t) \end{bmatrix} = 3\cos^2(3x) + 3\sin^2(3x) = 3 \neq 0$$

Which means that both solutions are valid for this ordinary differential equation. We can turn our original equation into y''(x) + 9y = 0 which is a homogeneous ordinary

differential equation. Our p(x) = 0, and because we know that $w(x) = Ce^{-\int p(x)dx}$, we know that the solution to our equation is 0.

Theorem 3.1.2

hat to we do when solving linear non-homogeneous ordinary differential equations?

- 1. Solve the associated homogeneous ODE $\to X_h(t)$, which will have many unknown constants
- 2. Find one solution to nonhomogeneous ODE $\to X_p(t)$

The general solution to the original nonhomogeneous ODE is

$$x(t) = x_h(t) + x_p(t)$$

Let's try out this new theorem on an example problem.

Problem 18. Consider the differential equation y'' - 5y' - 6y = 4x. We claim that

$$y(x) = c_1 e^{-x} + c_2 e^{6x} - \frac{2}{3} + \frac{5}{9}$$

Are $\{e^{-x}, e^{6x}\}$ linearly independent? Verify both solutions to equation (3.14).

$$w(x) = \begin{bmatrix} e^{-x} & e^{6x} \\ -e^{-x} & 6e^{6x} \end{bmatrix}$$
$$= 6e^{5x} + e^{5x} = 7e^{5x}$$

Now we need to verify that this is a solution:

$$y'(x) = -C_1 e^{-x} + 6C_2 e^{6x} - \frac{2}{3}$$
$$y''(x) = C_1 e^{-x} + 36C_2 e^{6x}$$

Let's see if the left hand side is equal to 4x like it was above.

$$y'' - 5y' - 6y = \left[C_1 e^{-x} + 36C_2 e^{6x}\right] - 5\left[-C_1 e^{-x} + 6C_2 e^{6x} - \frac{2}{3}\right]$$
$$-6\left[C_1 e^{-x} + C_2 e^{6x} - \frac{2}{3} + \frac{5}{9}\right]$$

$$= C_1[e^{-x} + te^{-x} - 6e^{-x}] + C_2[36e^{6x} - 30e^{6x} - 6e^{6x}] + \left[\frac{10}{3} + 4x - \frac{10}{3}\right]$$
$$= 0 + 0 + 4x.$$

§3.2 Homogeneous Linear Equations with Constant Coefficients

Homogeneous ordinary differential equations are ones that come in the form

$$ay'' + by' + cy = 0$$

The first method of solving homogeneous linear equations with constant coefficients is the method of lucky guess. In this method, we let $y(x) = e^{rx}$. If we take the multiple derivatives of e^{rx} , we can find what y'(x) and y''(x) equal. Now we can plug in the e^{rx} into our equation to get

$$ar^{2}e^{rx} + bre^{rx} + ce^{rx} = 0$$
$$e^{rx}[ar^{2} + br + c] = 0.$$
$$ar^{2} + br + c = 0$$

There are 3 distinct cases for the roots of this equation.

- 1. 2 real distinct roots.
- 2. 1 repeated root.
- 3. 2 complex conjugate roots (A + Bi).

Let's take a look at an example of this method in action.

Problem 19. Consider the equation

$$y'' - 5y' - 6y = 0.$$

We are going to let $y(x) = e^{rx}$. We can skip adding e^{rx} to the beginning of each term because we can just divide that over in our minds. After going through that we get

$$r^{2} - 5r - 6 = 0$$
$$(r - 6)(r + 1) = 0$$
$$r = 6 - 1$$

From this, we know that our two solutions are $y_1(x) = e^{6x}$, $y_2(x) = e^{-x}$. Our general solution is

$$y(x) = C_1 e^{6x} + C_2 e^{-x}$$

Now all we need to do is check the wronskian W(x) to make sure that $\{e^{r_1x}, e^{r_2x}\}$ are linearly independent.

Let's explore a case of a real repeated root. Let's look at the equation

$$y'' - 8y' + 16 = 0.$$

We can let $y(t) = e^{rt}$, to get

$$r^2 - 8r + 16 = 0$$
.

Our roots for r are 4 and 4. One of our solutions should be $y_1(t) = e^{4t}$, but because we have a repeated root, I claim that $y_2(t) = te^{4t}$. We can verify that this is a repeated root by putting it in a wronskian:

$$w(t) = \begin{vmatrix} e^{4t} & te^{4t} \\ 4e^{4t} & e^{4t} + 4te^{4t} \end{vmatrix}$$
$$= e^{8t} + 4te^{8t} - 4te^{8t}$$
$$= e^{8t} \neq 0.$$

Because the wronskian result was not equal to zero, we know that these are two linearly independent solutions. Our general solution in this case would be

$$y(t) = C_1 e^{4t} + C_2 t e^{4t}.$$

If we were to solve a 4th order ordinary differential equation, our general solution would look something like this:

$$y(t) = C_1 e^{2t} + C_2 e^{-3t} + C_3 t e^{-3t} + C_4 t^2 e^{-3t}.$$

As you can see, the main difference between the second and third order ordinary differential equations is the amount of roots you need to take. Let's take a look at the third case, having 2 complex conjugate values for $r = \alpha \pm i\beta$.

Consider the equation

$$y'' - 2y' + 5y = 0.$$

Let $y(t) = e^{rt}$. Our previous equation will turn into $r^2 - 2r + 5 = 0$, and we will need to use the quadratic formula for this case.

$$r = \frac{2 \pm \sqrt{4 - 20}}{2}$$
$$= \frac{2 \pm \sqrt{-16}}{2}$$
$$= 1 \pm 2i.$$

Because of Euler's identity,

$$e^{i\theta} = \cos(\theta) + i\sin(\theta),$$

we can determine that the general solution of this ordinary differential equation is

$$y(t) = C_1 e^{1t} \cos(2t) + C_2 e^{1t} \sin(2t).$$

Problem 20. Solve the homogeneous linear ordinary differential equation,

$$x''' + 3x'' - 4x' - 12x = 0.$$

Let $x(t) = e^{rt}$, this means that

$$x' = re^{rt}$$

$$x'' = r^{2}e^{rt}$$

$$x''' = r^{3}e^{rt}.$$

Our expanded out equation is

$$e^{rt}[r^3 + 3r^2 - 4r - 12] = 0.$$

From this, we can see that our roots are r = 2, -2, -3, which means that our general solution is

$$x(t) = C_1 e^{2t} + C_2 e^{-2t} + C_3 e^{-3t}.$$

§3.3 The Spring-Mass Equation

The spring-mass equation is a variation of Hooke's law, which states

$$F_{spring} = -kx(k > 0).$$

This tries to bring mass back to equilibrium where x(t) is the distance of the mass from its equilibrium position at time t. The differential version of the equation is

$$mx'' + bx' + kx = F(t).$$

Where mx'' is the acceleration, bx' is the resistance friction, kx is the spring force and F(t) is any extra force acting on the system. Where is gravity within this equation? It drops out because the stretching of the string due to gravity puts the block attached to the string into equilibrium.

When you first attach the mass and the mass stops moving,

$$m(0) = -k(s) - b(0) - mg \to mg = ks \to k.$$

Problem 21. A mass weighing 10 pounds stretches a string $\frac{1}{4}ft$. Assuming no dampening, find the amplitude period of oscillations of the mass if it is released from a point $\frac{1}{10}ft$ below it's equilibrium position with an initial upward velocity of $\frac{1}{20}\frac{ft}{s}$. Because the mass weighs 10 pounds, mg = 10, which means that $m = \frac{10}{32} = \frac{5}{16}\frac{lbs^2}{ft}$. The first thing we need to do is find K. This can be done by setting mg = ks and solving for

$$10 = \frac{1}{4}k$$

$$k = \frac{10}{\frac{1}{4}} = 40$$

$$[k] = \frac{10lb}{\frac{1}{ft}} = \left[\frac{lb}{ft}\right].$$

Our ordinary differential equation is going to be

$$\frac{5}{16}x'' + 40x = 0.$$

We are given the initial values of $x'(0) = -\frac{1}{20}$ and $x(0) = -\frac{1}{10}$. We get the ordinary differential equation

$$\rightarrow x'' + 128x = 0$$

٠

We are going to let $x(t) = e^{rt}$. This gives us

$$r^{2} + 128 = 0$$

 $r^{2} = -128$
 $r = \pm \sqrt{-128} = \pm_{8}\sqrt{2i}$.

The general solution to our ordinary differential equation in this situation is

$$x(t) = C_1 \cos\left(8\sqrt{2}t\right) + C_2 \sin\left(8\sqrt{2}t\right).$$

This gives us that the angular frequency of our system, $\beta = 8\sqrt{2} \left[\frac{1}{s}\right]$, and the period of our system, $T = \frac{\pi}{4\sqrt{2}}s$.

NOTE: In general, the natural frequency of a spring mass system is $\omega_0 = 8\sqrt{2}$. Now we need to apply our initial conditions to find C_1, C_2 .

$$x'(t) = -8\sqrt{2}C_1 \sin\left(8\sqrt{2}t\right) + 8\sqrt{2}C_2 \cos\left(8\sqrt{2}t\right)$$
$$x(0) = C_1 = -\frac{1}{10}$$
$$x'(0) = 8\sqrt{2}C_2 = \frac{1}{10}.$$

This gives us that our equation of motion is

$$x(t) = -\frac{1}{10}\cos\left(8\sqrt{2}t\right) + \frac{1}{160\sqrt{2}}\sin\left(8\sqrt{t}\right)$$

$$= R\sin\left(\beta t + \phi\right)$$

$$C_1 = R\sin\phi$$

$$C_2 = R\cos\phi$$

$$C_1^2 + C_2^2 = R^2$$

$$\sqrt{\left(-\frac{1}{10}\right)^2 + \left(\frac{1}{160\sqrt{2}}\right)^2} = R$$

$$\sqrt{\frac{513}{51200}} = R$$

$$R \approx 0.100ft$$

$$\phi = tan^{-1}\left(\frac{C_1}{C_2}\right) + \frac{1}{60}\sqrt{2}\sin(8\sqrt{2}t)$$

$$= tan^{-1}\left(-16\sqrt{2}\right)$$

$$\phi = -1.4226rad$$

$$x(t) = -\frac{1}{10}\cos\left(8\sqrt{2}t\right) + \frac{1}{160\sqrt{2}}\sin\left(8\sqrt{2}t\right)$$

$$\approx 0.1001\sin\left(8\sqrt{2}t - 1.5266\right).$$

x(t) is the displacement of mass from equilibrium at time t. The following is the general formula for a mass-spring system.

$$mx'' + bx' + kx = 0$$
 where $mbk > 0$.

Let's consider a dampening situation. If we let $x(t) = e^{rt}$, then

$$mr^{2} + br + k = 0$$

$$\rightarrow r = \frac{-b \pm \sqrt{b^{2} - 4mk}}{2m}.$$

• Repeated r values. For this case, we are going to get $b^2-4mk=0$, which is a critically dampened system. This would make $r=-\frac{b}{2m},-\frac{b}{2m}$, which would make our general solution

$$x(t) = C_1 e^{-\frac{b}{2m}t} + C_2 t e^{-\frac{b}{2m}t}.$$

If we take the $\lim_{t\to\infty} x(t)$, we can see that this equation goes to zero.

• 2 real distinct roots.

$$r_1 = \frac{-b + \sqrt{b^2 - 4mk}}{2m} < b, r_2 = \frac{-b - \sqrt{b^2 - 4mk}}{2m} < 0.$$

If $b^2 - 4mk > 0$, we have an over dampened system. $r < r_1 < 0$. The solution to this equation would be

$$x(t) = C_1 e^{r_1 t} + C_2 e^{r_2 t}.$$

The limit as this ordinary differential equation goes to infinity is zero.

• 2 real complex conjugate roots. If

$$r = -\frac{b}{2m} \pm i \frac{\sqrt{4mk - b^2}}{2m}.$$

Our $\alpha = -\frac{b}{2m}$, and our $\beta = i\frac{\sqrt{4mk-b^2}}{2m}$. Our general solution in this scenario would be

$$x(t) = C_1 e^{-\frac{b}{2m}t} \cos\left(i\frac{\sqrt{4mk - b^2}}{2m}t\right) + C_2 e^{-\frac{b}{2m}t} \sin\left(i\frac{\sqrt{4mk - b^2}}{2m}t\right).$$

$$\lim_{t \to \infty} x(t) = 0.$$

§3.4 Non homogeneous Linear Equations

§3.4.1 Method of Undetermined Coefficients

When can we use the method of undetermined coefficients? This method can be used with

- Most constant coefficient ordinary differential equations
- Nonhomogeneous term must be of the form:
 - Polynomials.
 - $-e^{cx}$
 - $-\sin(bx),\cos(bx)$

Theorem 3.4.1

After finding the homogeneous function $y_h(x)$:

- 1. Make your first guess at the form of $y_p(x)$ with unknown constants based on f(x).
- 2. Plug $y_p(x)$ into ordinary differential equation. (Left hand side equals right handside for all x's).
- 3. Solve linear system.
- 4. Plug back into the guess $y_p(x)$.
- 5. General solution is $y(x) = y_h(x) + y_p(x)$.

Problem 22. Consider the following ordinary differential equation.

$$x'' - 3x' - 4x = -7t.$$

In order to solve for $x_h(t)$, we set the right hand side equal to zero and let $x(t) = re^{rt}$.

$$r^{2} - 3r - 4 = 0$$
$$(r - 4)(r + 1) = 0$$
$$r = 4.1$$

This gives us that our general homogeneous solution is

$$x_h(t) = C_1 e^{4t} + C_2 e^{-t}$$
.

- 1. Now we make our initial guess of $x_p(t) = At + B$. Then we need to make sure that it does not match one of the homogeneous solutions from above.
- 2. $x'_{p}(t) = A, x''_{p}(t) = 0$

Now we must plug our $x_p(t)$ and $x_p''(t)$ into our original ordinary differential equation. This gives us

$$0 - 3A - 4(At + B) = -7t.$$

Now We need to separate all of the terms that have a t attached to them, and all of the solutions that are constant as

$$t: -4A = -7$$

 $1: -3A - 4B = 0$

3. Now we must simply solve for both A and B. After solving, we get that

$$x_p(t) = \frac{7}{4}t - \frac{21}{16}.$$

This gives us the general solution as

$$C_1 e^{4t} + C_2 e^{-t} + \frac{7}{4}t - \frac{21}{16}.$$

Just a caution, we need to make sure that we apply our initial and boundary conditions to our general solution only.

Problem 23. Consider the ordinary differential equation

$$x'' - 3x' - 4x = 6e^{-4t}.$$

We can see that $x_h(t) = C_1 e^{4t} + C_2 e^{-t}$. Our initial guess is going to be $x_p(t) = Ae^{-4t}$. Using our initial guess we can do the following

$$x_p'' - 3x_p - 4x_p = 6e^{-4t}$$
$$16Ae^{-4t} + 12Ae^{-4t} - 4Ae^{-4t} = 6e^{-4t}$$
$$24A = 6 : A = \frac{1}{4}x_p(t) = \frac{1}{4}e^{-4t}.$$

Because our initial guess matched on of the solutions in the homogeneous equation we need to modify our equation or else we will be incorrect, but how do we know how to modify our guess? We need to add a t and check again to make sure that our solution is not matching. Let's try again with Ate^{-4t}

$$x'_p = 4e^{4t} + 4Ate^{4t}$$
$$x''_p = 8Ae^{4t} + 16Ae^{4t}.$$

After plugging x_p and x_p'' into our ordinary differential equation and solving the system at the end, we get the general result as

$$x(t) = C_1 e^{4t} + C_2 e^{-t} + \frac{6}{5} t e^{4t}.$$

§3.4.2 Variation of Parameters

This method is more tedious and difficult than the method described in section (3.4.1), but it is more universal and will find a particular solution to any non-homogeneous linear ordinary differential equation.

Theorem 3.4.2

o solve something using variation of parameters,

- 1. Find $y_h(x) = C_1 y_1(x) + C_2 y_2(x)$.
- 2. Let $y(x) = v_1(x)y_1(x) + v_2(x)y_2(x)$, where $y_1(x), y_2(x)$ are from the homogeneous solution. Our goal is to find v_1, v_2 .
- 3. Solve system for v'_1, v'_2 . In order to do this, we can use the following system

$$y'_1(x)v'_1(x) + y'_2(x)v'_2(x) = f(x)$$

$$y_1(x)v'_1(x) + y_2(x)v'_2(x) = 0.$$

In order to do this, we need to use Cramer's rule. Cramer's rule tells us to

find

$$w(x) = \begin{vmatrix} y_1(x) & y_2(x) \\ y'_1(x) & y'_2(x) \end{vmatrix}$$
$$v'_1(x) = \frac{\begin{vmatrix} 0 & y_2 \\ f(x) & y'_2 \end{vmatrix}}{w(x)}$$
$$v'_2(x) = \frac{\begin{vmatrix} y_1 & 0 \\ y'_1 & f(x) \end{vmatrix}}{w(x)}.$$

- 4. Integrate to find $v_1(x)$ and $v_2(x)$. $v_1(x) = \Box + c$, $v_2(x) = \Box + c$.
- 5. Simplify.

Problem 24. Solve the ordinary differential equation $x'' - 2x' + x = \frac{e^t}{t}$. First we need to find the homogeneous ordinary differential equation by letting $x_h(t) = e^{rt}$ and solving for r. When all is done we end up with $x_1(t) = e^t$, $x_2(t) = te^t$.

- 1. Let $x(t) = v_1(t)e^t + v_2(t)te^t$
- 2. Set up a system of equations.

$$e^{t}v'_{1} + te^{t}v'_{2} = 0$$
$$e^{t}v'_{1} + (e^{t} + te^{t})v'_{2} = \frac{e^{t}}{t}.$$

Now we can solve this to get $v_1' = -1, v_2' = \frac{1}{t}$.

3. Now we need to integrate both $v_1(t)$ and $v_2(t)$ like so

$$v_1(t) = \int -1dt = -t + C_1$$

 $v_2(t) = \int \frac{1}{t}dt = \ln|t| + C_2.$

4. So,

$$x(t) = (-t + C_1)e^t + (\ln|t| + C_2)te^t$$

= $-te^t + te^t \ln|t| + C_1e^t + C_2te^t$.

Another way we can solve this is using the wronskian,

$$w(t) = \begin{vmatrix} e^t & te^t \\ e^t & te^t + e^t \end{vmatrix} = \dots = e^{2t}$$

$$v'_1(t) = \frac{\begin{vmatrix} 0 & te^t \\ \frac{e^t}{t} & te^t + e^t \end{vmatrix}}{w(t)} = \dots = -1$$

$$v'_2(t) = \frac{\begin{vmatrix} e^t & 0 \\ e^t & \frac{e^t}{t} \end{vmatrix}}{w(t)} = \dots = \frac{1}{t}.$$

- §3.5 The Forced Spring-Mass System
- §3.6 Solving Cauchy Euler Ordinary Differential Equations