Hayat AI - Emergency Protocols

Quick Reference Guide for Crisis Situations

1. Severe Bleeding Control

- Apply direct pressure to the wound with a clean cloth
- If bleeding continues, apply a tourniquet 2-3 inches above wound
- Note the time of tourniquet application
- Do not remove the tourniquet once applied
- Seek medical help immediately

■■ Tourniquets only for life-threatening limb bleeding

2. Burn Treatment

- Remove person from source of burn
- Cool burn with running water for 10-20 minutes
- Remove jewelry or tight clothing near the burn
- Cover with sterile, non-stick bandage
- Do not pop blisters or apply ice directly

■■ Never apply butter, oil, or ointments to fresh burns

3. Fracture Stabilization

- Check for breathing and consciousness
- Stop any bleeding with direct pressure
- Immobilize injured area with a splint
- Apply ice packs wrapped in cloth to reduce swelling
- Keep person still and comfortable

■■ Do not try to realign broken bones

4. CPR Basics

- Check responsiveness and breathing
- Call for emergency help if available
- Begin chest compressions: push hard and fast (100-120/min)
- Push at least 2 inches deep in center of chest
- Allow full chest recoil between compressions
- Continue until help arrives or person shows signs of life

■■ Ratio: 30 compressions to 2 rescue breaths if trained

Generated by Hayat AI - Offline Medical Assistant
This is not a substitute for professional medical advice