# SI Result 1 (a): Legendre Polynomial




See **Fig. 2 (d)** and **Fig. 2 (e)**

$$P_1(\cos\theta) = \cos\theta$$

$$P_2(\cos\theta) = (3\cos^2\theta - 1)/2$$

$$\langle P_n(\cos\theta)\rangle(z) = \int_0^\pi d\theta \sin\theta\, P_n(\cos\theta) p(\theta, z)$$

# SI Result 1 (b): Area per lipid







- $Mg^{2+}$ cannot induce membrane condensation due to its **hydration shell** **[7]**.
- $Ca^{2+}$ reduces repulsion between headgroups, inducing **membrane condensation** **[7]**.
- As Conc. of $CaCl_2$ increases, portion of interfacial water (**IW**) of $CaCl_2$ decreases.

(See $\rho_{H_2O}(z)$ in **Fig. 2 (f)**)

**[7]** Yi Dong, J. Phys. Chem. Lett., 13, 11237-11244 (2022)

# SI Result 1 (c): Lateral Displacement Distribution

$Li_3PS_4$










$2Dt\sqrt{2}$

Li ions in "Glass" phase diffuse faster than those in the crystalline!!

$$G_s(\boldsymbol{r},t) = (4\pi Dt)^{-3/2} e^{-r^2/4Dt}$$




$$r^2 G_s(\boldsymbol{r},t)$$

$$r_{\max} = 2Dt\sqrt{2}$$



# SI Result 1 (c): Lateral Displacement Distribution

$$G_s(\boldsymbol{r}_\parallel, t) \cong \sum_n f_n G_{\mathcal{N}}\left(\boldsymbol{r}_\parallel, t | D_\parallel^{(n)}\right) \qquad \sum_n f_n = 1$$

$$G_{\mathcal{N}}(\boldsymbol{r}_\parallel, t | D_\parallel) \equiv (4\pi D_\parallel t)^{-1} e^{-r_\parallel^2/4D_\parallel t}$$

$$r_\parallel \times G_{\mathcal{N}}(\boldsymbol{r}_\parallel, t | D_\parallel) \rightarrow \text{ maximum at } r_\parallel = \sigma = \sqrt{2D_\parallel t}$$

```
In[2]:= P[x_, σ_] := (2 π σ^2)^-1 Exp[-x^2 / (2 σ^2)]
                                  |지수 함수
```









# SI Result 1 (c): Lateral Displacement Distribution

$$G_s(\boldsymbol{r}_\parallel, t) \cong \sum_n f_n G_{\mathcal{N}}\left(\boldsymbol{r}_\parallel, t | D_\parallel^{(n)}\right) \qquad \sum_n f_n = 1$$

$$G_{\mathcal{N}}(\boldsymbol{r}_\parallel, t | D_\parallel) \equiv (4\pi D_\parallel t)^{-1} e^{-r_\parallel^2/4D_\parallel t}$$

$$f(\boldsymbol{r}_\parallel, t) \equiv \frac{r_\parallel G_s(\boldsymbol{r}_\parallel, t)}{\sigma\, p_{\mathcal{N}}(\sigma, t|\sigma)}\Bigg|_{\sigma=\sqrt{2\langle D_\parallel\rangle t}}$$









# SI Result 1 (c): Lateral Displacement Distribution



# SI Result 1 (d): Region dependent lateral diffusion coefficient




$D_{\parallel}(n)$: determined by umbrella sampling (1 μs-long)

$\mathrm{Layer}(n)$: defined by the nodes of $\langle P_2(\cos\theta)\rangle(z)$ in the salt-free case. (see **Fig. 2 (e)**)

# SI Result 1 (e): Experimental result

**Fig. 3**




[Experiment]
DMPC
37 $H_2O$/lipid molecule.
0.45 M conc.



Multilamellar lipid vesicle







## Quasi-elastic neutron scattering study of the effects of metal cations on the hydration water between phospholipid bilayers


Cite as: Appl. Phys. Lett. **116**, 133701 (2020); doi: 10.1063/1.5144012
Submitted: 31 December 2019 · Accepted: 26 February 2020 ·
Published Online: 30 March 2020


View Online Export Citation CrossMark


H. Seto[1,a)] and T. Yamada[2,b)]

AFFILIATIONS

[1]Institute of Materials Structure Science/J-PARC Center, High Energy Accelerator Research Organization, Tokai 319-1106, Japan
[2]Neutron Science and Technology Center, Comprehensive Research Organization for Science and Society, Tokai 319-1106, Japan

[a)]Author to whom correspondence should be addressed: hideki.seto@kek.jp
[b)]Electronic mail: t_yamada@cross.or.jp





**J-PARC MLF BL02 DNA: Dynamic Spectrometer**

# SI Result 1 (f). Second-order reorientational TCF $C_2(t) \equiv \langle P_2(\cos\varphi(t))\rangle$

L Bai, *J. Phys. Chem. C.*, 123, 21528-21537 (2019)




$$\tau_{\text{reor}} \equiv \int_0^\infty dt\, C_2(t)$$

**Salt-free**

**12.53 ps**

| | 0.1M | 0.4M | 0.8M |
|---|---|---|---|
| $MgCl_2$ | 15.03 ps | 30.92 ps | 31.60 ps |
| $CaCl_2$ | 14.56 ps | 23.03 ps | 35.91 ps |

# SI Result 2. Discrete Green's function $G^{(d)}(n,t|m)$




SI. 2

$x$ axis: $n$
$y$ axis: $m$
$t = 512$ ps

- The gap between membranes was divided into **7 discrete regions**.
- For sufficiently large $t$, $G^{(d)}(n,t|m)$ converges to $P_{st}(n)$.
- Higher Conc. leads to a longer timescale for this process. (i.e., $G^{(d)}(n,t|m) \neq P_{st}(n)$) (See **SI Video**)