


Trends in Invasive Melanoma Thickness in Norway, 1983–2019



Raju Rimal

Oslo Centre for Biostatistics and Epidemiology (OCBE)
Department of Biostatistics, UiO

Presented at: [Cancer Registry of Norway](#)

30 October 2024

ORIGINAL REPORT



Trends in Invasive Melanoma Thickness in Norway, 1983–2019

Raju RIMAL¹, Trude E. ROBSAHM², Adele C. GREEN^{3,4}, Reza GHIAVAND^{2,5}, Corina S. RUEEGG⁵, Assia BASSAROVA⁶, Petter GJERSVIK⁷, Elisabete WEIDERPASS⁸, Odd O. AALEN¹, Bjørn MØLLER⁹, Flavie PERRIER¹ and Marit B. VEIERØD¹

¹*Oslo Centre for Biostatistics and Epidemiology, Department of Biostatistics, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway*, ²*Department of Research, Cancer Registry of Norway, Norwegian Institute of Public Health, Oslo, Norway*, ³*Department of Population*

Health, QIMR Berghofer Medical Research Institute, Brisbane, Australia, ⁴*Cancer Research UK Manchester Institute, University of Manchester, Manchester, UK*, ⁵*Oslo Centre for Biostatistics and Epidemiology, Oslo University Hospital, Oslo, Norway*, ⁶*Department of Pathology, Oslo University Hospital – Ullevål, Oslo, Norway*, ⁷*Department of Dermatology, Institute of Clinical Medicine, University of Oslo, Oslo, Norway*,

⁸*International Agency for Research on Cancer, Lyon, France*, and ⁹*Department of Registration, Cancer Registry of Norway, Oslo, Norway*


Citation: Acta Derm Venereol 2024; 104: adv26110. DOI <https://doi.org/10.2340/actadv.v104.26110>.

Copyright: © 2024 The Author(s). Published by MJS Publishing, on behalf of the Society for Publication of Acta Dermato-Venereologica. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>).


Submitted: Nov 20, 2024; **Accepted after revision:** Nov 29, 2024; **Published:** Sep 2, 2024

BACKGROUND






Age-standardized incidence rates (World) in 2020



Age-standardized mortality rates (World) in 2020



Data source: GLOBOCAN 2020
 Graph production: Global Cancer Observatory (<http://gco.iarc.fr/>)
 © International Agency for Research on Cancer 2023

International Agency for Research on Cancer
 World Health Organization

Why this study and what we know



Understanding melanoma trends is:

- crucial for prevention and treatment strategies
- allocating resources effectively and
- prepare tailored interventions based on recent data




Tumour thickness

- Tumour thickness is the most important prognostic factor
- More thinner tumours are diagnosed than thicker


Research Objective

Describe population-based long-term trends in melanoma incidence and tumour thickness as well as by sex, age, calendar period, residential region, and anatomic site.

CASE SERIES




Data from Cancer Registry of Norway



- All histologically verified first primary invasive melanoma
- **Variables:** tumour thickness, sex, age at diagnosis, vital status, anatomic site, histopathological subtype



Tumour thickness categories



Source: <https://www.cancerresearchuk.org/about-cancer/melanoma/stages-types/tnm-staging>

Basic characteristics of data

Age and Thickness

	1983-1999	2008-2019
Age at diagnosis, Median (Q1, Q3)		
Women	56 (42, 71)	63 (50, 75)
Men	59 (46, 71)	67 (56, 76)

Tumour thickness, Median (Q1, Q3)

Women	1 (0.6, 2)	0.9 (0.5, 1.8)
Men	1.3 (0.74, 2.8)	1 (0.6, 2.3)

Missing tumour decreased from 3,714 (26.3%) in 1983–1999 to 1,742 (7.7%)

Notes

Women: 23,459 (52%) and
Men: 22,065 (48%)

Increased age at diagnosis

Reduced Tumour thickness at diagnosis

Men were diagnosed at older age and thicker tumour than women

ANALYSIS & RESULTS



Methods

- Multiple imputation



Methods

- Multiple imputation
- Age-adjusted incidence rates



Methods

- Multiple imputation
- Age-adjusted incidence rates
- Segmented regression




Melanoma incidence trend

By sex and T category

Increased incidence rates in all T-categories


T1 melanoma trend

- 1983–1989/90: Steep rise
- 1990–2004: Plateau
- 2005–2019: Sharp increase again



Melanoma incidence trend

By anatomic site




Melanoma incidence trend

By anatomic site

Overall:

T1 melanoma had the highest incidence across all sites.




Melanoma incidence trend

By anatomic site

Overall:


T1 plateau was not seen for all sites.



Melanoma incidence trend

By anatomic site


In women:
High and increasing incidence on the trunk and lower limbs with plateau in T1.



Melanoma incidence trend

By anatomic site

In men:
Highest incidence in trunk but lower incidence in lower limbs than women.




Melanoma incidence trend

By anatomic site


In men:

Higher and increasing head/neck incidence compared to women, mainly in thicker melanoma.




Melanoma incidence trend

By histopathological subtype



Melanoma incidence trend


By histopathological subtype



- Superficial spreading**
- Mirrors overall pattern.
 - Higher in women than men.
 - Stable/decreasing trend in thicker melanoma.
 - Increasing T2 in the recent years.

Melanoma incidence trend

By histopathological subtype



Nodular

- Higher incidence of thicker melanoma.
- Higher in men than women.
- Stable/decreasing recent incidence trend.
- Percentage in T3 and T4 was increasing.

Summary

- Steep increase in thin melanoma until 1989/90 followed by plateau and steep increase again after 2004/05.




Summary

- Steep increase in thin melanoma until 1989/90 followed by plateau and steep increase again after 2004/05.
- Higher and more increasing incidence in men than women.



Summary

- Steep increase in thin melanoma until 1989/90 followed by plateau and steep increase again after 2004/05.
- Higher and more increasing incidence in men than women.
- Nodular melanoma had higher incidence of thicker tumour, more in men compared to women.



Summary

- Steep increase in thin melanoma until 1989/90 followed by plateau and steep increase again after 2004/05.
- Higher and more increasing incidence in men than women.
- Nodular melanoma had higher incidence of thicker tumour, more in men compared to women.
- Need for increased melanoma awareness in men.
- Need for more awareness of new and rapidly growing pigmented lesions in older people



Collaborators



UNIVERSITY
OF OSLO




[International Agency for Research on Cancer](#)



Funded by:



Coauthors




EXTRA SLIDES



Melanoma incidence trend

By period and age at diagnosis





References

- Erdmann, Friederike, Joannie Lortet-Tieulent, Joachim Schüz, Hajo Zeeb, Rüdiger Greinert, Eckhard W. Breitbart, and Freddie Bray. 2013. “International Trends in the Incidence of Malignant Melanoma 1953-2008-Are Recent Generations at Higher or Lower Risk?” *International Journal of Cancer* 132 (2): 385–400. <https://doi.org/f4hgrm>.
- Gershenwald, Jeffrey E., and Richard A. Scolyer. 2018. “Melanoma Staging: American Joint Committee on Cancer (AJCC) 8th Edition and Beyond.” *Annals of Surgical Oncology* 25 (8): 2105–10. <https://doi.org/10.1245/s10434-018-6513-7>.
- Muggeo, Vito M. R. 2008. “Segmented: An r Package to Fit Regression Models with Broken-Line Relationships.” *R News* 8 (1): 20–25. <https://cran.r-project.org/doc/Rnews/>.
- Norway, Cancer Registry of. 2023. “Cancer in Norway 2022: Cancer Incidence, Mortality, Survival and Prevalence in Norway.” 0806-3621. Cancer Registry of Norway. https://www.kreftregisteret.no/globalassets/cancer-in-norway/2022/cin_report-2022.pdf.
- Robsahm, T. E., P. Helsing, Y. Nilssen, L. Vos, S. M. H. Rizvi, L. A. Akslen, and M. B. Veierod. 2018. “High Mortality Due to Cutaneous Melanoma in Norway: A Study of Prognostic Factors in a Nationwide Cancer Registry.” *Clin Epidemiol* 10: 537–48. <https://doi.org/gdkd9q>.
- Rubin, Donald B. 2004. *Multiple Imputation for Nonresponse in Surveys*. John Wiley & Sons.