

Centre for Bariatric Surgery

Gastrointestinal Motility Laboratory

Oesophageal Manometry Report

Name **karen loader**

UR

DOB **31/01/1962**

Date **Mon, 23 Jun 2008**

Indication

Symptoms of volume reflux, also requiring PPI therapy.
Good weight loss 10 years post LAP-BAND. Also
vertical lie of LAP-BAND.

Referring Doctor **Anna Korin**

22 The Avenue Windsor 3181

Technician **Cheryl Laurie**

LOS Relaxation **Present**

Basal	Nadir	% Relx'n
Median 16.3	Median 3.6	77.7
Max 19.4	Max 3.6	
Min 4.5	Min 3.6	


Oesophageal Peristalsis **Present**

5ml water right lateral Total **9**


Normal	6
Focal Failure of Peristalsis	1
Generalised Failure of Peristalsis	2

5ml water upright Total **10**

Normal	10



Normal Comparison



Comments

Oesophageal peristalsis was generally normal. There was a high intragastric pressure at the level of the LAP-BAND of 40-50 mm Hg with 3.9 mls in the Band. A 6-7 cm high pressure zone developed above the LAP-BAND. Quite high pressures developed within the pouch.

Diagnosis

Enlarged gastric pouch post Lap-Band (intrabdominal)

P Burton/G Hebbard