

Recognizing SMART Syndrome: Imaging Evolution, Pitfalls, and Differentiation from Recurrence

Texas Society of Neuroradiology

February 21-22, 2026

Junaid Kalair, M.D.

Resident, Diagnostic Radiology
Houston Methodist Hospital

Steve H. Fung, M.D.

Houston Radiology Associated
Associate Professor of Radiology, Houston Methodist Academic Institute
Clinical Associate Professor of Radiology, Weill Cornell Medical College


HOUSTON
MethodistSM
LEADING MEDICINE

Disclosures

- None

Clinical Background


- 62-year-old-male
- Remote history:
 - Left temporo-occipital GBM
 - Gross total resection, temozolomide, radiation (60 Gy)
 - Good response without further treatment for 19 years
- Recent presentation:
 - Dysarthria, confusion, headaches, recurrent falls
 - Recurrent focal seizures during admission
 - MRI: new unilateral cortical/leptomeningeal enhancement in prior RT field



Recent Imaging Findings


Treatment with corticosteroids and verapamil

ONSET




Day 1

PROGRESSION



Day 19

RESOLUTION



Day 43

New Unilateral Enhancement in Prior RT Field


- Mimics:
 - Recurrence/leptomeningeal disease
 - Infection
 - Patient had focal seizures → peri-ictal mimic considered
- SMART discriminator:
 - Unilateral gyriform cortical \pm leptomeningeal enhancement confined to RT field
 - Improvement/resolution on follow-up imaging
- Recognition can prevent unnecessary biopsy/therapy

Findings Against Malignancy

- No mass/ nodular solid component
- Enhancement **confined to irradiated territory**
- CSF cytology negative; spine MRI negative
- Non-masslike pattern + confinement favors SMART over recurrence

Diffusion Findings


- Initial MRI: Diffuse mild cortical DWI hyperintensity with equivocal ADC (not a discrete territorial infarct)
- Follow-up MRI: New focal left occipital restricted diffusion consistent with superimposed infarct
- Interpretation: Supports SMART-related neurovascular dysregulation with possible progression to infarction (PRES-like spectrum)



SMART Syndrome

Stroke-like Migraine After Radiation Therapy

Delayed radiation-induced neurovascular dysfunction
Years–decades after cranial irradiation; often
reversible




SMART Syndrome

Imaging Findings

- Unilateral gyriform cortical \pm leptomeningeal enhancement
- Confined to prior RT field
- Does not respect vascular territories
- Predominantly cortical involvement; subcortical white matter involvement may occur and is associated with worse recovery
- Cortical T2/FLAIR hyperintensity and swelling; typically resolves on follow-up

Stepwise Assessment of New Enhancement in Prior RT Field



Avoid unnecessary biopsy and misdirected therapy

SMART Syndrome

Treatment

- Corticosteroids to empirically treat acute inflammatory changes and cerebral edema
- Calcium channel blocker such as verapamil to treat vascular dysfunction, reduce cerebral vasospasm, manage migraine-like symptoms, and prevent long-term recurrence
- Antiplatelet agents such as aspirin and blood pressure medications in cases with superimposed infarction
- Antimigraine and antiepileptic medications such as carbamazepine to control migraines and seizures
- L-arginine has been proposed to improve endothelial function and stroke-like episodes

Summary

- SMART can present decades after cranial irradiation
- Hallmark: unilateral cortical \pm leptomeningeal enhancement confined to RT field
- Often reversible; incomplete clinical or imaging recovery reported
- Steroids + verapamil; clinical improvement with near-resolution on follow-up

References

[1] Black DF, Bartleson JD, Bell ML, Lachance DH. SMART syndrome: stroke-like migraine attacks after radiation therapy. *Cephalalgia*. 2006;26(9):1137-1142. doi:10.1111/j.1468-2982.2006.01184.x

[2] Black DF, Morris JM, Lindell EP, Krecke KN, Worrell GA, Bartleson JD, et al. Stroke-like migraine attacks after radiation therapy (SMART) syndrome is not always completely reversible: a case series. *AJNR Am J Neuroradiol*. 2013;34(12):2298-2303. doi:10.3174/ajnr.A3602

[3] Zheng Q, Yang L, Tan L-M, Qin L-X, Wang C-Y, Zhang H-N, et al. Stroke-like migraine attacks after radiation therapy syndrome. *Chin Med J (Engl)*. 2015;128(15):2097-2101. doi:10.4103/0366-6999.161393

[4] Ota Y, Nemoto M, Gyoten T, Liao E, Shah G, Capizzano AA. Comprehensive update and review of clinical and imaging features of SMART syndrome. *AJNR Am J Neuroradiol*. 2023;44(6):626-639. doi:10.3174/ajnr.A7859