Quadratic $L_0 = \left(\frac{1}{2}(y - f(y))^2\right)$ - Huber (d=1) Huber (d=3) Smooth-L1 — Huber (d=5) - Huber (d=10) 20 Huber (4)301 $\frac{\mathrm{d}L_1(x)}{\mathrm{d}x} = \left\{ \begin{array}{cc} 1 & \text{if } x \ge 0 \\ -1 & \text{otherwise} \end{array} \right.$ (5)Regression 15 -error term (e) 10

Losses

Huber Loss

100

200

Quadratic $L_0 = \left(\frac{1}{2}(y - f(y))^2\right)$ - Huber (d=1) Huber (d=3) Smooth-L1 - Huber (d=10) 20 Huber (4) $\frac{\mathrm{d}L_1(x)}{\mathrm{d}x} = \left\{ \begin{array}{cc} 1 & \text{if } x \ge 0 \\ -1 & \text{otherwise} \end{array} \right.$ (5)Regression 15 error term (e) 10+ Losses

Huber Loss

100

200

Quadratic $L_0 = \int_{-2}^{1} (y - f(y))^2$ - Huber (d=1) Huber (d=3) Smooth-L1 - Huber (d=10) 20 Huber (4) $\frac{\mathrm{d}L_1(x)}{\mathrm{d}x} = \left\{ \begin{array}{cc} 1 & \text{if } x \ge 0 \\ -1 & \text{otherwise} \end{array} \right.$ (5)Regression 15 -error term (e) 10


Losses

Huber Loss


100

200


$$smooth_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1\\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$\mathrm{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$\mathrm{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$\mathrm{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$\mathrm{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise}, \end{cases}$$




$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



- Advantages of both L1 loss and L2 loss.
- Less sensitive to outliers than L2 loss
- Prevents Exploding Gradients
- Well suited for most regression tasks

- Advantages of both L1 loss and L2 loss
- Less sensitive to outliers than L2 loss
- Prevents Exploding Gradients
- Well suited for most regression tasks

- Advantages of both L1 loss and L2 loss
- Less sensitive to outliers than L2 loss
- Prevents Exploding Gradients
- Well suited for most regression tasks

- Advantages of both L1 loss and L2 loss
- Less sensitive to outliers than L2 loss
- Prevents Exploding Gradients
- Well suited for most regression tasks

- Advantages of both L1 loss and L2 loss
- Less sensitive to outliers than L2 loss
- Prevents Exploding Gradients
- Well suited for most regression tasks

- Advantages of both L1 loss and L2 loss
- Less sensitive to outliers than L2 loss
- Prevents Exploding Gradients
- Well suited for most regression tasks

- Advantages of both L1 loss and L2 loss
- Less sensitive to outliers than L2 loss
- Prevents Exploding Gradients
- Well suited for most regression tasks

$$L_2(x)=x^2$$
 $L_1(x)=|x|$ $ext{smooth}_{L_1}(x)=egin{cases} 0.5x^2 & ext{if } |x|<1 \ |x|-0.5 & ext{otherwise} \end{cases}$

$$L_1(x) = \int 1 \quad ext{if } x \geq 0$$

$$L_2(x)=x^2$$
 $L_1(x)=oxed{|x|}$ smooth $_{L_1}(x)=egin{cases} 0.5x^2 & ext{if } |x|<1 \ |x|-0.5 & ext{otherwise} \end{cases}$

$$rac{\mathrm{d}L_2(x)}{\mathrm{d}x}=2x$$

$$rac{\mathrm{d} L_1(x)}{\mathrm{d} x} = \left\{egin{array}{ll} 1 & ext{if } x \geq 0 \ -1 & ext{otherwise} \end{array}
ight.$$

$$rac{\mathrm{d} \; \mathrm{smooth}_{L_1}}{\mathrm{d} x} = \left\{ egin{array}{cc} x & \mathrm{if} \; |x| \leq 1 \ \pm 1 & \mathrm{otherwise} \end{array}
ight.$$

$$L_2(x)=x^2$$
 $L_1(x)=|x|$ $ext{smooth}_{L_1}(x)=egin{cases} 0.5x^2 & ext{if } |x|<1 \ |x|-0.5 & ext{otherwise} \end{cases}$

$$\frac{\mathrm{d}L_2(x)}{\mathrm{d}x} = 2x$$

$$rac{\mathrm{d} L_1(x)}{\mathrm{d} x} = \left\{egin{array}{ll} 1 & ext{if } x \geq 0 \ -1 & ext{otherwise} \end{array}
ight.$$

$$\frac{\mathrm{d} \; \mathrm{smooth}_{L_1}}{\mathrm{d} x} = \left\{ \begin{array}{cc} & \text{if } |x| \leq 1 \\ \pm 1 & \text{otherwise} \end{array} \right.$$

$$L_2(x)=x^2$$


$$L_1(x)=|x|$$

$$\operatorname{smooth}_{L_1}(x) = \begin{cases} \frac{0.5x^2}{|x| - 0.5} & \text{if } |x| < 1 \\ & \text{otherwise} \end{cases}$$

$$\frac{\mathrm{d}L_2(x)}{\mathrm{d}x}$$
 = 2x

$$\frac{\mathrm{d}L_1(x)}{\mathrm{d}x} = \begin{cases} 1 & \text{if } x \ge 0 \\ -1 & \text{otherwise} \end{cases}$$






$$L_2(x)=x^2$$


$$L_1(x)=|x|$$

$$\operatorname{smooth}_{L_1}(x) = \begin{cases} \frac{0.5x^2}{|x| - 0.5} & \text{if } |x| < 1 \\ \text{otherwise} \end{cases}$$

$$\frac{\mathrm{d}L_2(x)}{\mathrm{d}x} = 2x$$

$$\frac{\mathrm{d}L_1(x)}{\mathrm{d}x} = \begin{cases} 1 & \text{if } x \ge 0 \\ -1 & \text{otherwise} \end{cases}$$






$$L_2(x)=x^2$$


$$L_1(x)=|x|$$

$$\operatorname{smooth}_{L_1}(x) = \begin{cases} \frac{0.5x^2}{|x| - 0.5} & \text{if } |x| < 1 \\ & \text{otherwise} \end{cases}$$

$$\frac{\mathrm{d}L_2(x)}{\mathrm{d}x}$$
 = 2x

$$\frac{\mathrm{d}L_1(x)}{\mathrm{d}x} = \begin{cases} 1 & \text{if } x \ge 0 \\ -1 & \text{otherwise} \end{cases}$$





$$smooth_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1\\ |x| - 0.5 & \text{otherwise,} \end{cases}$$

$$smooth_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$

$$smooth_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$

Why 1?


Can I use 2 or 5?

$$smooth_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$


Why 1?

Can I use 2 or 5?

$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1, \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



$$\mathrm{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$

$$l_n = \begin{cases} 0.5(x_n - y_n)^2/\underline{beta}, & \text{if } |x_n - y_n| < \underline{beta} \\ |x_n - y_n| - 0.5 * \underline{beta}, & \text{otherwise} \end{cases}$$

$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1, \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



$$l_n = egin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \ |x_n - y_n| - 0.5 * beta, & ext{otherwise} \end{cases}$$

$$smooth_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1, \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \\ |x_n - y_n| - 0.5 * beta, & ext{otherwise} \end{cases}$$

$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1, \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$



$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \\ |x_n - y_n| - 0.5 * beta, & ext{otherwise} \end{cases}$$

$$\operatorname{smooth}_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1, \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$





$$l_n = egin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \ |x_n - y_n| - 0.5 * beta, & ext{otherwise} \end{cases}$$


$$smooth_{L_1}(x) = \begin{cases} 0.5x^2 & \text{if } |x| < 1 \\ |x| - 0.5 & \text{otherwise,} \end{cases}$$




$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & \text{if } |x_n - y_n| < beta \\ |x_n - y_n| - 0.5 * beta, & \text{otherwise} \end{cases}$$









https://github.com > torch + issues 1


Smooth L1 loss vs Huber loss · Issue #579 · torch/nn - GitHub

22-Jan-2016 — I would say that the Huber loss really is parameterised by delta, as it defines the boundary between the squared and absolute costs. I don't ...

https://github.com > torch > issues [

Smooth L1 loss vs Huber loss · Issue #579 · torch/nn - GitHub

22-Jan-2016 — I would say that the Huber loss really is parameterised by delta, as it defines the boundary between the squared and absolute costs. I don't ...



$$l_n = egin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \ |x_n - y_n| - 0.5*beta, & ext{otherwise} \end{cases}$$

Smooth-L1

$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$

$$l_n = egin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \ |x_n - y_n| - 0.5*beta, & ext{otherwise} \end{cases}$$

Smooth-L1

$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$

$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & \text{if } |x_n - y_n| < \underline{beta} \\ |x_n - y_n| - 0.5 * beta, & \text{otherwise} \end{cases}$$

Smooth-L1

$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$

$$l_n = egin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < \underbrace{beta} \ |x_n - y_n| - 0.5*beta, & ext{otherwise} \end{cases}$$

Smooth-L1

$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$

$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & \text{if } |x_n - y_n| < beta \\ |x_n - y_n| - 0.5 * beta, & \text{otherwise} \end{cases}$$

Smooth-L1



$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta*(|x_n - y_n| - 0.5*delta), & ext{otherwise} \end{cases}$$

$$l_n = egin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \ |x_n - y_n| - 0.5*beta, & ext{otherwise} \end{cases}$$


Smooth-L1

$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$

$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & \text{if } |x_n - y_n| < beta \\ |x_n - y_n| - 0.5 * beta, & \text{otherwise} \end{cases}$$

Smooth-L1

$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$



$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & ext{if } |x_n - y_n| < beta \\ |x_n - y_n| - 0.5 * beta, & ext{otherwise} \end{cases}$$

Smooth-L1

$$l_n = \begin{cases} 0.5(x_n - y_n)^2, & \text{if } |x_n - y_n| < delta \\ delta*(|x_n - y_n| - 0.5*delta), & \text{otherwise} \end{cases}$$




$$l_n = \begin{cases} 0.5(x_n - y_n)^2/beta, & \text{if } |x_n - y_n| < beta \\ |x_n - y_n| - 0.5 * beta, & \text{otherwise} \end{cases}$$

Smooth-L1


$$l_n = \begin{cases} 0.5(x_n - y_n)^2, & \text{if } |x_n - y_n| < delta \\ delta*(|x_n - y_n| - 0.5*delta), & \text{otherwise} \end{cases}$$




$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$




$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$



$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$



$$l_n = egin{cases} 0.5(x_n - y_n)^2, & ext{if } |x_n - y_n| < delta \ delta * (|x_n - y_n| - 0.5 * delta), & ext{otherwise} \end{cases}$$



Implementation

```
def smoothllloss(x, y):
    if abs(x-y)<1: return 1/2*(x-y)**2
    else: return abs(x-y)-1/2</pre>
```