

Lethality and Centrality in Protein Networks

H. Jeong, S.P. Mason, A.-L. Barabási & Z.N. Oltvai
Nature, Vol. 411, 3 May 2001 [Brief Communication]

Core Claim

The phenotypic consequence of deleting a single gene in *S. cerevisiae* is strongly predicted by the **topological position** of its protein product in the protein–protein interaction (PPI) network. Specifically: **highly connected (“hub”) proteins are disproportionately essential for survival.**

Dataset

The *S. cerevisiae* PPI network: **1,870 proteins** (nodes) connected by **2,240 direct physical interactions** (edges), derived from combined two-hybrid analyses. Lethality data from 1,572 mutants with known phenotypic profiles (Proteome database).

Network Architecture

$P(k)$ follows a **power law with an exponential cut-off** at $k_c \approx 20$:

$$P(k) \sim k^{-\gamma} \cdot e^{-k/k_c}$$

The cut-off means proteins with >20 interactions are slightly fewer than a pure scale-free prediction. The same topology is found in the *H. pylori* PPI network, suggesting **evolutionary conservation** of large-scale biological network structure. A few hub proteins mediate interactions among many low-connectivity proteins.

Key Quantitative Findings

Connectivity	Fraction of all proteins	% Essential (lethal knockout)
$k \leq 5$ (low)	$\sim 93\%$	$\sim 21\%$
$k > 15$ (hubs)	$\sim 0.7\%$	$\sim 62\%$

Pearson correlation $r = 0.75$ between lethality fraction and connectivity k .
Hub proteins are $\sim 3\times$ more likely to be essential than low-connectivity proteins.

Simulation Results

- **Random protein removal** (simulating random mutation): network diameter unaffected — consistent with yeast’s known empirical tolerance to individual gene deletions.
- **Targeted removal of most-connected proteins**: network diameter increases rapidly — confirming the hub-dependence of network integrity.
- These results directly mirror findings from the Albert et al. 2000 paper on error/attack tolerance of scale-free networks.

Mechanistic Interpretation

The robustness of yeast to random mutations is not purely a consequence of genetic redundancy or biochemical backup—it has a **topological component**. Because most proteins have few interactions, random deletions almost always remove peripherally connected proteins that do not disrupt global network structure. Hub proteins, however, are irreplaceable connectors; their loss fragments the interaction network, disrupting multiple functional modules simultaneously.

Broader Implications

- **Evolutionary argument:** The convergent appearance of scale-free topology in both metabolic and PPI networks across organisms implies *evolutionary selection* for this architecture.
- **Drug targets:** Hub proteins are prime candidates for therapeutic intervention—targeting them maximally disrupts network function (analogous to attacking hub nodes in the Internet).
- **Systems biology view:** Individual protein function and cellular robustness cannot be understood in isolation; *topological context* within the interaction network is essential.
- Future PPI studies in other organisms are predicted to reveal the same scale-free topology and the same centrality–lethality correlation.

Notable Statement

“The most highly connected proteins in the cell are the most important for its survival.”

Robustness against mutations in yeast “is also derived from the *organization* of interactions and the topological positions of individual proteins”—not just from biochemical redundancy alone.