

GATE

CRASH COURSE

ALL BRANCHES

**Engineering
Mathematics**

**Calculus (Part 02)
(Lec 05)**

By – Dr. Puneet Sharma Sir



Topics *to be covered*

① Mean Value Theorems

100% { ② Increasing Decreasing funcⁿ

③ Maxima - Minima (Single Variable)

④ " " (two variables)



Lagrange's M.V. Th.



$A(a, f(a))$

$B(b, f(b))$

Consider $f(x)$ is defined in $[a, b]$ s.t

$f(x)$ is continuous as well as Diff then

∃ at least one point c in (a, b) for which

$$\text{Slope of chord } AB = \frac{f(b) - f(a)}{b - a}$$


$$\text{Slope of tangent at } c = f'(c)$$

so A.T. Diagram,

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

ie Slope of tangent at c = Slope of chord AB .
ie tangent at c is \parallel to the chord AB

Rolle's M.V.Th



$A(a, f(a))$

$B(b, f(b))$

Consider $f(x)$ is defined $[a, b]$ s.t

$f(x)$ is Cont & Diff & also we have $f(a)=f(b)$

Then ∃ at least one c $b < c < a$ & b where

tangent is \parallel to x axis

or Horizontal tangent exist

or $f'(c) = 0$

Cauchy M.V.Th → Consider $f(x)$ & $g(x)$ are two funcⁿ defined in $[a, b]$

s.t ① Both $f(x)$ & $g(x)$ are cont $[a, b]$

② " " " are diff (a, b)

③ $g'(x) \neq 0 \forall x \in (a, b)$

then \exists exist at least one c in (a, b) for which

$$\boxed{\frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)}}$$

Q. $f(x) = \frac{\sin x}{e^x}$; $[0, \pi]$ then using R.T. the value of $c = ?$

(a) $\pi/2$

(b) $\pi/4$

(c) $\pi/6$

(d) 0

$f(x)$ is cont as well as diff (obvious)

$\leftarrow f(0) = 0 = f(\pi)$ i.e. all 3 R. Condⁿ are satisfied

So By R.T. $f'(c) = 0$

$$\frac{\cos c - \sin c}{e^c} = 0$$

$$\sin c = \cos c$$

$$\tan c = 1 \Rightarrow c = \frac{\pi}{4}$$

$$f(x) = \frac{\sin x}{e^x}$$

$$f'(x) = \frac{e^x(\cos x) - \sin x(e^x)}{(e^x)^2}$$

$$= \frac{\cos x - \sin x}{e^x}$$

Q. The Point of the Curve $f(x) = \cos x - 1$, $\left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$ where tangent is \parallel to x axis?

(a) $\left(\frac{\pi}{2}, 1\right)$

(b) $(\pi, 2)$

(c) $(\pi, -2)$

(d) $\left(\frac{\pi}{4}, \pi\right)$

it is cont as well diff (obvious) & $f\left(\frac{\pi}{2}\right) = -1 = f\left(\frac{3\pi}{2}\right)$

So By R.Th, $f'(c) = 0$

$$[-\sin c] = 0$$

$$c = n\pi, n \in \mathbb{I}$$

$$c = \dots -2\pi, -\pi, 0, \pi, 2\pi, 3\pi \dots$$

only $c = \pi$ lies in the given Domain

$$f(c) = \cos c - 1$$

$$= \cos \pi - 1 = -1 - 1 = -2$$

$$P(c, f(c)) = (\pi, -2)$$

Q. For $y = \sqrt{x-2}$, $[2, 3]$ Find the eqn of tangent which is || to chord joining end points?

Sol: $f(x) = \sqrt{x-2}$ ①

$$f'(x) = \frac{1}{2\sqrt{x-2}}$$

$$a=2, f(2)=0$$

$$b=3, f(3)=1$$

using LMVT

$$f'(c) = \frac{f(3)-f(2)}{3-2}$$

$$m = 1$$

$$\frac{1}{2\sqrt{c-2}} = \frac{1-0}{1}$$

$$\sqrt{c-2} = \frac{1}{2}$$

$$c-2 = \frac{1}{4}$$

$$c = \frac{9}{4}$$

$$\begin{aligned} f(c) &= \sqrt{c-2} \\ &= \sqrt{\frac{9}{4}-2} \\ &= \frac{1}{2} \end{aligned}$$

So Point where tangent is || to chord
 $= (c, f(c)) = \left(\frac{9}{4}, \frac{1}{2}\right)$

R. eqn of tangent is

$$y - \frac{1}{2} = m \left(x - \frac{9}{4}\right)$$

$$\frac{2y-1}{2} = \frac{4x-9}{4}$$

$$4y-4 = 8x-18$$

$$8x-4y-14=0 \Rightarrow \boxed{4x-2y=7}$$

Q2 Consider $f(x) = \ln x$ & $g(x) = \ln\left(\frac{1}{x}\right)$ defined in $[1, 2]$
 then using C.M.V.Th, $c = ?$

(a) \times 1

(b) 1.5

(c) 1.25

(d) Any value b/w 1 & 2

(e) \times 2

$$f(x) = \ln x, \quad g(x) = -\ln x$$

$$f'(x) = \left(\frac{1}{x}\right), \quad g'(x) = \left(-\frac{1}{x}\right) \Rightarrow g'(x) \neq 0 \forall x \in (1, 2)$$

$$\text{By C.M.V.Th, } \frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)}$$

$$\frac{f(2) - f(1)}{g(2) - g(1)} = \left(\frac{1/c}{-1/c}\right) \Rightarrow \frac{\ln 2 - 0}{-\ln 2 - 0} = -1 \Rightarrow \text{Identically } -1 = -1$$

$$f(x) = (1+x) \log(1+x); [0, 1]$$

$$a=0, f(a)=f(0)=0$$

$$b=1, f(b)=2 \log 2$$

$$\text{i.e. } (a, f(a)) = (0, 0)$$

$$(b, f(b)) = (1, 2 \log 2)$$

Slope of chord AB is

$$m = \frac{2 \log 2 - 0}{1 - 0}$$

$$f'(c) = 2 \log 2$$

By applying Lagrange's mean value for the function $f(x) = (1+x) \log(1+x)$ on $[0, 1]$ the value of $c \in (0, 1)$ is

(a) $\frac{4}{e}$

(b) $\frac{1}{e}$

(c) $\frac{4-e}{e}$

(d) $\frac{1-e}{e}$

$$f'(c) = 2 \log 2$$

$$1 + \log(1+c) = 2 \log 2$$

$$\log(1+c) = \log(2)^2 - 1$$

$$\log(1+c) = \log 4 - \log e$$

$$\log(1+c) = \log\left(\frac{4}{e}\right)$$

$$f(x) = (1+x) \log(1+x)$$

$$f'(x) = (1+x) \left[\frac{1}{1+x} \right] + \log(1+x) [1]$$

$$= 1 + \log(1+x)$$

$$1+c = \frac{4}{e}$$


$$c = \frac{4}{e} - 1 = \frac{4-e}{e}$$

MAXIMA-MINIMA (of single variable)

⊗ W.K. that for $y=f(x)$, $\frac{dy}{dx} = \boxed{f'(x) = \tan \theta}$ = Slope of tangent at any Random point x



$$\boxed{f'(x) > 0} \text{ always}$$




$$\boxed{f'(x) < 0}$$

& for Horizontal tangent

$$\boxed{f'(x) = 0}$$

Hence for Inc. funcⁿ $\Leftrightarrow f'(x) > 0$ always
 & for Dec funcⁿ $\Leftrightarrow f'(x) < 0$ "



Local Max points = C, E, B

" " Values = $f(C), f(E), f(B)$



Max. Value = $f(E)$

Local min points = A, D, F

" " Values = $f(A), f(D), f(F)$

Min Value = $f(F)$

Short Cut Method to check Max-Min:
(use the sign of $f'(x)$)



Turning Points / Critical Point

Put $f'(x) = 0$ & try to find x

say it is α then

$x = \alpha$ is called T-Point.

1st Condⁿ for Maxima-Minima is $f'(x) = 0$

(*) Corner points are not T-Points (True)

(*) Extreme points → points where Maxima or minima occurs called E-Points.

(*) Corner points are Extreme points (T)

(*) Shortcut Method of finding Maxima or Minima is applicable only for T-Points.

(*) Max-Min occurs alternately (T)

(*) Long Method:-

Let $x = \alpha$ is the T-Point then

$f'(\alpha) = 0$
 $\begin{cases} f''(\alpha) < 0 \Rightarrow \text{Maxima} \\ f''(\alpha) > 0 \Rightarrow \text{Minima} \end{cases}$

The maximum value of
 $f(x) = x^3 - 9x^2 + 24x + 5$ in the interval $[1, 6]$ is

- (a) 21 (b) 25
 (c) 41 (d) 46

$$f'(x) = 3x^2 - 18x + 24$$


$$= 3(x^2 - 6x + 8)$$

$$f'(x) = 3(x-4)(x-2)$$

∴ Points are $f'(x) = 0$

$$x = 2 \text{ \& } 4$$


No. of Extremes = four



$$f(2) = 25$$

$$f(6) = 41$$


So Points of Max $\nearrow x=2$
 $\nwarrow x=6$



The maximum value of $(-\infty, \infty)$
 $f(x) = x^3 - 9x^2 + 24x + 5$ is ____.

$$f'(x) = 3(x-4)(x-2)$$

$$x = 2, 4$$



No. of Extremes = two



- (a) 21
- (b) 25
- (c) 41
- (d) DNE



The maximum value of $f(x) = 2x^3 - 9x^2 + 12x - 3$ in the interval $0 \leq x \leq 3$ is 6.

$$f'(x) = 6x^2 - 18x + 12$$

$$= 6(x^2 - 3x + 2)$$

$$f'(x) = 6(x-1)(x-2)$$

For T-Points, $f'(x) = 0$
 $x = 1 \text{ \& } 2$



So Points of Maxima are $x = 1 \text{ \& } 3$

& Local Max. values $\rightarrow f(1) = 2$

$f(3) = 6$

So Absolute Max Value = 6 & it occurs at $x = 3$

No. of Extremas = 4
 (min & Max Both)

A point on the curve is said to be an extremum if it is a local minimum (or) a local maximum. The number of distinct extrema for the curve $3x^4 - 16x^3 + 24x^2 + 37$ is _____.

(a) 0

(b) 1

(c) 2

(d) 3

it only $x=0$ is Extreme point.

$$f(x) = 3x^4 - 16x^3 + 24x^2 + 37 ; (-\infty, \infty)$$

$$f'(x) = 12x^3 - 48x^2 + 48x$$

$$= 12x(x^2 - 4x + 4)$$

$$f'(x) = 12x(x-2)^2$$

T. Points; $x=0, 2$

if we have only one Extrema which occurs at $x=0$



$$f'(-1) = 12(-1)(-1-2)^2 = -ve$$

$$f'(1) = 12(1)(1-2)^2 = +ve$$

$$f'(3) = 12(3)(3-2)^2 = +ve$$

The minimum value of the function $f(x) = \frac{1}{3}x(x^2 - 3)$ in the interval $-100 \leq x \leq 100$ occurs at $x = \underline{\hspace{2cm}}$.

- (a) 1
(b) -1
(c) -100
(d) None

$$f(x) = \frac{1}{3}x(x^2 - 3) = \frac{1}{3}(x^3 - 3x)$$

$$f'(x) = \frac{1}{3}[3x^2 - 3] = x^2 - 1$$

T-Points, Put $f'(x) = 0$
 $x = \pm 1$




$$f'(x) = (x+1)(x-1)$$

Minima occur at $\begin{cases} x = 1 \\ x = -100 \end{cases}$

Min Values $\begin{cases} f(1) = -2/3 \\ f(-100) = < < < 0 \end{cases}$

Maxima-Minima of funcⁿ of two variables

$$[Z = f(x, y)] \begin{cases} \text{Point} = (x, y) \\ \text{Value} = Z \end{cases}$$



1. Condⁿ for Maxima-Minima

$$\boxed{\frac{\partial Z}{\partial x} = 0 \quad , \quad \frac{\partial Z}{\partial y} = 0}$$

① ②

Critical Points → Points obtained by solving (1) & (2) are called critical points. let after solving we get $x=a$ & $y=b$ then $P(a,b)$ is c.p.

S. Condⁿ for Max-Minima → funcⁿ must satisfy Lagrange's Condⁿ.

Lagrange's Condⁿ Let $\left(\frac{\partial^2 z}{\partial x^2}\right)_p = r$, $\left(\frac{\partial^2 z}{\partial x \partial y}\right)_p = s$, $\left(\frac{\partial^2 z}{\partial y^2}\right)_p = t$

(1) if $rt - s^2 > 0$ & $r > 0$ then $p(a, b)$ is point of Minima

(2) if $rt - s^2 > 0$ & $r < 0$ " " " " Maxima

(3) if $rt - s^2 < 0$ then " " " " Inflexion (\approx Saddle Point)

(4) if $rt - s^2 = 0$ then Not possible to check Nature of point in terms of Maxima or Minima

(*) Saddle point \Rightarrow we have Neither Maxima nor minima
 \Leftarrow

$$z = f(x, y) = 4x^2 + 6y^2 - 8x - 4y + 8$$

$$\frac{\partial z}{\partial x} = 8x - 8, \quad \frac{\partial^2 z}{\partial x^2} = 8$$

$$\frac{\partial z}{\partial y} = 12y - 4, \quad \frac{\partial^2 z}{\partial y^2} = 12$$

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial y} \right) = \frac{\partial}{\partial x} (12y - 4) = 0$$

Critical Points ;

$$\frac{\partial z}{\partial x} = 0 \quad \& \quad \frac{\partial z}{\partial y} = 0$$

$$x = 1 \quad \& \quad y = \frac{1}{3}$$

Given a function $f(x, y) = 4x^2 + 6y^2 - 8x - 4y + 8$. The optimum value of $f(x, y)$ is

- (a) a minimum equal to $10/3$
- (b) a maximum equal to $10/3$
- (c) a minimum equal to $8/3$
- (d) a maximum equal to $8/3$

Optimum Value

→ Extreme Value

{ Min Value & Max Value both }

ie $P(1, \frac{1}{3})$ is Critical Point.

$$\begin{aligned} r &= (Z_{xx})_p = 8 \\ s &= (Z_{xy})_p = 0 \\ t &= (Z_{yy})_p = 12 \end{aligned} \quad \left. \begin{aligned} &\because r = 8 > 0 \\ &\& \Delta = r^2 - s^2 = (8)^2 - (0)^2 \\ &= 64 > 0 \end{aligned} \right\} \text{So } P(1, \frac{1}{3}) \text{ is Point of Minimum}$$

$$\text{Min Value}(z) = (f(x, y))_p = [4x^2 + 6y^2 - 8x - 4y + 8]_p = \dots = \frac{10}{3}$$

The total cost (C_T) of an equipment in terms of the operation variables x and y is

$$C_T = 2x + \frac{12000}{xy} + y + 5$$

The optimal value of C_T , rounded to 1 decimal place, is _____.

$$Z = 2x + \frac{12000}{xy} + y + 5$$

$$Z_x = 2 - \frac{12000}{x^2 y}$$

$$Z_y = -\frac{12000}{xy^2} + 1$$

For C. Points: $\frac{\partial Z}{\partial x} = 0$ & $\frac{\partial Z}{\partial y} = 0$

$$2 - \frac{12000}{x^2 y} = 0 \quad \& \quad -\frac{12000}{xy^2} + 1 = 0$$

$$\boxed{x^2 y = 6000} \quad \& \quad \boxed{xy^2 = 12000}$$

$$\frac{x^2 y}{xy^2} = \frac{6000}{12000} \Rightarrow \frac{x}{y} = \frac{1}{2}$$

if $x:y = 1:2$ $\begin{cases} x = k \text{ (let)} \\ y = 2k \end{cases}$

For Bq ①, $(k)^2(2k) = 6000 \Rightarrow k^3 = 3000$

$k = 14.4$ $\begin{cases} x = k = 14.4 \\ y = 2k = 28.8 \end{cases}$

$$Z = 2(14.4) + \frac{12000}{(14.4)(28.8)} + 28.8 + 5 = 91.5$$

The word 'Thank' is written in a large, bold, yellow, cursive-style font. A yellow arrow starts from the top of the 'T', extends horizontally to the right, and then curves downwards to point at the end of the word 'Thank'.

Thank

THANK

Keep Hustling!

