

GATE

DATA SCIENCE + CS & IT

**Engineering
Mathematics**

SUPER 1500

Lec : 05

Calculus

By – Dr. Puneet Sharma Sir



Topics *to be covered*

CALCULUS

- ① Maxima - Minima of Curve ($y = f(x)$)
- ② " " of Surface ($z = f(x, y)$)




MAXIMA-MINIMA of $y=f(x)$

① Point = x & Value = y , ② N-Cond: $f'(x) = 0$ or undefined

③ S-Cond: (M-I) 1st Derivative test (M-II) 2nd Derivative test.

④ we will also check the Maxima-Minima at corner points.



⑤ Concave up: $f''(x) > 0$ & Concave Down $f''(x) < 0$

⑥ S-Inc: $f'(x) > 0$ & S-Dec: $f'(x) < 0$

⑦ Saddle Point \Rightarrow ~~N.M.N.M~~

The maximum value of

$y = f(x) = x^3 - 9x^2 + 24x + 5$ in the interval $[1, 6]$ is

(a) 21

(b) 25

(c) 41


(d) 46

$$f'(x) = 3x^2 - 18x + 24$$

$$= 3(x^2 - 6x + 8)$$

$$f'(x) = 3(x-4)(x-2)$$


∴ Points are $x = 2, 4$



Max Value will occur

at $x = 2, f(2) = 25$

$x = 6, f(6) = 41$



The maximum value of $(-\infty, \infty)$
 $f(x) = x^3 - 9x^2 + 24x + 5$ is ____.


(a) 21 (b) 25

(c) 41 (d) ~~DNE~~

∴ Points are $x = 2 \text{ \& \; } 4$

$$f'(x) = 3(x-2)(x-4)$$

∴ Local Max Value



$$f'(1) = +ve$$

$$f'(3) = -ve$$

$$f'(5) = +ve$$

$$= f(2) = 25$$

Q

Max Value of $f(x) = -x^2 + 4x - 4$ will be?

(a) ~~∞~~ (b) 0
 (c) 25 (d) 2

Qs Max Value of $f(x) = -x^2 + 4x - 4$ will be (a) ∞ (b) 0

Sol: $f(x) = -(x^2 - 4x + 4)$, $(-\infty, \infty)$ (c) 25 (d) 2


$$y = f(x) = -(x-2)^2$$

Max value occurs at $x=2$


$$\& \text{ Value} = f(2) = 0$$

(M-II)

$$f'(x) = -2(x-2) \text{ at T.P in } x=2$$



Q2 the Max Value of $y = x^2$ in $[1, 5]$ will be _____



Min Value = 1

Max " = 25

The maximum value of $f(x) = 2x^3 - 9x^2 + 12x - 3$ in the interval $0 \leq x \leq 3$ is _____.


$$f(x) = 2x^3 - 9x^2 + 12x - 3$$

$$f'(x) = 6x^2 - 18x + 12$$

$$= 6(x^2 - 3x + 2)$$

$$f'(x) = 6(x-1)(x-2)$$

$$\therefore \text{Pare } x = 1 \text{ \& } 2$$



$$f(1) = 2 - 9 + 12 - 3 = 2$$

$$f(3) = 2(27) - 9(9) + 12(3) - 3 = 6$$

A point on the curve is said to be an extremum if it is a local minimum (or) a local maximum. The number of distinct extrema for the curve $3x^4 - 16x^3 + 24x^2 + 37$ is _____.

(a) 0

☒ (b) 1

(c) 2

(d) 3

$$f(x) = 3x^4 - 16x^3 + 24x^2 + 37$$

$$f'(x) = 12x^3 - 48x^2 + 48x$$

$$= 12x(x^2 - 4x + 4)$$

$$f'(x) = 12x(x-2)^2$$

∴ Points are $x = 0, 2, 2$



The range of values of k for which the function

$f(x) = (k^2 - 4)x^2 + 6x^3 + 8x^4$ has a local maxima

at point $x = 0$ is

- (a) $k < -2$ or $k > 2$ (b) $k \leq -2$ or $k \geq 2$
 (c) $-2 < k < 2$ (d) $-2 \leq k \leq 2$

$$f(x) = (k^2 - 4)x^2 + 6x^3 + 8x^4$$

$$f'(x) = (k^2 - 4)(2x) + 18x^2 + 32x^3$$

$$f''(x) = (k^2 - 4)(2) + 36x + 96x^2$$

At $x=0$, using N.Cand^y $f'(0) = 0$

$$0 = 0$$

S. Cand^y at $(x=0)$ is $f''(0) < 0$

$$2(k^2 - 4) + 0 + 0 < 0$$

$$(k-2)(k+2) < 0$$


$$-2 < k < 2$$

Consider the function $f(x) = 2x^3 - 3x^2$ in the domain $[-1, 2]$. The global minimum of $f(x)$ is _____

$$f(x) = x^2(2x - 3)$$

$$f'(x) = 6x^2 - 6x = 6x(x - 1)$$

T.P $x = 0, 1$




L-min = $f(-1) = -5 = \text{global min}$
L-min = $f(1) = -1 = \text{local min}$

As x varies from -1 to $+3$, which one of the following describes the behavior of the function $f(x) = x^3 - 3x^2 + 1$?

- (a) $f(x)$ increases monotonically.
- (b) $f(x)$ increases, then decreases and increases again.
- (c) $f(x)$ decreases, then increases and decreases again.
- (d) $f(x)$ increases and then decreases.


$$f(x) = x^3 - 3x^2 + 1$$
$$f'(x) = 3x^2 - 6x$$
$$f'(x) = 3x(x-2)$$

FP at $x = 0, 2$



Let $\max\{a, b\}$ denote the maximum of two real numbers a and b . Which of the following statement(s) is/are **TRUE** about the function $f(x) = \max\{3 - x, x - 1\}$?

- ☒ (a) It is continuous on its domain.
- ☒ (b) It has a local minimum at $x = 2$.
- ☒ (c) It has a local maximum at $x = 2$.
- ☒ (d) It is **differentiable** on its domain.




$$y = 3 - x \quad \& \quad y = x - 1$$

$$x + y = 3 \qquad x - y = 1$$

$$\frac{x}{3} + \frac{y}{3} = 1 \qquad \frac{x}{1} + \frac{y}{-1} = 1$$

$$x - 1 = 3 - x \Rightarrow x = 2 \quad \& \quad y = 1$$



Local Minima :

A, D, F, B

Global "

D

For real values of x , the minimum value of the function $f(x) = \exp(x) + \exp(-x)$ is

- (a) 2 (b) 1
(c) 0.5 (d) 0

$$f(x) = e^x + e^{-x}$$

$$= e^x + \frac{1}{e^x}$$

$$= R^+ + \frac{1}{R^+}$$

$$\boxed{f(x) \geq 2} \Rightarrow f_{\min} = 2$$

Th: if $x \in R^+$ then $\boxed{x + \frac{1}{x} \geq 2}$

if $x=1$ then $1 + \frac{1}{1} = 2$

if $x=3$ then $3 + \frac{1}{3} > 2$


if $x=2$ then $2 + \frac{1}{2} > 2$

if $x=0.2$, then $0.2 + \frac{1}{0.2} = 0.2 + 5 > 2$
.....

The maximum area (in square units) of a rectangle whose vertices lie on the ellipse $x^2 + 4y^2 = 1$ is _____.

(M-I) $x^2 + 4y^2 = 1$

$$\frac{x^2}{1^2} + \frac{y^2}{(1/2)^2} = 1$$



$$A = (AB) \times (AD) = (2x)(2y) = 4xy \quad \text{--- (1)}$$

let $u = A^2 = 16x^2y^2$ where $A = \sqrt{u}$
or $A_{\max} = \sqrt{u_{\max}} = ?$

Now Try to Calculate Max Value of u

$$u = 16x^2y^2$$

$$= 16(x^2(1-x^2))$$

$$u = 4(x^2 - x^4) \quad \text{--- (2)}$$

$$\frac{du}{dx} = 0$$

$$4(2x - 4x^3) = 0 \quad \left| \quad x \neq 0, \pm \frac{1}{\sqrt{2}} \right.$$


$$8x(1 - 2x^2) = 0 \quad \left| \quad \frac{d^2u}{dx^2} = (8 - 48x^2) \right.$$

$$1 \left(\frac{d^2 u}{dx^2} \right)_{x=\frac{1}{\sqrt{2}}} = -ve \Rightarrow u \text{ will be max at } x = \frac{1}{\sqrt{2}}$$

$$\hookleftarrow u_{\text{max}} = \left(4(x^2 - x^4) \right)_{x=\frac{1}{\sqrt{2}}} = 4 \left(\frac{1}{2} - \frac{1}{4} \right) = 4 \left(\frac{2-1}{4} \right) = 1$$

$$\therefore A_{\text{max}} = \sqrt{u_{\text{max}}} = \sqrt{1} = 1$$

M-II



Sol. $x^2 + 4y^2 = 1 \Rightarrow \frac{x^2}{1^2} + \frac{y^2}{(1/2)^2} = 1$

∴ Random Point of Ellipse
is $P(\cos \theta, \frac{1}{2} \sin \theta)$


$$\text{Area} = L \times B = (2 \cos \theta)(\sin \theta) = \sin 2\theta$$

$$\boxed{A = \sin 2\theta} \Rightarrow \frac{dA}{d\theta} = 2 \cos 2\theta, \quad \frac{d^2A}{d\theta^2} = -4 \sin 2\theta$$

T-Points $\frac{dA}{d\theta} = 0 \Rightarrow 2\theta = \frac{\pi}{2} \Rightarrow \theta = \frac{\pi}{4}$

$\left(\frac{d^2A}{d\theta^2}\right)_{\theta = \frac{\pi}{4}} = -4 < 0$ ∴ $\theta = \frac{\pi}{4}$ is Point of Maxima

∴ (Max Value of A) _{$\theta = \frac{\pi}{4}$} $= \sin 2\left(\frac{\pi}{4}\right) = 1$



For a right angled triangle, if the sum of the length of the hypotenuse and a side is kept constant, in order to have maximum area of the triangle, the angle between the hypotenuse and the side is

- (a) 120° (b) 60°
(c) 30° (d) 45°

$B = x, \theta = ?$

ATQ $\boxed{\sqrt{x^2 + y^2} + x = k} \text{ (const)}$

$$x^2 + y^2 = (k - x)^2$$

$$x^2 + y^2 = k^2 + x^2 - 2kx$$

$$y^2 = k^2 - 2kx \quad \text{--- (1)}$$

$$A = \frac{1}{2}xy \Rightarrow u = A^2 = \frac{1}{4}(x^2y^2) = \frac{1}{4}x^2(k^2 - 2kx)$$

for Max Area, u must be Max $\because A = \sqrt{u}$

$$\Rightarrow \frac{du}{dx} = 0$$

$$\frac{1}{4}(2k^2x - 2k(3x^2)) = 0$$

$$kx - 3x^2 = 0$$

$$x = k/3 \Rightarrow y = \frac{k}{\sqrt{3}}$$

$$\tan \theta = \frac{y}{x} = \frac{k/\sqrt{3}}{k/3}$$

$$\tan \theta = \sqrt{3} \Rightarrow \theta = 60^\circ$$

The best approximation of the minimum value attained by $e^{-x} \sin(100x)$ for $x > 0$ is _____

$$f(x) = e^{-x} \sin(100x)$$

w.k. that e^{-x} is always +ve i.e. $e^{-x} > 0$ i.e. Min Value of $e^{-x} > 0$

So Min Value of $f(x)$ depends on $\sin(100x)$

& w.k. that, Min of $\sin(100x) = -1$ ($\because -1 \leq \sin x \leq 1$)

$$\sin 100x = \sin\left(\frac{3\pi}{2}\right)$$

$$100x = \frac{3\pi}{2}$$

i.e. Point of Minimum, $x = \frac{3\pi}{200}$



$$\begin{aligned} \& \text{ Min Value} &= f\left(\frac{3\pi}{200}\right) = e^{-\frac{3\pi}{200}}(-1) \\ &= -e^{-\frac{3\pi}{200}} = -0.954 \end{aligned}$$

Maxima-Minima of funcⁿ of two variables

S. Condⁿ

$$[Z = f(x, y)] \begin{cases} \rightarrow \text{Point} = (x, y) \\ \rightarrow \text{Value} = Z \end{cases}$$

① if $rt - s^2 > 0$ & $r > 0$ then $P(a, b)$ is point of minima

② if $rt - s^2 > 0$ & $r < 0$ then ... of Maxima

③ if $rt - s^2 < 0$ then ... of Inflexion

④ if $rt - s^2 = 0$ then case fails

N. Condⁿ $\frac{\partial Z}{\partial x} = 0$ & $\frac{\partial Z}{\partial y} = 0$

The total cost (C_T) of an equipment in terms of the operation variables x and y is

$$C_T = 2x + \frac{12000}{xy} + y + 5$$

The optimal value of C_T , rounded to 1 decimal place, is ____.

$$Z_x = 2 - \frac{12000}{x^2 y}$$

$$Z_{xx} = \frac{24000}{x^3 y}$$

$$Z_y = -\frac{12000}{xy^2} + 1$$

$$Z_{yy} = \frac{24000}{xy^3}$$

$$Z_{xy} = \frac{12000}{x^2 y^2}$$

$$Z_x = 0 \text{ \& } Z_y = 0$$

$$2 - \frac{12000}{x^2 y} = 0 \text{ \& } -\frac{12000}{xy^2} + 1 = 0$$

$$x^2 y = 6000 \text{ \& } xy^2 = 12000$$

$$\boxed{\frac{x}{y} = \frac{1}{2}}$$

$$\begin{aligned} x &= \alpha \text{ (let)} \\ y &= 2\alpha \text{ (let)} \end{aligned}$$

$$\text{By (1); } 2 = \frac{12000}{\alpha^3}$$

$$\alpha^3 = 6000 \Rightarrow \alpha = 14.2$$

$$\text{Hence } x = \alpha = 14.3$$

$$y = 2x = 28.6$$

$$\begin{aligned} \text{Hence } C_T &= \left(2x + \frac{12000}{xy} + y + 5 \right) \\ &= 91.5 \end{aligned}$$

(14.2, 28.4)

The function $f(x, y) = 2x^4 + y^2 - x^2 - 2y$ has a relative_____.

- (a) maxima at $\left(\frac{1}{2}, 1\right)$ (b) ☒ minima at $\left(\frac{1}{2}, 1\right)$
 (c) maxima at $(0, 1)$ (d) minima at $(0, 1)$

C. Points are: $\frac{\partial z}{\partial x} = 0$ & $\frac{\partial z}{\partial y} = 0$

$$2x(4x^2 - 1) = 0 \quad \& \quad 2(y - 1) = 0$$

$$x = 0, \pm \frac{1}{2}, \quad y = 1$$

So C. Points are $(0, 1), \left(\frac{1}{2}, 1\right), \left(-\frac{1}{2}, 1\right)$

At $P(0, 1) \rightarrow r = (24x - 2)_P = -2, t = 2, s = 0$
 $\therefore rt - s^2 = (-2)(2) - (0)^2 < 0 \Rightarrow$ saddle point.

At $Q\left(\frac{1}{2}, 1\right) \rightarrow r = (24x - 2)_Q = 10, t = 2, s = 0$
 $rt - s^2 = (10)(2) - (0)^2 > 0 \quad \& \quad r > 0$

$$Z = 2x^4 + y^2 - x^2 - 2y$$

$$\frac{\partial Z}{\partial x} = 8x^3 - 2x, \quad \frac{\partial^2 Z}{\partial x^2} = 24x - 2$$

$$\frac{\partial Z}{\partial y} = 2y - 2, \quad \frac{\partial^2 Z}{\partial y^2} = 2$$

$$\frac{\partial^2 Z}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial Z}{\partial y} \right) = \frac{\partial}{\partial x} (2y - 2) = 0$$

Find the absolute maxima and minima of the function *value*

$z = f(x, y) = x^2 - xy - y^2 - 6x + 2$ on the rectangular plate $0 \leq x \leq 5$, $-3 \leq y \leq 0$

$$\frac{\partial z}{\partial x} = (2x - y - 6), \quad \frac{\partial^2 z}{\partial x^2} = 2$$


$$\frac{\partial z}{\partial y} = (-x - 2y), \quad \frac{\partial^2 z}{\partial y^2} = -2$$

$$\frac{\partial^2 z}{\partial x \partial y} = -1$$

C. points are $\frac{\partial z}{\partial x} = 0$ & $\frac{\partial z}{\partial y} = 0$
 $2x - y - 6 = 0$ & $-x - 2y = 0$

$$\begin{cases} 2x - y = 6 \\ x + 2y = 0 \end{cases} \Rightarrow x = \frac{12}{5}, y = -\frac{6}{5} \text{ is } P\left(\frac{12}{5}, -\frac{6}{5}\right) \text{ C. Point}$$

Now At P; $xt - s^2 < 0$ so $P\left(\frac{12}{5}, -\frac{6}{5}\right)$ is saddle point



Corner points are
 $(0,0)$, $(5,0)$, $(0,-3)$, $(5,-3)$

$$f(x, y) = x^2 - xy - y^2 - 6x + 2$$

$$f(0, 0) = 0 - 0 - 0 - 0 + 2 = 2$$

$$f(5, 0) = 25 - 0 - 0 - 30 + 2 = -3$$

$$f(0, -3) = 0 - 0 - 9 - 0 + 2 = -7 \quad (\text{Global Min Value}) \text{ \& it occurs at } (0, -3)$$

$$f(5, -3) = 25 + 15 - 9 - 30 + 2 = 3 \quad (\text{ " Max Value }) \text{ \& " " at } (5, -3)$$

The word 'Thank' is written in a large, bold, yellow, cursive-style font. A yellow arrow starts from the top of the 'T', extends horizontally to the right, and then curves downwards to point at the end of the word 'Thank'.

Thank

THANK



Keep Hustling!