

# Species population

## Solving the Lotka-Volterra equations

Andreas Evensen

Computational Physics  
Stockholm University

May 28, 2024

# Table of contents

- 1 Introduction
- 2 Theory
- 3 Method
- 4 Result
- 5 Further investigation
- 6 Conclusion

# Introduction

How does the reproduction of any species depend on its environment?  
What are the factors that makes a species population thrive, or decline.

# Introduction

How does the reproduction of any species depend on its environment?  
What are the factors that makes a species population thrive, or decline.

## ① Nutrition

# Introduction

How does the reproduction of any species depend on its environment?  
What are the factors that makes a species population thrive, or decline.

- 1 Nutrition
- 2 Predators

# Introduction

How does the reproduction of any species depend on its environment?  
What are the factors that makes a species population thrive, or decline.

- 1 Nutrition
- 2 Predators
- 3 Climate

How does the reproduction of any species depend on its environment?  
What are the factors that makes a species population thrive, or decline.

- ① Nutrition
  - ② Predators
  - ③ Climate
- Can one use this information to model the population of any two or more species?

Suppose we want to model a system of two species,  $X$  and  $Y$ .



Suppose we want to model a system of two species,  $X$  and  $Y$ .

- 1 Species  $X$  has unlimited food.

Suppose we want to model a system of two species,  $X$  and  $Y$ .

- ① Species  $X$  has unlimited food.
- ② The food source of species  $Y$  is entirely species  $X$ .

Suppose we want to model a system of two species,  $X$  and  $Y$ .

- 1 Species  $X$  has unlimited food.
- 2 The food source of species  $Y$  is entirely species  $X$ .
- 3 The rate of which the population for the two species increases/decreases is proportional to their current population.

Suppose we want to model a system of two species,  $X$  and  $Y$ .

- ① Species  $X$  has unlimited food.
- ② The food source of species  $Y$  is entirely species  $X$ .
- ③ The rate of which the population for the two species increases/decreases is proportional to their current population.
- ④ The ecological properties remains unchanged.

Suppose we want to model a system of two species,  $X$  and  $Y$ .

- 1 Species  $X$  has unlimited food.
- 2 The food source of species  $Y$  is entirely species  $X$ .
- 3 The rate of which the population for the two species increases/decreases is proportional to their current population.
- 4 The ecological properties remains unchanged.
- 5 Species  $Y$  always 'hunt' species  $X$ .

Suppose we want to model a system of two species,  $X$  and  $Y$ .

- 1 Species  $X$  has unlimited food.
- 2 The food source of species  $Y$  is entirely species  $X$ .
- 3 The rate of which the population for the two species increases/decreases is proportional to their current population.
- 4 The ecological properties remains unchanged.
- 5 Species  $Y$  always 'hunt' species  $X$ .
- 6 Each of the two species population are described by only time, i.e. the entire population of  $Y$  partake in 'hunting' species  $X$ .

These assumptions gives rise to the following equations:

$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xy \\ \frac{dy}{dt} &= \delta xy - \gamma y.\end{aligned}\tag{1}$$

These assumptions gives rise to the following equations:

$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xy \\ \frac{dy}{dt} &= \delta xy - \gamma y.\end{aligned}\tag{1}$$

- A pair of nonlinear first order differential equation named **Lotka-Volterra equations**, or Pray-Predator equations.



These assumptions gives rise to the following equations:

$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xy \\ \frac{dy}{dt} &= \delta xy - \gamma y.\end{aligned}\tag{1}$$

- A pair of nonlinear first order differential equation named **Lotka-Volterra equations**, or Pray-Predator equations.
- Can be modified by introducing time-dependent variables,  $\alpha(t)$ ,  $\beta(t)$ ,  $\delta(t)$ , and  $\gamma(t)$ .

These assumptions gives rise to the following equations:

$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xy \\ \frac{dy}{dt} &= \delta xy - \gamma y.\end{aligned}\tag{1}$$

- A pair of nonlinear first order differential equation named **Lotka-Volterra equations**, or Pray-Predator equations.
- Can be modified by introducing time-dependent variables,  $\alpha(t)$ ,  $\beta(t)$ ,  $\delta(t)$ , and  $\gamma(t)$ .
- Can be modified to the Arditi–Ginzburg equations.

Rewriting our system we obtain:

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = \underbrace{\begin{pmatrix} \alpha - \beta^* y & -\beta^* x \\ \delta^* y & \delta^* x - \gamma \end{pmatrix}}_{\mathbf{J}^*(x,y)} \cdot \begin{pmatrix} x \\ y \end{pmatrix}, \quad (2)$$

where  $\beta^* = 0.5\beta$ ,  $\delta^* = 0.5\delta$ .

Rewriting our system we obtain:

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = \underbrace{\begin{pmatrix} \alpha - \beta^* y & -\beta^* x \\ \delta^* y & \delta^* x - \gamma \end{pmatrix}}_{\mathbf{J}^*(x,y)} \cdot \begin{pmatrix} x \\ y \end{pmatrix}, \quad (2)$$

where  $\beta^* = 0.5\beta$ ,  $\delta^* = 0.5\delta$ .

- 1 Fixed point at  $(0,0)$  (Saddle point).

Rewriting our system we obtain:

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = \underbrace{\begin{pmatrix} \alpha - \beta^* y & -\beta^* x \\ \delta^* y & \delta^* x - \gamma \end{pmatrix}}_{\mathbf{J}^*(x,y)} \cdot \begin{pmatrix} x \\ y \end{pmatrix}, \quad (2)$$

where  $\beta^* = 0.5\beta$ ,  $\delta^* = 0.5\delta$ .

- 1 Fixed point at  $(0, 0)$  (Saddle point).
- 2 Fixed point at  $\left(\frac{\gamma}{\delta^*}, \frac{\alpha}{\beta^*}\right)$  (Oscillatory behavior).

# Method

Solutions to eq 1, can be obtained via Explicit Euler, or any order Runge-Kutta method.

# Method

Solutions to eq 1, can be obtained via Explicit Euler, or any order Runge-Kutta method.



$$k_1 = f(t_n, x_n)$$


$$k_2 = f\left(t_n + \frac{dt}{2}, x_n + dt \cdot \frac{k_1}{2}\right)$$

$$k_3 = f\left(t_n + \frac{dt}{2}, x_n + dt \cdot \frac{k_2}{2}\right)$$

$$k_4 = f(t_n + dt, x_n + dt \cdot k_3)$$


$$x_{n+1} = x_n + \frac{dt}{6} (k_1 + 2k_2 + 2k_3 + k_4). \quad (3)$$

## Comparison of the two method





## Comparison of the two method




- Choose Runge-Kutta method with  $dt = 0.001$ .


# Solutions

Variables:  $\alpha = 1.0$ ,  $\beta = 0.1$ ,  $\delta = 0.8$ , and  $\gamma = 1.4$ .

Initial condition:  $x(0) = 8$ ,  $y(0) = 10$ .



(a) Population in time




(b) Phase diagram

Figure: Population and phase diagram


# Solutions

Variables:  $\alpha = 1.0$ ,  $\beta = 0.5$ ,  $\delta = 1.0$ , and  $\gamma = 0.9$ .

Initial condition:  $x(0) = 10$ ,  $y(0) = 5$ .



(a) Population in time




(b) Phase diagram

Figure: Population and phase diagram


# Solutions

Variables:  $\alpha = 1.1$ ,  $\beta = 0.5$ ,  $\delta = 0.3$ , and  $\gamma = 0.9$ .

Initial condition:  $x(0) = 10$ ,  $y(0) = 5$ .



(a) Population in time




(b) Phase diagram

Figure: Population and phase diagram


# Extinction

Variables:  $\alpha = 1.5$ ,  $\beta = 0.1$ ,  $\delta = 0.1$ , and  $\gamma = 0.1$ .

Initial condition:  $x(0) = 0$ ,  $y(0) = 10$ .



(a) Population in time



(b) Phase diagram

Figure: Population and phase diagram

# New set of equations

Suppose we instead want three species,  $X$ ,  $Y$ , and  $Z$ .

# New set of equations

Suppose we instead want three species,  $X$ ,  $Y$ , and  $Z$ .

- Species  $Y$  lives in protection of species  $X$ .

# New set of equations

Suppose we instead want three species,  $X$ ,  $Y$ , and  $Z$ .

- Species  $Y$  lives in protection of species  $X$ .
- 

$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xz \\ \frac{dy}{dt} &= \eta xy - \chi yz \\ \frac{dz}{dt} &= \delta xz - \gamma z.\end{aligned}\tag{4}$$



# Three species

This gives rise to a problem.

# Three species

This gives rise to a problem.

- Problem becomes more stiff.

# Three species


This gives rise to a problem.

- Problem becomes more stiff.


- | $\alpha$ | $\beta$ | $\delta$ | $\gamma$ | $\eta$ | $\chi$ |
|----------|---------|----------|----------|--------|--------|
| 1.1      | 0.5     | 0.3      | 0.9      | 0.1    | 0.1    |

# Three species solution

Initial condition:  $x(0) = 10$ ,  $y(0) = 1$ , and  $z(0) = 5$ .



(a) Population in time



(b) Phase diagram

Figure: Population and phase diagram

# Generalized Lotka-Volterra equations

Suppose now instead of protection, we say that one herbivore and two carnivores.

$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xy - \zeta xy \\ \frac{dy}{dt} &= \eta xy - \chi yz \\ \frac{dz}{dt} &= \delta xz + \iota yz - \gamma z.\end{aligned}\tag{5}$$


# Generalized Lotka-Volterra equations

Suppose now instead of protection, we say that one herbivore and two carnivores.


$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xy - \zeta xy \\ \frac{dy}{dt} &= \eta xy - \chi yz \\ \frac{dz}{dt} &= \delta xz + \iota yz - \gamma z.\end{aligned}\tag{5}$$

$\alpha$	$\beta$	$\delta$	$\gamma$	$\eta$	$\chi$	$\eta$	$\iota$
1.1	0.5	0.3	0.9	0.1	0.1	0.1	0.1

# Generalized Lotka-Volterra equations



(a) Population in time



(b) Phase diagram

Figure: Population and phase diagram for generalized Lotka-Volterra solution

# Time dependent variables

- Can we modify the equation to take into account seasonal changes?



- Can we modify the equation to take into account seasonal changes?
- 

$$\begin{aligned}\frac{dx}{dt} &= \alpha(t)x - \beta(t)xy \\ \frac{dy}{dt} &= \delta(t)xy - \gamma(t)y.\end{aligned}\tag{6}$$

# Time dependent variables

How do we then model the seasons?

# Time dependent variables

How do we then model the seasons?




$$\alpha(t) = \alpha_1 \cos(t) + \alpha_1$$

$$\beta(t) = H^{-1}(t) \cdot \beta_1$$


$$\delta(t) = H^{-1}(t) \cdot \delta_1$$

$$\gamma(t) = H(t) \cdot \gamma_1$$

# Time dependent variables



(a) Population in time



(b) Phase diagram

Figure: Population and phase diagram with time dependent variables

# Conclusion

What have we learned?

# Conclusion

What have we learned?

- Solution methods to a pair of nonlinear first order differential equations.

# Conclusion

What have we learned?

- Solution methods to a pair of nonlinear first order differential equations.
- Generalization to more than two equations.

What have we learned?

- Solution methods to a pair of nonlinear first order differential equations.
- Generalization to more than two equations.
- The unperturbed model is fixed in an oscillatory behavior.



What have we learned?

- Solution methods to a pair of nonlinear first order differential equations.
- Generalization to more than two equations.
- The unperturbed model is fixed in an oscillatory behavior.
- The model becomes more stiff when introducing more variables.

What have we learned?

- Solution methods to a pair of nonlinear first order differential equations.
- Generalization to more than two equations.
- The unperturbed model is fixed in an oscillatory behavior.
- The model becomes more stiff when introducing more variables.
- Naive implementation of seasons brakes symmetry

What have we learned?

- Solution methods to a pair of nonlinear first order differential equations.
- Generalization to more than two equations.
- The unperturbed model is fixed in an oscillatory behavior.
- The model becomes more stiff when introducing more variables.
- Naive implementation of seasons brakes symmetry
- Questions?

Thanks for listening