

## 7. ER Pelvic Protocol

### **ER Pelvic Protocol (This need to be supervised by staff/facilitator/Second Call)**

#### **Without Contrast**

3 PL localizer free breathing  
T2 Haste sagittal  
T2 Haste coronal  
T2 Haste axial  
T2 Haste fs coronal  
T2 Haste fs Axial  
T2 Haste fs Sagittal  
T2 TrueFISP axial breath hold  
T1 Vibe Dixon fs axial (4 points)  
Diffusion b50, 800

### **(Don't inject pregnant patients)- This need to be supervised by staff/facilitator/Second Call**

#### **With Contrast**

3 PL localizer free breathing  
T2 Haste sagittal  
T2 Haste coronal  
T2 Haste axial  
T2 Haste fs coronal  
T2 Haste fs Axial  
T1 Vibe Dixon fs axial (4 points)  
Diffusion b50, 800  
T1 VIBE fs axial breath hold  
T1 VIBE fs sagittal breath hold  
T1 VIBE fs coronal breath hold

#### **Contrast**

T1 VIBE fs axial +c (50s)  
T1 VIBE fs axial +c (90s)  
T1 VIBE fs sagittal+c (2 min)  
T1 VIBE fs coronal+c (3 min)  
T1 tse axial +c ( 4 min)