



BIO 160 : INVERTEBRATE SYSTEMATICS/ ZOOLOGY

LESSON 1. INTRODUCTION / PROTISTA CONCEPT

ARKOH, M. A

THEORETICAL AND APPLIED BIOLOGY DEPT
FACULTY OF BIOSCIENCES, C. O. S



Edit with WPS Office

**3 CREDIT COURSE; 2 LECTURE
HOURS, 3 PRACTICAL HOURS PER
WEEK**

**COURSE OUTLINE:
Prostista concept.**

Systematics of selected Inverts -
- Their Classification,
- Phylogeny,



www.knust.edu.gh



Edit with WPS Office

- Biology,
- Adaptive radiation
- and Economic importance.

INVERTS: Protozoa, Porifera, Coelenterata (Cnidaria & Ctenophora), Platyhelminthes/ Nematini and Aschelminthes/ Nematoda

REFERENCES

Monger, G. & Sangster (2010).

Systematics and classification.

Kershaw, R.D(2012). Animal diversity.

Pichenik, J.A (2015). Invertebrate Zoology

Hickmann, Larsen & Roberts (2015). Animal diversity.



Cleveland, Larry & Allan (2014).
Animal biology.

Purves, Orians & Heller (2015).
Life (science of biology).

Barnes, Callow & Goldwig (2012)
. The invertebrate



INVERTEBRATE ZOOLOGY

Organisms are classified under 5 broad Kingdoms.

Monera, Protocista, Plantae,
Fungi and Animalia

Inverts are animals without bones!

Sub-kingdom Protozoa is an animal though, could be

considered as unicellular



PROTISTA CONCEPT

Before life started, and living things grouped into Kingdoms, the world was void.

Then under high pressure under water depth, simple elements (H, N, C, P, etc) combined to form monomers (Coz, PO₄, NO₂, etc)



The monomers continued to combine into polymers (higher molecular compounds) till a DNA/RNAs were formed. This was around 4.5 billions year ago.

This phenomenon was chemogenesis.

The Polymers then started pulsating.



www.knust.edu.gh



Edit with WPS Office

Then around 3 billion years, the DNAs/RNAs building food around themselves to be able to get energy from simple substances.

They then built a wall around this food store.

Resulted in Prokaryotic cell formation.



www.knust.edu.gh

Around 1500-1400BC, the primitive cells grouped their scattered nuclei materials together and build a wall around it for protection and easy coordination.

Eukaryotic cells then, appeared.



www.knust.edu.gh



Edit with WPS Office

These cells then became complex through cell multiplications, tissue and organ formations.

The resulting organisms then adapted to water and other diverse environments through body modifications till the human species was reached.



Early systematics put together a dendrogram for pictorial presentation.

Whittaker started in 1969,
then followed by Margulis
1982 and Barnes in 1984



Classification of Kingdom

Animalia

KINGDOM : ANIMALIA

3 SUB-KINGDOMS : PROTOZOA

MESOZOA

METAZOA

*PROTOZOA :

SARCOMASTIGOPHORA,

CILIOPHORA & APICOMPLEXA.



www.knust.edu.gh



Edit with WPS Office

MEZOZOA : all organisms are extinct. Falls worms.

METAZOA: PARAZOA & EUMATOZOA

-Parazoa – multicellular but loose and uncoordinated cells.
Has only 1 phylum ; Porifera

EUMATOZOA : Invertebrata & Vertebrata

***INVERTEBRATA** : Radiata & Bilateria

-RADIATA : Coelenterata (Cnidaria & Ctenophora)

-BILATERIA : Acoelomata, Pseudo-coelomata & Coelomata

ACOELOMATA : Platyhelminthes
& Nemertini

PSEUDO-COELOMATA :
Nematoda & Aschelminthes

COELOMATA : Annelida,
Arthropoda & Molluska



www.knust.edu.gh



Edit with WPS Office

PHYLOGENETIC TREE OF INVERTS

draw !



www.knust.edu.gh



Edit with WPS Office

Thanks



www.knust.edu.gh



Edit with WPS Office