

MATH157: ALGEBRA

CHAPTER 2

Indices - Logarithm and Exponential Functions

February 01, 2021



Outline

1 Indices

- Index form
- Index laws

2 Exponential Functions

- Exponential equations
- Exponential Graphs
- Exponential Functions

3 Logarithm Functions



$8 = 2 \times 2 \times 2$ can be rewritten as 2^3 . This is read as '2 (raised) to the power 3.

Definition

Let x be a real number. If there exist two real numbers a and n such that $x = a^n$, then

- *the expression a^n is the **index form** of x ;*
- *a is called the **base**;*
- *n is called the **index** (or the power/exponent).*



Property (Index laws)

Let n and m be two integers, a and b two real numbers.

$$(1) a^n \times a^m = a^{n+m}.$$

$$(2) \frac{a^n}{a^m} = a^{n-m} \text{ provided } a \neq 0$$

$$(3) (a^n)^m = a^{nm}.$$

$$(4) (ab)^m = a^m b^m.$$

$$(5) \left(\frac{a}{b}\right)^m = \frac{a^m}{b^m} = a^m b^{-m} \text{ provided } b \neq 0.$$

$$(6) a^0 = 1 \text{ provided } a \neq 0.$$



Application of (1) $a^n \times a^m = a^{n+m}$.

Example

$$3^4 \times 3^5 =$$



Application of (1) $a^n \times a^m = a^{n+m}$.

Example

$$3^4 \times 3^5 = 3^9.$$

$$11^{-4} \times 11^5 =$$



Application of (1) $a^n \times a^m = a^{n+m}$.

Example

$$3^4 \times 3^5 = 3^9.$$

$$11^{-4} \times 11^5 = 11^1.$$

$$4^{-2} \times 4^{-1} =$$



Application of (1) $a^n \times a^m = a^{n+m}$.

Example

$$3^4 \times 3^5 = 3^9.$$

$$11^{-4} \times 11^5 = 11^1.$$

$$4^{-2} \times 4^{-1} = 4^{-3}.$$

Application of (2) $\frac{a^n}{a^m} = a^{n-m}$.

Example

$$\frac{3^4}{3^5} =$$



Application of (1) $a^n \times a^m = a^{n+m}$.

Example

$$3^4 \times 3^5 = 3^9.$$

$$11^{-4} \times 11^5 = 11^1.$$

$$4^{-2} \times 4^{-1} = 4^{-3}.$$

Application of (2) $\frac{a^n}{a^m} = a^{n-m}$.

Example

$$\frac{3^4}{3^5} = 3^{4-5} = 3^{-1}.$$

$$\frac{11^{-4}}{11^5} =$$



Application of (1) $a^n \times a^m = a^{n+m}$.

Example

$$3^4 \times 3^5 = 3^9.$$

$$11^{-4} \times 11^5 = 11^1.$$

$$4^{-2} \times 4^{-1} = 4^{-3}.$$

Application of (2) $\frac{a^n}{a^m} = a^{n-m}$.

Example

$$\frac{3^4}{3^5} = 3^{4-5} = 3^{-1}.$$

$$\frac{11^{-4}}{11^5} = 11^{-4-5} = 11^{-9}.$$

$$\frac{4^{-2}}{4^{-1}} =$$



Application of (1) $a^n \times a^m = a^{n+m}$.

Example

$$3^4 \times 3^5 = 3^9.$$

$$11^{-4} \times 11^5 = 11^1.$$

$$4^{-2} \times 4^{-1} = 4^{-3}.$$

Application of (2) $\frac{a^n}{a^m} = a^{n-m}$.

Example

$$\frac{3^4}{3^5} = 3^{4-5} = 3^{-1}.$$

$$\frac{11^{-4}}{11^5} = 11^{-4-5} = 11^{-9}.$$

$$\frac{4^{-2}}{4^{-1}} = 4^{-2+1} = 4^{-1}.$$



Application of (4) $(a^n)^m = a^{nm}$.

Example

$$(3^4)^5 =$$



Application of (4) $(a^n)^m = a^{nm}$.

Example

$$(3^4)^5 = 3^{4 \times 5} = 3^{20}.$$

$$(11^{-4})^5 =$$



Application of (4) $(a^n)^m = a^{nm}$.

Example

$$(3^4)^5 = 3^{4 \times 5} = 3^{20}.$$

$$(11^{-4})^5 = 11^{-20}.$$

$$(4^{-2})^{-1} =$$



Application of (4) $(a^n)^m = a^{nm}$.

Example

$$(3^4)^5 = 3^{4 \times 5} = 3^{20}.$$

$$(11^{-4})^5 = 11^{-20}.$$

$$(4^{-2})^{-1} = 4^2.$$



Property

$$(7) \ a^{\frac{1}{n}} = \sqrt[n]{a} \text{ and } a^{\frac{m}{n}} = (\sqrt[n]{a})^m \text{ provided } a \geq 0.$$

Example

$$9^{\frac{3}{2}} =$$



Property

$$(7) \ a^{\frac{1}{n}} = \sqrt[n]{a} \text{ and } a^{\frac{m}{n}} = (\sqrt[n]{a})^m \text{ provided } a \geq 0.$$

Example

$$9^{\frac{3}{2}} = (\sqrt{9})^3 = 27 \text{ (or } (3^2)^{\frac{3}{2}} = 3^3 = 27).$$

$$32^{\frac{3}{5}} =$$



Property

(7) $a^{\frac{1}{n}} = \sqrt[n]{a}$ and $a^{\frac{m}{n}} = (\sqrt[n]{a})^m$ provided $a \geq 0$.

Example

$$9^{\frac{3}{2}} = (\sqrt{9})^3 = 27 \text{ (or } (3^2)^{\frac{3}{2}} = 3^3 = 27).$$

$$32^{\frac{3}{5}} = (2^5)^{\frac{3}{5}} = 8.$$

Exercise

- Find $\frac{3^3 \times 3^7}{3^2}$, $\frac{2^{-3} \times 2^7}{4^2}$ and $(27 \times 16)^{\frac{2}{3}}$
- Show that $\frac{8a^2b^2}{3a^3b} \div \frac{4ab^2}{9a^3b^5} = 6ab^4$ and $36 \times 2^{m-2} + 2^{m+3} - 2^m = 2^{m+4}$.
- Express in the simplest product of index forms: (a) $\frac{8^3 \times 5^4 \times 7^3}{5^3 \times 7^5 \times 2^6}$,
 (b) $\frac{(0.6)^2 \times 12^5 \times 54^3}{9^2 \times 5^3 \times (0.8)^3 \times (0.4)^4}$, (c) $\frac{(3^4 \times 2^{-3})^3}{(9^{-1} \times 2^2)^4} \cdot \frac{10^2 \times 3^2}{8 \times 5^2} \div \sqrt{\frac{2^5 \times 3^9}{6}}$,



Outline

1 Indices

- Index form
- Index laws

2 Exponential Functions

- Exponential equations
- Exponential Graphs
- Exponential Functions

3 Logarithm Functions



Definition

*Equations of the form $a^x = b$ are called **exponential equations**.*

If $a > 0$ and the index form of b is a^n , then $a^x = a^n$ implies that $x = n$.

Example

Solve (1) $2^x = 8$ and (2) $3^{-4x} - 1 = 8$.



Definition

*Equations of the form $a^x = b$ are called **exponential equations**.*

If $a > 0$ and the index form of b is a^n , then $a^x = a^n$ implies that $x = n$.

Example

Solve (1) $2^x = 8$ and (2) $3^{-4x} - 1 = 8$.

(1) Since $8 = 2^3$, $2^x = 8 \implies 2^x = 2^3 \implies x = 3$.



Definition

*Equations of the form $a^x = b$ are called **exponential equations**.*

If $a > 0$ and the index form of b is a^n , then $a^x = a^n$ implies that $x = n$.

Example

Solve (1) $2^x = 8$ and (2) $3^{-4x} - 1 = 8$.

(1) *Since $8 = 2^3$, $2^x = 8 \implies 2^x = 2^3 \implies x = 3$.*

(2) $3^{-4x} - 1 = 8 \implies 3^{-4x} = 3^2 \implies -4x = 2 \implies x = -1/2$

Exercise

Find x for which (a) $3^{-2x} = 27^{x+1}$, (b) $5^{3x} = 5^{7x-2}$, (c) $5^{5-9x} = \frac{1}{25^{x-2}}$,

(d) $21^{-1+2x} - 21^{x^2} = 0$, (e) $4^x \times 3^{2x} = 6$, (f) $4^{x+1} + 2^{2x} = 20$.

Find x and y solutions of $2^{5x+1} = 16^{y+1}$ and $2^{x-13} = 32^{1-y}$.



Activity

Use the calculator to find out the values of 2^x for $x \in \{-4, -3, -1, 0, 1, 2, 3\}$. Show the points of coordinates $(x, 2^x)$ on a graph.



Activity

Use the calculator to find out the values of 2^x for $x \in \{-4, -3, -1, 0, 1, 2, 3\}$. Show the points of coordinates $(x, 2^x)$ on a graph.

Solution




Figure: Exponential graph



Definition

- The real number $e = 2.7182818284590452353602874713527 \dots$ is called **Euler's number**.

The function that associates to any real number x , the real number $y = e^x$ is called **exponential function**.



Definition

- The real number $e = 2.7182818284590452353602874713527 \dots$ is called **Euler's number**.

The function that associates to any real number x , the real number $y = e^x$ is called **exponential function**.

Property

$$e^0 = 1, e^1 = e \text{ and } e^x > 0, x \in \mathbb{R}.$$

$$e^x e^y =$$



Definition

- The real number $e = 2.7182818284590452353602874713527 \dots$ is called **Euler's number**.

The function that associates to any real number x , the real number $y = e^x$ is called **exponential function**.

Property

$$e^0 = 1, e^1 = e \text{ and } e^x > 0, x \in \mathbb{R}.$$

$$e^x e^y = e^{x+y},$$

$$(e^x)^y =$$



Definition

- The real number $e = 2.7182818284590452353602874713527 \dots$ is called **Euler's number**.

The function that associates to any real number x , the real number $y = e^x$ is called **exponential function**.

Property

$$e^0 = 1, e^1 = e \text{ and } e^x > 0, x \in \mathbb{R}.$$

$$e^x e^y = e^{x+y},$$

$$(e^x)^y = e^{xy},$$

$$e^{-x} =$$



Definition

- The real number $e = 2.7182818284590452353602874713527 \dots$ is called **Euler's number**.

The function that associates to any real number x , the real number $y = e^x$ is called **exponential function**.

Property

$$e^0 = 1, e^1 = e \text{ and } e^x > 0, x \in \mathbb{R}.$$

$$e^x e^y = e^{x+y},$$

$$(e^x)^y = e^{xy},$$

$$e^{-x} = \frac{1}{e^x}.$$

Exercise

Solve (a) $e^{\frac{1}{2}x^2+1} = e^{x\sqrt{2}}$ and (b) $e^{2x} - 3e^x + 2 = 0$.



Outline

1 Indices

- Index form
- Index laws

2 Exponential Functions

- Exponential equations
- Exponential Graphs
- Exponential Functions

3 Logarithm Functions



Definition (Natural logarithm function)

Natural logarithm function \ln is the inverse-function of the exponential function e .

That is, if $y > 0$,

Exponential form

$$e^x = y$$



Natural logarithm form

$$x = \ln y$$

For any two positive real numbers x and y , we have

$$(a) \ln(xy) = \ln x + \ln y, (b) \ln(x^n) = n \ln x, \text{ and } (c) \ln\left(\frac{x}{y}\right) = \ln x - \ln y.$$

For instance

$$e^{x-1} = 3 \implies (\ln [e^{x-1}] = \ln 3 \implies)x - 1 = \ln 3 \implies x = 1 + \ln 3.$$

$$\text{Solve in } \mathbb{R} : e^{x^2-4x+1+\ln 6} = 6.$$



Definition (Logarithm function)

If b and x are two positive real numbers, then **base b logarithm** of x is denoted by $\log_b(x)$ and is defined by $\log_b(x) = \frac{\ln x}{\ln b}$.

- $b = e \implies \log_e x = \frac{\ln x}{\ln e} = \ln x$.
- $b = 10 \implies \log_{10} x = \frac{\ln x}{\ln 10} = \log x$.
- If $b \neq e$ and $b \neq 10$ we use \log_b .

Note that $\log_b 1 = 0$ and $\log_b b = 1$.

Property (Conversion)

For $b > 0$ and $y > 0$,

Exponential form

$$b^x = y$$

$$\iff$$

Logarithmic form

$$x = \log_b y.$$



Example

The logarithmic form of :

$$5^2 = 25 \text{ is } 2 = \log_5 25$$

$$11^x = 21 \text{ is } x = \log_{11} 21.$$

Property

For $b > 0, x > 0$ and $y > 0$

$$(a) \log_b 1 = 0, \log_b b = 1 \text{ and } \log_b x = \frac{\ln x}{\ln b}.$$

$$(b) \log_b(xy) = \log_b x + \log_b y,$$

$$(c) \log_b \left(\frac{x}{y} \right) = \log_b x - \log_b y.$$

$$(d) \log_b(x^n) = n \log_b x.$$



Example

We want to simplify (1) $\log_4 10 + \log_4 2$,

(2) $\log_2(\sqrt{3} + 1) + \log_2(\sqrt{3} - 1)$ *and expand* (3) $\log \left(\frac{10x^4}{3y^2} \right)$.

We have

(1) $\log_4 10 + \log_4 2 =$



Example

We want to simplify (1) $\log_4 10 + \log_4 2$,

(2) $\log_2(\sqrt{3} + 1) + \log_2(\sqrt{3} - 1)$ *and expand* (3) $\log \left(\frac{10x^4}{3y^2} \right)$.

We have

(1) $\log_4 10 + \log_4 2 = \log_4(10 \times 2) = \log_4(20)$.

(2) $\log_2(\sqrt{3} + 1) + \log_2(\sqrt{3} - 1) =$



Example

We want to simplify (1) $\log_4 10 + \log_4 2$,

(2) $\log_2(\sqrt{3} + 1) + \log_2(\sqrt{3} - 1)$ *and expand* (3) $\log \left(\frac{10x^4}{3y^2} \right)$.

We have

$$(1) \log_4 10 + \log_4 2 = \log_4(10 \times 2) = \log_4(20).$$

$$(2) \log_2(\sqrt{3} + 1) + \log_2(\sqrt{3} - 1) = \log_2 \left[(\sqrt{3} + 1)(\sqrt{3} - 1) \right] = \log_2(2) = 1.$$

$$(3) \log \left(\frac{10x^4}{3\sqrt{y}} \right) =$$



Example

We want to simplify (1) $\log_4 10 + \log_4 2$,

(2) $\log_2(\sqrt{3} + 1) + \log_2(\sqrt{3} - 1)$ *and expand* (3) $\log \left(\frac{10x^4}{3y^2} \right)$.

We have

$$(1) \log_4 10 + \log_4 2 = \log_4(10 \times 2) = \log_4(20).$$

$$(2) \log_2(\sqrt{3} + 1) + \log_2(\sqrt{3} - 1) = \log_2 [(\sqrt{3} + 1)(\sqrt{3} - 1)] = \log_2(2) = 1.$$

$$(3) \log \left(\frac{10x^4}{3\sqrt{y}} \right) = \log(10x^4) - \log(3\sqrt{y}) = \log 10 + 4 \log x - \log 3 - \frac{1}{2} \log y$$

$$= 1 - \log 3 + 4 \log x - \frac{1}{2} \log y$$



Exercise

Solve

(1) $(x - 2) \log_4(x - 2) = 0,$

(2) $\log(x^2 + x - 1) = 0,$

(3) $x - y = 2 - b$ *and* $\log_b(x + y) = 1$ *provided* $b > 0.$

Given $\log_7 2 = \alpha, \log_7 3 = \beta$ *and* $\log_7 5 = \gamma$, *express in terms of* α, β *and* γ

(4) $\log_7 6.$

(5) $\log_7 \frac{15}{2}.$

