

Real Analysis Assignment 6

Joel Savitz

October 20, 2020

Problem 1:

Claim 1. *The minimal subsequential limit of some bounded infinite sequence x_n can be computed by the formula $\sup_k (\inf_{n>k} x_n)$*

Proof. Since x_n is bounded, it is bounded above. Denote the upper bound N , then $\forall n, x_n \leq N$. Define a subsequence $y_k = \inf_{n>k} x_n$. Then, y_1 is the greatest lower bound of $\{x_2, x_3, x_4, \dots\}$. Since $\{x_3, x_4, x_5, \dots\}$ is a subset of $\{x_2, x_3, x_4, \dots\}$, we have that y_1 is also a lower bound of the latter set, but since $y_2 = \inf_{n>2} x_n$, we must have $y_1 \leq y_2$. For some y_k , we have y_{k-1} is a lower bound of $\{x_{k+1}, x_{k+2}, \dots\}$, since $y_{k-1} = \inf_{n>k-1} x_n$, but since $y_k = \inf_{n>k} x_n$, we must have $y_k \geq y_{k-1}$. Thus $\forall k, y_k \leq y_{k+1}$, i.e. y_k is monotone increasing. Then since $\forall n > k, y_k \leq x_n$ and $\forall n, x_n \leq N$, we have by transitivity that $y_k \leq N$. Thus y_k is bounded above. Since y_k is a monotone increasing sequence bounded above, we have by the monotone convergence theorem that $\lim y_k$ exists, and it is equal to $\sup y_k$. Let $A = \lim y_k$. Since y_k has a limit A every subsequence of y_k has the same limit A .

Now, pick some $n_1 \in \mathbb{N}$. Then $y_{n_1} = \inf\{x_{n_1}, x_{n_1+1}, x_{n_1+2}, \dots\}$. Since y_{n_1} is the greatest lower bound of $\{x_{n_1}, x_{n_1+1}, x_{n_1+2}, \dots\}$, we must have that $y_{n_1} + 1$ is not a lower bound of that set. Then, since $y_{n_1} + 1$ is not a lower bound, $\exists n_2 > n_1, y_{n_1} \leq x_{n_2} < y_{n_1} + 1$. Continuing in the same way, we have $y_{n_2} = \inf\{x_{n_2}, x_{n_2+1}, x_{n_2+2}, \dots\}$, and since y_{n_2} is the greatest lower bound of the set, we have that $y_{n_2} + \frac{1}{2}$ is not a lower bound of the set, thus $\exists n_3 > n_2, y_{n_2} \leq x_{n_3} < y_{n_2} + \frac{1}{2}$. We can continue this indefinitely, such that for any n_{k-1} , we have that $y_{n_{k-1}} = \inf\{x_{n_{k-1}}, x_{n_{k-1}+1}, x_{n_{k-1}+2}, \dots\}$, and so $y_{n_{k-1}} + \frac{1}{k-1}$ is not a lower bound of the set, and thus $\exists n_k > n_{k-1}, y_{n_{k-1}} \leq x_{n_k} < y_{n_{k-1}} + \frac{1}{k-1}$. Applying the squeeze theorem and limit laws, we have:

$$\lim(y_{n_{k-1}}) \leq \lim x_{n_k} < \lim(y_{n_{k-1}} + \frac{1}{k-1}) \quad (1)$$

$$\lim(y_{n_{k-1}}) \leq \lim x_{n_k} < \lim(y_{n_{k-1}}) + \lim(\frac{1}{k-1}) \quad (2)$$

$$A \leq \lim x_{n_k} < A + 0 \quad (3)$$

$$\implies \lim x_{n_k} = A \quad (4)$$

Thus A is a subsequential limit of x_n .

Consider some other subsequential limit of x_n , suppose $\lim x_{m_k} = B$. Then:

$$y_{m_k-1} = \inf\{x_{m_k}, x_{m_k+1}, x_{m_k+2}, \dots\} \quad (5)$$

$$\forall m_k, y_{m_k-1} \leq x_{m_k} \quad (6)$$

$$\lim y_{m_k-1} \leq \lim x_{m_k} \quad (7)$$

$$A \leq B \quad (8)$$

Therefore, A is the minimal subsequential limit of x_n , and it is the value of $\sup_k (\inf_{n>k} x_n)$. \square

Problem 2:

Claim 2. If a_n and b_n are Cauchy sequences, then $a_n b_n$ is a Cauchy sequence.

Proof. If a sequence is Cauchy, then it is bounded, and if a sequence is bounded, then it is bounded above. Thus, $\exists A, B \in \mathbb{R}$, where $\forall n, a_n \leq A$ and $\forall n, b_n \leq B$. By the definition of a Cauchy sequence, we have that

$$\forall \epsilon > 0, \exists N_\epsilon, \forall n > N_\epsilon, m > N_\epsilon, |a_n - a_m| < \epsilon \quad (9)$$

and likewise

$$\forall \epsilon > 0, \exists M_\epsilon, \forall n > M_\epsilon, m > M_\epsilon, |b_n - b_m| < \epsilon \quad (10)$$

Let $\epsilon > 0$. Thus we have the following:

$$\exists N_{\frac{\epsilon}{2|B|}}, \forall n > \frac{\epsilon}{2|B|}, m > \frac{\epsilon}{2|B|}, |a_n - a_m| < \frac{\epsilon}{2|B|} \quad (11)$$

$$\exists M_{\frac{\epsilon}{2|A|}}, \forall n > \frac{\epsilon}{2|A|}, m > \frac{\epsilon}{2|A|}, |b_n - b_m| < \frac{\epsilon}{2|A|} \quad (12)$$

Let $K_\epsilon = \max(N_{\frac{\epsilon}{2|B|}}, M_{\frac{\epsilon}{2|A|}})$. Then, $\forall n > K_\epsilon, m > K_\epsilon$, we have:

$$|B||a_n - a_m| < \frac{\epsilon}{2} \quad (13)$$

$$|A||b_n - b_m| < \frac{\epsilon}{2} \quad (14)$$

$$\implies |B||a_n - a_m| + |A||b_n - b_m| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \quad (15)$$

Since $\forall n, a_n \leq A$ and $\forall n, b_n \leq B$, we have:

$$|B||a_n - a_m| + |A||b_n - b_m| \geq |b_m||a_n - a_m| + |a_n||b_n - b_m| \quad (16)$$

$$|b_m||a_n - a_m| + |a_n||b_n - b_m| = |a_n b_n - a_n b_m| + |a_n b_m - a_m b_m| \quad (17)$$

And by the triangle inequality:

$$|a_n b_n - a_n b_m| + |a_n b_m - a_m b_m| \geq |a_n b_n - a_n b_m + a_n b_m - a_m b_m| \quad (18)$$

$$|a_n b_n - a_n b_m + a_n b_m - a_m b_m| = |a_n b_n - a_m b_m| \quad (19)$$

Therefore, by transitivity, $|a_n b_n - a_m b_m| < \epsilon$. Thus $\forall \epsilon > 0, \exists K_\epsilon, \forall n > K_\epsilon, m > K_\epsilon, |a_n b_n - a_m b_m| < \epsilon$, and this is exactly the definition that $a_n b_n$ is a Cauchy sequence. \square

Problem 3:

Define a sequence $x_n = \sqrt{n}$.

Part (a):

Claim 3. $\lim(x_{n+2} - x_n) = 0$

Proof. Let $\epsilon > 0$. By the Archimedian property, $\exists N_{\frac{4}{\epsilon^2}} \in \mathbb{N}, N_{\frac{4}{\epsilon^2}} > \frac{4}{\epsilon^2}$. Thus, $\forall n > N_{\frac{4}{\epsilon^2}}, n > \frac{4}{\epsilon^2}$. Therefore:

$$\frac{4}{n} < \epsilon^2 \quad (20)$$

$$\frac{2}{\sqrt{n}} < \epsilon \quad (21)$$

$$\frac{2}{\sqrt{2+n} + \sqrt{n}} < \frac{2}{\sqrt{n}} \quad (22)$$

$$\frac{2}{\sqrt{2+n} + \sqrt{n}} = \frac{n+2-n}{\sqrt{2+n} + \sqrt{n}} \quad (23)$$

$$= \sqrt{n+2} - \sqrt{n} \quad (24)$$

If we assume that for some natural n , $\sqrt{n+2} \leq \sqrt{n}$, we see that $n+2 \leq n$ and thus $2 \leq 0$, which is absurd, therefore $\sqrt{n+2} > \sqrt{n}$, and so $\sqrt{n+2} - \sqrt{n} = |\sqrt{n+2} - \sqrt{n}|$. By the transitivity of the above order relations, we then have $|\sqrt{n+2} - \sqrt{n}| = |\sqrt{n+2} - \sqrt{n} - 0| < \epsilon$. Thus we have demonstrated that $\forall \epsilon > 0, \exists N_\epsilon, \forall n > N_\epsilon, |\sqrt{n+2} - \sqrt{n} - 0| < \epsilon$, which is exactly the definition that $\lim(x_{n+2} - x_n) = 0$. \square

Part (b):

Claim 4. $x_n = \sqrt{n}$ is not a Cauchy sequence.

Proof. Let $\epsilon = 1$ and let N be any natural number. Define $n = 4(N+1)^2$ and $m = (N+1)^2$. We see that:

$$|x_n - x_m| = |\sqrt{n} - \sqrt{m}| = |\sqrt{4(N+1)^2} - \sqrt{(N+1)^2}| \quad (25)$$

$$|2(N+1) - (N+1)| \quad (26)$$

$$|2N+2 - N-1| \quad (27)$$

$$|N+1| \quad (28)$$

Since $N \geq 1$, we have $N+1 \geq 1+1 = 2 \geq 1 = \epsilon$. Thus we have demonstrated that $\exists \epsilon > 0, \forall N, \exists n > N, m > N, |x_n - x_m| \geq \epsilon$, which is the negation of the definition of a Cauchy sequence, therefore $x_n = \sqrt{n}$ is not a Cauchy sequence. \square