

# HIGHER SCHOOL CERTIFICATE TRIAL EXAMINATION 2020



## Chemistry

### General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Draw diagrams using pencil
- Board-approved calculators may be used
- Use the Data Sheet and Periodic Table provided
- Use the multiple-choice answer sheet provided
- Write your NESA Number at the top of this page and on the multiple-choice answer sheet

| SECTION | TOTALS | MARKS |
|---------|--------|-------|
| Part A  | 20     |       |
| Part B  | 80     |       |
| TOTAL   | 100    |       |

### Total marks – 100

#### 100 marks

This section has two parts, Part A and Part B

#### Part A – 20 marks

- Attempt Questions 1-20
- Allow about 35 minutes for this part

#### Part B – 80 marks

- Attempt Questions 21-35
- Allow about 2 hours and 25 minutes for this part

### Disclaimer

Every effort has been made to prepare this Examination in accordance with the Board of Studies documents. No guarantee or warranty is made or implied that the Examination paper mirrors in every respect the actual HSC Examination question paper in this course. This paper does not constitute 'advice' nor can it be construed as an authoritative interpretation of Board of Studies intentions. No liability for any reliance, use or purpose related to this paper is taken. Advice on HSC examination issues is only to be obtained from the NSW Board of Studies.

Use the multiple-choice answer sheet.

Select the alternative A,B,C or D that best answers the question. Fill in the response oval completely.

Sample:  $2 + 4 =$  (A) 2 (B) 6 (C) 8 (D) 9

A ☐ B ☒ C ☐ D ☐

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

A ☒ B ☒ C ☐ D ☐

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word **correct** and drawing an arrow as follows.

A ☒ B ☒ C ☐ D ☐  
correct

## Part A

20 marks

Attempt Questions 1–20

Allow about 35 minutes for this part

Use the multiple-choice answer sheet for Questions 1-20

- 1 Which pair of compounds represents an acid and its conjugate base?

|    | <i>Acid</i>                      | <i>Conjugate base</i>                        |
|----|----------------------------------|----------------------------------------------|
| A. | H <sub>2</sub> SO <sub>4</sub>   | SO <sub>4</sub> <sup>2-</sup>                |
| B. | CH <sub>3</sub> COO <sup>-</sup> | CH <sub>3</sub> COOH                         |
| C. | CH <sub>3</sub> NH <sub>2</sub>  | CH <sub>3</sub> NH <sub>3</sub> <sup>+</sup> |
| D. | HCrO <sub>4</sub> <sup>-</sup>   | CrO <sub>4</sub> <sup>2-</sup>               |

- 2 Which of the following most accurately describes how Aboriginal and Torres Strait Islander peoples use solubility principles in the preparation of food sources, such as the fruits of cycads?

- A. The starch in the fruits of cycads is more soluble than the toxins found in these fruits.
- B. The toxins in the ground-up fruits are soluble in running water.
- C. Placing the cycad fruits in water removes hard to digest food products such as fibre.
- D. The skins of the cycad fruits easily dissolve in water leaving essential nutrients.

- 3 Ammonia can be produced by the reaction of hydrogen and nitrogen. When this reaction takes place in a sealed container of fixed volume, an equilibrium system is established.

The equation for the reaction is:




If the pressure and volume remain constant when the temperature is increased, the forward reaction rate will


- A. increase and the [NH<sub>3</sub>] will decrease.
- B. increase and the [NH<sub>3</sub>] will increase.
- C. decrease and the [NH<sub>3</sub>] will decrease.
- D. decrease and the [NH<sub>3</sub>] will remain the same.

4 The structural formula representing 2-methyl-3-heptyne is:


A.




B.



C.



D.



5 Which alternative most correctly represents the chemical reaction for the equilibrium expression shown below?

$$\frac{[\text{Ni}(\text{CO})_4]}{[\text{CO}]^4}$$

- A.  $\text{Ni}(\text{CO})_4(\text{g}) \rightleftharpoons 4\text{CO}(\text{g})$
- B.  $\text{Ni}(\text{CO})_4(\text{g}) \rightleftharpoons \text{Ni}(\text{g}) + 4\text{CO}(\text{g})$
- C.  $\text{Ni}(\text{s}) + 4\text{CO}(\text{g}) \rightleftharpoons \text{Ni}(\text{CO})_4(\text{g})$
- D.  $4\text{Ni}(\text{s}) + 4\text{CO}(\text{g}) \rightleftharpoons 4\text{Ni}(\text{CO})(\text{g})$

6 An isomer of butanoic acid is

- A.  $\text{CH}_3\text{CH}_2\text{COOH}$
- B.  $\text{CH}_3\text{CH}_2\text{COOCH}_3$
- C.  $\text{CH}_3\text{COCH}_2\text{CH}_3$
- D.  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$

7 The following table shows the colour changes and pH ranges of three indicators:

| <i>Indicator</i> | <i>Colour change<br/>(low pH to high pH)</i> | <i>pH range</i> |
|------------------|----------------------------------------------|-----------------|
| bromophenol blue | yellow to blue                               | 3.0–4.5         |
| methyl red       | red to yellow                                | 4.5–6.3         |
| alizarin         | yellow to red                                | 10.2–12.0       |

The indicators were used to test a liquid. The following table shows the final colours of the liquid:

| <i>Indicator</i> | <i>Final colour</i> |
|------------------|---------------------|
| bromophenol blue | blue                |
| methyl red       | yellow              |
| alizarin         | yellow              |

- A. vinegar (pH 2.1)
- B. rain water (pH 5.2)
- C. distilled water (pH 7.0)
- D. bleach (pH 12.1)


8 What is the concentration of barium ions in a saturated solution of barium carbonate?

- A.  $5.08 \times 10^{-5} \text{ mol L}^{-1}$
- B.  $1.60 \times 10^{-3} \text{ mol L}^{-1}$
- C.  $1.16 \times 10^{-3} \text{ mol L}^{-1}$
- D.  $0.005 \text{ mol L}^{-1}$

9 Which of the following organic compounds is classified as an organic base?

- A. Propane
- B. Propanol
- C. Propanamide
- D. Propanamine

10 Consider the reaction




What is the  $K_c$  for the reaction shown below, at  $150^\circ\text{C}$ ?



- A.  $1.6 \times 10^{-2}$
- B.  $4.1 \times 10^{-3}$
- C.  $6.9 \times 10^{-5}$
- D.  $8.03 \times 10^{-5}$

11 The diagram below shows the mass spectrum for an organic compound.



The compound is


- A. ethanoic acid
- B. ethanol
- C. propanol
- D. propanoic acid

- 12 The pH of a  $0.01 \text{ mol L}^{-1}$  solution of a monoprotic acid was measured by a student and found to be 4.

What proportion of the acid molecules has been converted to ions?

- A. 0%
- B. 1%
- C. 90%
- D. 100%


- 13 Which graph best represents the electrical conductivity changes that occur when an aqueous solution of sulfuric acid is titrated with an aqueous solution of barium hydroxide?




- 14 During a precipitation titration investigation, a  $10.0 \text{ mL}$  sample of a solution containing chloride ions was titrated with  $23.5 \text{ mL}$  of  $0.125 \text{ mol L}^{-1} \text{ AgNO}_3$  to reach the end-point. Calculate the mass of chloride ions present in the original sample.

- A.  $0.00294 \text{ g}$
- B.  $0.0520 \text{ g}$
- C.  $0.104 \text{ g}$
- D.  $0.208 \text{ g}$

- 15 The gas carbon monoxide reacts with fluorine gas to form the gas carbonyl difluoride,  $\text{COF}_2$ . The reaction is exothermic.



Scientists studying this reaction measured the concentration of each of these gases in a sealed 2.0 L reaction vessel over a period of time. The results are shown below.




What changes were imposed on this equilibrium at 15 minutes and 30 minutes?

|    | <i>Change at 15 min</i>                     | <i>Change at 30 min</i>                         |
|----|---------------------------------------------|-------------------------------------------------|
| A. | Fluorine gas was removed from the vessel    | The temperature of the vessel was decreased     |
| B. | Fluorine gas was removed from the vessel    | The temperature of the vessel was increased     |
| C. | The temperature of the vessel was decreased | Carbonyl difluoride gas was added to the vessel |
| D. | The temperature of the vessel was increased | Carbonyl difluoride gas was added to the vessel |

- 16 A sample of Teflon, polytetrafluorethene, was found to have a molecular mass of  $1.9 \times 10^4 \text{ g mol}^{-1}$ .

On average, how many monomer units are there in this polymer?

- A. 190
- B. 186
- C. 258
- D. 212



- I When 2 drops of NaOH solution was added to a sample, a white precipitate was formed.
- II After excess NaOH solution was added the precipitate disappeared.
- III When 2 drops, then excess  $\text{NH}_3$  solution was added, a white precipitate appeared.
- IV After dilute  $\text{H}_2\text{SO}_4$  solution was added, the result was a colourless solution.

The ion present in the solution is:

- A.  $\text{Pb}^{2+}$
- B.  $\text{Al}^{3+}$
- C.  $\text{Mg}^{2+}$
- D.  $\text{Zn}^{2+}$

**18** A student wishes to distinguish between samples of pentan-1-ol and pentanoic acid. Which of the following tests would NOT allow the 2 compounds to be identified?

- A. Adding bromine water to both
- B. Adding an acidified solution of potassium permanganate to both
- C. Testing the pH of solutions of both
- D. Adding sodium carbonate solution to both

**19** An organic compound of formula  $C_4H_8O$

I – could be an alkanal or an alkanone

II – could be formed by oxidation of an alkanol

III – could be formed by reduction of an alkanoic acid

IV – could react with an alkanoic acid to form an ester

Which of the above statements are correct?

- A. I, and IV only
- B. I, II and III
- C. I and III only
- D. III and IV only

**20** Three chemical formulas are shown below.

Compound A:  $CH_3(CH_2)_{15}COONa$

Compound B:  $CH_3(CH_2)_5CONH_2$

Compound C:  $CH_3(CH_2)_3COCH_3$

What are the names of the chemical families to which these compounds belong?

|    | <i>Compound A</i> | <i>Compound B</i> | <i>Compound C</i> |
|----|-------------------|-------------------|-------------------|
| A. | carboxylic acid   | amide             | aldehyde          |
| B. | soap              | amine             | aldehyde          |
| C. | ester             | amine             | ketone            |
| D. | soap              | amide             | ketone            |

## Part B

80 marks

Attempt Questions 21-35

Allow about 2 hours and 25 minutes for this section

Answer the questions in the spaces provided. These spaces provide guidance for the expected length of response.


Show all relevant working in questions involving calculations.

Extra writing space is provided on pages 29 to 32. If you use this space, clearly indicate which question you are answering.


### Question 21 (6 marks)

Nitrogen dioxide is brown and dinitrogen tetroxide is colourless. They form an equilibrium mixture as shown by the following equation:

6



A sealed gas syringe can be used to investigate the properties of a fixed mass of gas. An equimolar mixture of nitrogen oxide and dinitrogen tetroxide was set up as shown in X in the following diagram. The conditions were then varied as shown in Y and Z.



Complete the table by describing the colour of the gas mixtures in X, Y and Z. Include any comparisons to the initial colour of X and justify your answers.

|          | <i>Colour</i> | <i>Justification</i> |
|----------|---------------|----------------------|
| <i>X</i> |               |                      |
| <i>Y</i> |               |                      |
| <i>Z</i> |               |                      |

**Question 22 (5 marks)**

A student makes up a 200.0 mL solution of barium hydroxide. The final solution is found to have a pH of 12. The student then adds 0.20 mol L<sup>-1</sup> hydrochloric acid solution to neutralise the barium hydroxide solution.

- (a) Write the balanced equation for the neutralisation reaction. **1**

.....

- (b) Calculate the concentration (in mol L<sup>-1</sup>) of hydroxide ions in the original solution. **2**

.....

.....

- (c) Calculate the volume of hydrochloric acid required to complete the neutralisation reaction. **2**

.....


.....

.....

.....

**Question 23** (6 marks)

The reaction to produce hydrogen from carbon monoxide and steam is shown.



Use the data supplied in the table to answer the following questions.

|                            | $\Delta H_f^\circ$<br>$\text{kJ mol}^{-1}$ | $S^\circ$<br>$\text{J K}^{-1} \text{mol}^{-1}$ |
|----------------------------|--------------------------------------------|------------------------------------------------|
| $\text{CO}_{(g)}$          | -111                                       | 198                                            |
| $\text{H}_2\text{O}_{(g)}$ | -242                                       | 189                                            |
| $\text{CO}_{2(g)}$         | -393                                       | 214                                            |
| $\text{H}_{2(g)}$          | 0                                          | 131                                            |

- (a) Calculate the enthalpy change for this reaction. 1

---

---

---

- (b) Calculate the entropy change for this reaction. 1

---

---

---

- (c) Determine whether this reaction is spontaneous at 600K under standard conditions. 2

---

---

---

---

- (d) Given that when a reaction has reached equilibrium  $\Delta G = 0$ , calculate the minimum temperature required for this reaction to reach equilibrium under standard conditions AND state when the reaction will be spontaneous. 2

---

---

---

---

---

---

**Question 24 (8 marks)**

A solution of hydrochloric acid was standardised by titration against a sodium carbonate solution using the following procedure.

- All glassware was rinsed correctly to remove possible contaminants.
- Hydrochloric acid was placed in the burette.
- 25.0 mL of sodium carbonate solution was pipetted into the conical flask.

The titration was performed and the hydrochloric acid was found to be  $0.200 \text{ mol L}^{-1}$ .

- (a) Identify the substance used to rinse the conical flask and justify your answer. **2**

---

---

---

---

- (b) Explain why sodium carbonate solution, rather than sodium hydroxide solution, is used to standardise the hydrochloric acid solution. **2**

---

---

---

**Question 24 continues on the next page**

Question 24 (continued)

- (c) Seashells contain a mixture of carbonate compounds, including  $\text{CaCO}_3$ .

4

The standardised hydrochloric acid was used to determine the percentage by mass of carbonate ions in a seashell using the following procedure.

- A 0.145 g sample of the seashell was placed in a conical flask.
- 50.0 mL of the standardised hydrochloric acid was added to the conical flask.
- At the completion of the reaction, the mixture in the conical flask was titrated with  $0.250 \text{ mol L}^{-1}$  sodium hydroxide.

The volume of sodium hydroxide used in the titration was 29.5 mL.

Calculate the percentage by mass of carbonate ions in the sample of the seashell.

---

---

---

---

---

---

---

---

---


---

**End of Question 24**


**Question 25** (7 marks)

(a) Using IUPAC nomenclature, name the compounds shown below.


**3**



Compound 1



Compound 2



Compound 3

Compound 1:

Compound 2:

Compound 3:

(b) Compare the intermolecular forces in the above 3 molecules and predict the order of boiling points (lowest to highest) of these molecules. Explain your prediction.

**4**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

### Question 26 (4 marks)

Compare Arrhenius' theory of acids and bases with the Brønsted-Lowry theory.

4

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

**Question 27 (4 marks)**

50.0 mL of a solution of  $0.300 \text{ mol L}^{-1}$  NaOH was added to a 50.0 mL solution of  $0.250 \text{ mol L}^{-1}$  HNO<sub>3</sub> in a styrofoam cup.

The initial temperature of both solutions was  $21.0^{\circ}\text{C}$ . The final temperature of the resulting solution was  $22.5^{\circ}\text{C}$ .

- (a) Calculate the experimental heat of neutralisation from these results. **3**

---

---

---

---

---

---

- (b) Suggest ONE change to the experimental design used above that would improve the accuracy of the investigation. **1**

---

---

**Question 28** (4 marks)

Hydrofluoric acid (HF) ionises in water to give a dissociation constant ( $K_a$ ) value of  $6.9 \times 10^{-4}$  at  $25^\circ\text{C}$ .

- (a) Write an equation for the ionisation of hydrofluoric acid in water. 1

.....

- (b) Calculate the  $pK_a$  value for the ionisation of hydrofluoric acid. 1

.....

.....

- (c) Dissociation  $pK_a$  values at  $25^\circ\text{C}$  for some common acids are listed below. 2

| <i>Acid</i>                                 | <i>pK<sub>a</sub></i> |
|---------------------------------------------|-----------------------|
| ammonium ion ( $\text{NH}_4^+$ )            | 9.25                  |
| carbonic acid ( $\text{H}_2\text{CO}_3$ )   | 6.35                  |
| ethanoic acid ( $\text{CH}_3\text{COOH}$ )  | 4.75                  |
| phosphoric acid ( $\text{H}_3\text{PO}_4$ ) | 2.15                  |
| sulphurous acid ( $\text{H}_2\text{SO}_3$ ) | 1.82                  |

Discuss the relative strength of hydrofluoric acid.

.....

.....

.....

.....

.....

**Question 29 (3 marks)**

Quantitatively compare the solubility of lead(II) sulfate in water with its solubility in a 0.10 mol L<sup>-1</sup> solution of sodium sulfate, 25°C.

**3**

---

---

---

---

---


---

---

**Question 30 (4 marks)**

An equimolar solution of carbonic acid and hydrogen carbonate ions has reached equilibrium with a pH of 6.4.

4



Explain how the addition of both a small volume of dilute acid and base can result in no change in the overall pH of this solution.

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

**Question 31** (4 marks)

Explain how the surfactant properties of the sodium salts of long chain fatty acids help to clean grease from dirty dishes. Use a diagram to support your answer.

**4**

---

---

---

---

---

---

---

---

**Question 32 (4 marks)**

In an addition reaction, 3-hexene is reacted with water and dilute sulfuric acid under heated conditions.

(a) Name the product formed and draw its structural formula. **2**

(b) Name and draw the structural formula for all possible products of the reaction between hydrogen chloride and propene. **2**

.....

.....

.....

.....

.....

.....

**Question 33 (8 marks)**

Yeast was added to a 150 mL solution of glucose and water in a conical flask and the apparatus weighed.

- (a) Describe the optimal conditions required for fermentation. **2**

---

---

---

---

- (b) Write a balanced chemical equation for the fermentation of glucose. **1**

---

- (c) The mixture was left under these optimal conditions for 24 hours. When weighed again, the mass of the mixture had reduced by 2.38 g. **2**

What was the mass of ethanol produced?

---

---

---

---

---

- (d) Discuss the benefits and limitations of using ethanol obtained by the fermentation of sugars as an alternative liquid fuel. **3**

---

---

---

---

---


**Question 34** (8 marks)

Brass is an alloy of copper and zinc.

To determine the percentage of copper in a particular sample of brass, an analyst prepared a number of standard solutions of copper(II) ions and measured their absorbance using an atomic absorption spectrometer (AAS). The results are given in the table.

| <i>Cu<sup>2+</sup> concentration<br/>(mg L<sup>-1</sup>)</i> | <i>Absorbance</i> |
|--------------------------------------------------------------|-------------------|
| 0                                                            | 0                 |
| 50.00                                                        | 0.060             |
| 100.0                                                        | 0.120             |
| 200.0                                                        | 0.240             |
| 300.0                                                        | 0.360             |
| 400.0                                                        | 0.480             |
| 500.0                                                        | 0.600             |

- (a) Draw and label the absorbance versus concentration calibration curve for Cu<sup>2+</sup>.

**3**

**Question 34 continues on the next page**

Question 34 (continued)

A 19.8 mg sample of the brass was dissolved in acid, and the solution was made up to 100 mL in a volumetric flask. The absorbance of this test solution was found to be 0.150.

- (b) Calculate the percentage by mass of copper in the brass sample. **3**

---

---

---

---

---

---

- (c) When using AAS techniques, the presence of  $\text{Zn}^{2+}$  in the sample does not affect the measurement of  $\text{Cu}^{2+}$  in the sample. **2**

Explain this observation.

---

---

---

---

---

---

**End of Question 34**

**Question 35 (5 marks)**

To determine the sulfate content of a sample of fertiliser by gravimetric analysis requires the sample to be dissolved and the sulfate ions precipitated with an appropriate cation before being filtered, dried and then weighed.

Several assumptions have to be made at particular steps in the process.

- (a) Identify any *two* assumptions made in this process and outline how these assumptions may affect the final result. **2**

---

---

---

---

---

- (b) From an initial sample of fertiliser weighing 4.27 g, a dried precipitate of  $\text{BaSO}_4(s)$  weighed 3.12 g. **3**

Calculate the percentage sulfate content of this fertiliser sample.

---

---

---

---

---

---

**End of Paper**

**BLANK PAGE**

**If you use this space, clearly indicate which question you are answering**

[illegible]

Blank lined paper for writing.

**Extra writing space**  
**If you use this space, clearly indicate which question you are answering**

Area with horizontal dotted lines for extra writing space.

Blank lined area for writing.

# 2020 Higher School Certificate Chemistry Trial Examination Marking Criteria

## Part A

| Question | Answer |
|----------|--------|
| 1        | D      |
| 2        | B      |
| 3        | A      |
| 4        | D      |
| 5        | C      |
| 6        | B      |
| 7        | C      |
| 8        | A      |
| 9        | D      |
| 10       | C      |
| 11       | D      |
| 12       | B      |
| 13       | B      |
| 14       | C      |
| 15       | A      |
| 16       | A      |
| 17       | B      |
| 18       | A      |
| 19       | B      |
| 20       | D      |

**Part B – 80 marks****Question 21 (6 marks)**

| Criteria                                         | Marks |
|--------------------------------------------------|-------|
| • Correctly completes all SIX cells of the table | 6     |
| • Correctly completes FIVE cells of the table    | 5     |
| • Correctly completes FOUR cells of the table    | 4     |
| • Correctly completes THREE cells of the table   | 3     |
| • Correctly completes TWO cells of the table     | 2     |
| • Correctly completes ONE cell of the table      | 1     |

**Sample answer**


|          | <i>Colour</i>                                                       | <i>Justification</i>                                                                                                                                                           |
|----------|---------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>X</i> | light brown                                                         | A mixture of colourless $\text{N}_2\text{O}_4$ and brown $\text{NO}_2$ gives a light brown equilibrium mixture.                                                                |
| <i>Y</i> | lighter brown (lighter than <i>X</i> )<br>NOT Completely Colourless | The system has shifted to the right (fewer gas molecules), decreasing the amount of brown $\text{NO}_2$ in the resulting equilibrium mixture (Le Châtelier's principle).       |
| <i>Z</i> | brown (darker than <i>X</i> )                                       | The forward reaction is exothermic. Increasing temperature shifts the reaction to the left, increasing the amount of brown $\text{NO}_2$ in the resulting equilibrium mixture. |

**Marker comments**

In equilibrium, both reactants and products exist so a shift to  $\text{N}_2\text{O}_4$  does not make the system colourless, only lighter brown.

**Question 22 (5 marks)****22 (a) (1 mark)**

| Criteria                                                                               | Marks |
|----------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Writes a correctly balanced equation</li> </ul> | 1     |

**Sample answer****22 (b) (2 marks)**

| Criteria                                                                                                                                                     | Marks |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Correctly calculates the pOH of the solution</li> <li>Correctly calculates the hydroxide ion concentration</li> </ul> | 2     |
| <ul style="list-style-type: none"> <li>Correctly calculates the pOH of the solution</li> </ul>                                                               | 1     |

**Sample answer**

$$\text{pOH} + \text{pH} = 14$$

$$\therefore \text{pOH} = 14 - 12 = 2$$

$$\therefore [\text{OH}^-] = 10^{-2} = 0.01 \text{ mol L}^{-1}$$

**22 (c) (2 marks)**

| Criteria                                                                                           | Marks |
|----------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Correctly calculates the volume of acid required</li> </ul> | 2     |
| <ul style="list-style-type: none"> <li>Substitutes values into the correct formula</li> </ul>      | 1     |

**Sample answer**

$$C_1V_1 = C_2V_2$$

$$0.01 \times 0.2 = 0.2 \times V_2$$

$$\therefore V_2 = 0.01 \text{ L (10.0 mL of acid)}$$

**Marker comments**

$\text{Ba(OH)}_2$  produces 2 moles of  $\text{OH}^-$  ions per mole.

**Question 23 (6 marks)****23 (a) (1 mark)**

| Criteria                                   | Marks |
|--------------------------------------------|-------|
| • Calculates the enthalpy change correctly | 1     |

**Sample answer**

$$\Delta H = (-393 + 0) - (-242 + -111) = -393 + 353$$

$$\Delta H = -40 \text{ kJ mol}^{-1}$$

**23 (b) (1 mark)**

| Criteria                                  | Marks |
|-------------------------------------------|-------|
| • Calculates the entropy change correctly | 1     |

**Sample answer**

$$\Delta S = (214 + 131) - (198 + 189)$$

$$\Delta S = -42 \text{ J K}^{-1} \text{ mol}^{-1}$$

**23 (c) (2 marks)**

| Criteria                                                                                             | Marks |
|------------------------------------------------------------------------------------------------------|-------|
| • Calculates $\Delta G$ correctly and states that the reaction is spontaneous at 600 K with evidence | 2     |
| • Calculates $\Delta G$ correctly                                                                    | 1     |

**Sample answer**

$$\Delta G = \Delta H - T\Delta S$$

$$= -40 - (600 \times -0.042)$$

$$= -14.8 \text{ kJ mol}^{-1}$$

Yes, the reaction is spontaneous at 600 K as  $\Delta G < 0$

**23 (d) (2 marks)**

| Criteria                                                                         | Marks |
|----------------------------------------------------------------------------------|-------|
| • Correctly calculates the minimum temperature for reaction to reach equilibrium | 2     |
| • States a criterion for the reaction to be spontaneous                          |       |
| • Correctly calculates the minimum temperature for reaction to reach equilibrium | 1     |
| OR                                                                               |       |
| • States a criterion for the reaction to be spontaneous                          |       |

**Sample answer**

$$\text{When } \Delta G = 0$$

$$T = \Delta H / \Delta S$$

$$T = -40 / -0.042$$

$$= 952.38 \text{ K}$$

The reaction will be spontaneous when the reaction temperature is below 952.38 K (when Gibbs Free energy change is negative)

### **Marker comments**

Entropy change is in Joules so must be converted to kJ so must be changed to 0.042 kJ in part “c”. Since  $\Delta G = \Delta H - T\Delta S$  a higher temperature will make  $\Delta G$  less.

### **Question 24 (8 marks)**

#### **24 (a) (2 marks)**

| Criteria                                                                                                  | Marks |
|-----------------------------------------------------------------------------------------------------------|-------|
| • Identifies deionised water as the substance used to rinse the conical flask<br>AND justifies the answer | 2     |
| • Identifies deionised water as the substance used to rinse the conical flask                             | 1     |

### **Sample answer**

Standard sodium carbonate is the solution to be placed in the flask. The flask must be cleaned and then rinsed with deionised water. If water remains in the flask, it will not change the number of moles of sodium carbonate in the flask. If either hydrochloric acid or sodium carbonate were used, the moles of the wash remaining in the flask would alter the titration results and hence make the calculation invalid.

#### **24 (b) (2 marks)**

| Criteria                                                                                                             | Marks |
|----------------------------------------------------------------------------------------------------------------------|-------|
| • Explains, giving at least 2 valid reasons, why sodium carbonate is used rather than sodium hydroxide as a standard | 2     |
| • Outlines one valid reason                                                                                          | 1     |

### **Sample answer**

A primary standard solution is one that can be used as the starting point in a series of titrations to determine the concentration of other solutions. The primary standard used to make the primary standard solution is normally a crystalline solid, which can be weighed out accurately (will not absorb water or carbon dioxide from the atmosphere) and has a relatively high molar mass (so that errors in transferring the solid are minimised in terms of moles). Sodium carbonate is suitable as it can be obtained pure, as crystals, can be weighed out accurately without absorbing water or reacting with carbon dioxide from the air and has a higher molar mass (106 g/mol) than NaOH (40 g/mol). Sodium hydroxide is not obtainable as pure crystals (lumps, which absorb water and react with  $\text{CO}_2$ ).


**24 (c) (4 marks)**

| Criteria                                                                        | Marks |
|---------------------------------------------------------------------------------|-------|
| • Calculates correctly the percentage by mass of carbonate ions in the seashell | 4     |
| • Determines the mass of carbonate ions in the seashell                         | 3     |
| • Determines the moles of HCl in excess                                         | 2     |
| • Determines the moles of HCl added to the flask                                | 1     |

**Sample answer**

There were 4 steps in the process:

- The sample (0.145 g) was placed in a conical flask.
- 50.0 mL of standard hydrochloric acid (0.200 mol/L) was pipetted into the flask
- The excess HCl was titrated with NaOH to determine the no. of moles of HCl in excess and hence the no. of moles and mass of carbonate ions
- The percentage of carbonate ions, by mass, was determined



Moles HCl added initially to react and dissolve carbonate ions  
 $= cV = 0.200 \times 0.050 = 0.0100 \text{ mol}$

Moles NaOH required for titration excess HCl  $= 0.0295 \times 0.250 = 0.007375 \text{ mol}$

Hence moles of excess HCl  $= 0.007375$  (as NaOH and HCl react in 1:1 ratio)

Moles of HCl which had reacted with seashell  $= 0.0100 - 0.00738 = 0.00262 \text{ mol}$

Moles  $\text{CO}_3^{2-} = \frac{1}{2}$  moles HCl (from balanced equation)

Hence moles  $\text{CO}_3^{2-}$  in seashell  $= 0.00131 \text{ mol}$

Mass of  $\text{CO}_3^{2-}$  in seashell  $= n \times M = 0.00131 \times 60.01 = 0.0786 \text{ g (3 s.f.)}$

%  $\text{CO}_3^{2-}$  in seashell  $= 0.0786/0.145 \times 100 = 54.2\%$

**Marker comments**

This final answer required the correct significant figures. A lot of students used the molar mass of calcium carbonate rather than carbonate ( $\text{CO}_3^{2-}$ ).

**Question 25 (7 marks)****25 (a) (3 marks)**

| Criteria                          | Marks |
|-----------------------------------|-------|
| • Names THREE compounds correctly | 3     |
| • Names TWO compounds correctly   | 2     |
| • Names ONE compound correctly    | 1     |

**Sample answer**

Compound 1 = propan-1-amine

Compound 2 = 2,2-dimethylpropane

Compound 3 = propanamide

**25 (b) (4 marks)**

| Criteria                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Marks |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>• Predicts the correct order of boiling points</li> <li>• Explains thoroughly the impact of the different intermolecular forces</li> <li>• Identifies that Compound 1 has hydrogen bonding as the strongest intermolecular force</li> <li>• Identifies that Compound 2 has dispersion (temporary dipole-dipole forces) only</li> <li>• Identifies that Compound 3 has hydrogen bonding (2 hydrogen bonds form between adjacent molecules) and forms dimers</li> </ul> | 4     |
| <ul style="list-style-type: none"> <li>• Predicts the correct order of boiling points</li> <li>• Explains thoroughly the impact of the different intermolecular forces</li> <li>• Identifies the intermolecular forces in 2 of the 3 compounds</li> </ul>                                                                                                                                                                                                                                                    | 3     |
| <ul style="list-style-type: none"> <li>• TWO of:               <ul style="list-style-type: none"> <li>• Predicts the correct order of boiling points</li> <li>• Explains thoroughly the impact of the different intermolecular forces</li> <li>• Identifies the intermolecular forces in 2 of the 3 compounds</li> </ul> </li> </ul>                                                                                                                                                                         | 2     |
| <ul style="list-style-type: none"> <li>• ONE of:               <ul style="list-style-type: none"> <li>• Predicts the correct order of boiling points</li> <li>• Explains thoroughly the impact of the different intermolecular forces</li> <li>• Identifies the intermolecular forces in 1 of the 3 compounds</li> </ul> </li> </ul>                                                                                                                                                                         | 1     |

**Sample answer**


The order of increasing boiling points is Compound 2, Compound 1, Compound 3.

The stronger the intermolecular forces, the higher the boiling point, as greater energy is needed to separate the liquid molecules to form a gas.


Compound 2 is non-polar and has only weak intermolecular forces (dispersion or temporary dipole-dipole forces) caused by the electrical interaction of molecules as they collide (protons from 1 molecule being attracted to electrons from the other as the molecules are temporarily distorted on collision).

Compound 1 is polar and would experience hydrogen bonding forces, as well as weaker temporary and permanent dipolar forces) as molecules interact. These are strong intermolecular forces where the electronegativity of the nitrogen results in a very polar bond

with hydrogen in the N–H group. This hydrogen is attracted to the nitrogen of a neighbouring amine molecule. The geometry of the molecule only allows 1 H-bond per pair of molecules at any instant.



Compound 3 is an amide and has the very polar – CONH functional group. The hydrogen atom of the –CONH can form a hydrogen bond with an oxygen of the neighbouring amide molecule. The planar nature of this –CONH group allows 2 H-bonds per pair of molecules. Hence the intermolecular forces and thus boiling points are highest in Compound 3 and lowest in Compound 2.



### **Marker comments**

Boiling point depends on inter-molecular forces such as dispersion forces (weakest), dipole-dipole interaction and Hydrogen bonding (strongest). No covalent bonds are broken when a molecule changes state only the inter-molecular forces.

**Question 26 (4 marks)**


| Criteria                                                                                                                                                                                                                         | Marks |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"><li>• Explains in detail similarities and differences between Arrhenius definition of acids and bases and Brønsted- Lowry</li><li>• Supports answer with clear, well-chosen examples</li></ul> | 4     |
| <ul style="list-style-type: none"><li>• Explains similarities and differences between Arrhenius definition of acids and bases and Brønsted- Lowry</li><li>• Supports answer with examples</li></ul>                              | 3     |
| <ul style="list-style-type: none"><li>• Explains at least one similarity and one difference between Arrhenius definition of acids and bases and Brønsted- Lowry</li></ul>                                                        | 2     |
| <ul style="list-style-type: none"><li>• Gives either a similarities OR differences between Arrhenius definition of acids and bases and Brønsted- Lowry</li></ul>                                                                 | 1     |

**Sample answer**

Svante Arrhenius theorised that acids were compounds that produced hydrogen ions in water. Strong acids ionised completely while weak acids only ionised slightly. He said bases were substances which produced hydroxide ions in solution. This definition failed to take into account the basic nature of metal oxides and ammonia. The concept of acid was further developed by two chemists, Johannes Brønsted and Thomas Lowry, who said that acids are chemical species that transferred a proton to another species. This included the same acid compounds as the Arrhenius definition. The Brønsted- Lowry definition of bases as species which accept protons included the basic compounds excluded by the Arrhenius definition.

**Marker comments**

It is always better to use equations to illustrate your answer. E.g. Arrhenius



**Question 27 (4 marks)****27 (a) (3 marks)**

| Criteria                                                                                                                                                                                                                          | Marks |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Heat energy produced in reaction is calculated</li> <li>Number of moles H<sub>2</sub>O produced is calculated</li> <li>Heat of neutralisation subsequently calculated correctly</li> </ul> | 3     |
| <ul style="list-style-type: none"> <li>Two of the above steps correct</li> </ul>                                                                                                                                                  | 2     |
| <ul style="list-style-type: none"> <li>One step correct</li> </ul>                                                                                                                                                                | 1     |

**Sample answer**

$$\begin{aligned}
 q &= mc\Delta T \\
 &= 100 \times 4.18 \times (22.5 - 21.0) \\
 &= 627 \text{ J} \\
 &= 0.627 \text{ kJ}
 \end{aligned}$$

$$\begin{aligned}
 \text{No. moles H}_2\text{O produced} &= c \times v \text{ for HNO}_3 \text{ (limiting reagent)} \\
 &= 0.250 \times 0.0500 \\
 &= 1.25 \times 10^{-2} \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{Heat of neutralisation} &= 0.627 \text{ kJ} / 1.25 \times 10^{-2} \\
 &= 50.2 \text{ kJ}
 \end{aligned}$$

**Marker comments**

This is a limiting reagent question. Only 0.0125 moles of acid and base can react.

**27 (b) (1 mark)**


| Criteria                                                                                             | Marks |
|------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>A change suggested that would improve the accuracy</li> </ul> | 1     |

**Sample answer**

e.g. Use larger quantities; use a calorimeter with closed lid; etc

**Question 28 (4 marks)****28 (a) (1 mark)**

| Criteria                                                        | Marks |
|-----------------------------------------------------------------|-------|
| • Writes a correctly balanced equation for the ionisation of HF | 1     |

**Sample answer****Marker comments**

One-way arrows were not penalised, however, better answers identified that HF was a weak acid and therefore a two-way arrow was appropriate.

**28 (b) (1 mark)**

| Criteria                                               | Marks |
|--------------------------------------------------------|-------|
| • Calculates the $pK_a$ value for the ionisation of HF | 1     |

**Sample answer**

$$pK_a = -\log_{10}(K_a) = -\log_{10}(6.9 \times 10^{-4}) = 3.16$$

**Marker comments**

Some misconception around significant figures (did not affect mark) – similar to pH values, only the decimal places in  $pK_a$  are significant, therefore 3.16 has 2 sig. figs.

**28 (c) (2 marks)**

| Criteria                                                       | Marks |
|----------------------------------------------------------------|-------|
| • Associates a lower $pK_a$ value with stronger acids          | 2     |
| • Compares strength of HF with acids listed in the table       |       |
| • Makes a relevant statement about the relative strength of HF | 1     |

**Sample answer**

The  $pK_a$  value for HF lies between ethanoic acid and phosphoric acid. As the values for  $pK_a$  decrease the strength of the acid increases. Therefore, hydrofluoric acid is a stronger acid than ethanoic acid but is a weaker acid than phosphoric acid.

**Marker comments**

Additional information provided by students (not marked) showed some misconceptions regarding value of  $pK_a$  associated with strong acids. Note that none of the listed acids are strong; all are varying levels of weak (strong acids have negative  $pK_a$  values (typically  $< -2$ )). Also, being polyprotic is irrelevant to strength of an acid, e.g. sulfuric acid is diprotic and strong, whereas citric acid is triprotic and very weak.

**Question 29 (3 marks)**

| Criteria                                                                                                                                                                                                  | Marks |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Correctly calculates solubility in water and in 0.10 M Na<sub>2</sub>SO<sub>4</sub></li> <li>Compares the two solubilities</li> </ul>                              | 3     |
| <ul style="list-style-type: none"> <li>Calculates solubility in water and in 0.10 M Na<sub>2</sub>SO<sub>4</sub> with one error AND</li> <li>Compares the two solubilities</li> </ul>                     | 2     |
| <ul style="list-style-type: none"> <li>Provides some relevant steps to calculating solubility AND/OR</li> <li>Qualitatively explains that solubility will be lower in solution of a common ion</li> </ul> | 1     |

**Sample answer**

$$K_{sp} \text{ PbSO}_4 = [\text{Pb}^{2+}] [\text{SO}_4^{2-}] = 2.53 \times 10^{-8}$$

The solubility of PbSO<sub>4</sub> in water =  $[\text{Pb}^{2+}] = [\text{SO}_4^{2-}] = \sqrt{K_{sp}} = \sqrt{(2.53 \times 10^{-8})}$   
= **1.59 x 10<sup>-4</sup> mol/L**

Let the solubility of PbSO<sub>4</sub> in a 0.10 mol/L solution of sodium sulfate = s

$$K_{sp} \text{ PbSO}_4 = [\text{Pb}^{2+}] [\text{SO}_4^{2-}] = (s) (0.10 + s) = 2.53 \times 10^{-8}$$

Since s is small by comparison with 0.10,

$$K_{sp} \text{ PbSO}_4 = [\text{Pb}^{2+}] [\text{SO}_4^{2-}] = (s) (0.10) = 2.53 \times 10^{-8}$$

$$s = 2.53 \times 10^{-8} / (0.10) = 2.53 \times 10^{-7} \text{ mol/L}$$

The solubility of PbSO<sub>4</sub> in 0.10 M SO<sub>4</sub><sup>2-</sup> = **2.53 x 10<sup>-7</sup> mol/L**

Hence PbSO<sub>4</sub> is much less soluble in a 0.10 mol/L common ion solution of sulfate ion than in water.

**Marker comments**

Many students didn't conduct any calculations and only qualitatively compared solubility. There was also a lot of confusion around the distinction between solubility and  $K_{sp}$ , as many students incorrectly assumed the terms are interchangeable.


**Question 30 (4 marks)**

| Criteria                                                                                                                                                                                                                                                                                                       | Marks |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Identifies the solution H<sub>2</sub>CO<sub>3</sub>/HCO<sub>3</sub><sup>-</sup> as a buffer</li> <li>Explains in detail how the addition of a small amount of acid or base can shift the equilibrium of the system to absorb the changes and maintain the pH</li> </ul> | 4     |
| <ul style="list-style-type: none"> <li>Identifies the solution H<sub>2</sub>CO<sub>3</sub>/HCO<sub>3</sub><sup>-</sup> as a buffer</li> <li>Explains how the addition of a small amount of acid or base can shift the equilibrium of the system to absorb the changes and maintain the pH</li> </ul>           | 3     |
| <ul style="list-style-type: none"> <li>Identifies the solution is a buffer</li> <li>Refers to LCP/equilibrium shift to maintain pH</li> </ul>                                                                                                                                                                  | 2     |
| <ul style="list-style-type: none"> <li>Identifies the solution is a buffer</li> </ul>                                                                                                                                                                                                                          | 1     |


**Sample answer**

An equimolar solution of  $\text{H}_2\text{CO}_3/\text{HCO}_3^-$  is a buffer. This is a solution consisting of a weak acid and its conjugate base.

If a small amount of acid such as  $\text{HCl}$  is added to this buffer then the equilibrium will shift to the left as the increase in hydrogen ions will cause the reaction to shift in such a way as to counteract the change. The additional  $\text{H}^+$  will react with the  $\text{HCO}_3^-$  (aq) to form  $\text{H}_2\text{CO}_3$ (aq) and the pH will remain constant.



If a small amount of base such as  $\text{NaOH}$  is added to this buffer then the equilibrium will shift to the right as the increase in hydroxide will cause the reaction to shift in such a way as to counteract the change. The additional  $\text{OH}^-$  will react with the  $\text{H}_3\text{O}^+$  to form water and the  $\text{H}_2\text{CO}_3$  will produce more  $\text{HCO}_3^-$  (aq) and the pH will remain constant.



### ***Marker comments***

Writing an equation was not required, however reference must have been made to the specific reaction given, and better answers used an equation to describe the neutralisation of added  $\text{OH}^-$ .

**Question 31 (4 marks)**


| Criteria                                                                                                                                                                                                                                                                                                         | Marks |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"><li>Provides a detailed description of the structure of a soap molecule</li><li>Correctly outlines how soap molecules interact with both water and grease</li><li>Refers specifically to the formation of micelles</li><li>Includes an appropriate, labelled diagram</li></ul> | 4     |
| <ul style="list-style-type: none"><li>Most of the above</li></ul>                                                                                                                                                                                                                                                | 3     |
| <ul style="list-style-type: none"><li>Some of the above</li></ul>                                                                                                                                                                                                                                                | 2     |
| <ul style="list-style-type: none"><li>Provides some relevant information</li></ul>                                                                                                                                                                                                                               | 1     |

**Sample answer**


The sodium salts of long chain fatty acids consist of a non-polar hydrophobic 'tail' and a polar, hydrophilic, negatively charged 'head' consisting of the anionic carboxylate group.



A micelle forms when soap molecules assemble so that the long hydrophobic tails all point inwards and the polar heads all sit on the outside of the micelle.



When in contact with grease, the hydrophobic tails interact via dispersion forces, while the anionic heads remain in contact with the water via ion-dipole forces. With agitation, the soap molecules surround particles of grease to lift them off surfaces and trap them within micelles which become suspended in the water. The anionic heads from adjacent micelles repel each other so the grease cannot recombine and are easily rinsed away.

**Marker comments**


Any one diagram was acceptable as long as it was accurate and labelled. Better answers referred to dispersion forces and ion-dipole forces as the forces behind the interaction of the soap molecules with both water and grease.

**Question 32 (4 marks)****32 (a) (2 marks)**

| Criteria                                                                                                                                               | Marks |
|--------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"><li>• Correct name supplied</li><li>• Correct structural formula drawn</li></ul>                                     | 2     |
| <ul style="list-style-type: none"><li>• One of the above</li></ul> OR <ul style="list-style-type: none"><li>• Identifies hexanol is produced</li></ul> | 1     |

**Sample answer**


3-hexanol (or hexan-3-ol)

**32 (b) (2 marks)**


| Criteria                                                                                                                                                                                                              | Marks |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"><li>• Correct name and structural formula given with no incorrect additional products or duplication</li></ul>                                                                      | 2     |
| <ul style="list-style-type: none"><li>• Either correct name OR structural formula provided</li></ul> OR <ul style="list-style-type: none"><li>• Correct answer but with extra incorrect additional products</li></ul> | 1     |

**Sample answer**

1-chloropropane:



2-chloropropane:



**Question 33 (8 marks)****33 (a) (2 marks)**


| Criteria                                          | Marks |
|---------------------------------------------------|-------|
| • Two conditions, including anaerobic, identified | 2     |
| • Any one condition identified                    | 1     |

**Sample answer**

Anaerobic (stop air getting in); warm (around 25–35°C); sugar dissolved in water.

**33 (b) (1 mark)**

| Criteria                                | Marks |
|-----------------------------------------|-------|
| • Appropriate balanced equation written | 1     |

**Sample answer****33 (c) (2 marks)**

| Criteria                                                          | Marks |
|-------------------------------------------------------------------|-------|
| • Correct answer given                                            | 2     |
| • An appropriate calculation performed but error or omission made | 1     |

**Sample answer**

Assume mass loss =  $\text{CO}_2(\text{g})$  liberated

$$n(\text{CO}_2) = 2.38 \text{ g} / 44.01 \text{ g mol}^{-1} = 0.05408 \text{ mol}$$

$$n(\text{CO}_2) = n(\text{C}_2\text{H}_5\text{OH})$$

$$\text{mass}(\text{C}_2\text{H}_5\text{OH}) = 0.05408 \times 46.068 = 2.49 \text{ g}$$

**Marker comments**

Many students didn't attribute mass loss to amount of gas formed and instead assumed it was equal to mass of glucose reacted.

**33 (d) (3 marks)**

| Criteria                                                                                                                          | Marks |
|-----------------------------------------------------------------------------------------------------------------------------------|-------|
| • At least three benefits and limitations (including at least one of each) of the use of ethanol as an alternative fuel discussed | 3     |
| • Response shows evidence of good understanding of the issue                                                                      |       |
| • A benefit and a limitation of the use of ethanol as an alternative fuel discussed                                               | 2     |
| • A benefit or a limitation is described                                                                                          | 1     |

**Sample answer**

Sugars for fermentation to produce ethanol are obtained from plant material or biomass. In many cases, the plant material is sourced from a crop that would otherwise be used as a food (e.g. rice or corn), thus increasing the price of the food, having a detrimental effect for poor populations. The distillation of the fermentation mixture to obtain pure ethanol is energy intensive, however burning waste biomass is often used for this. The biomass grown for


ethanol consumes CO<sub>2</sub> from the atmosphere, recycling that CO<sub>2</sub> released upon the combustion of the ethanol, making it less carbon emitting overall than fossil fuel liquids such as petrol or diesel.

**Question 34 (8 marks)**

**34 (a) (3 marks)**

| Criteria                                                                                                                                                             | Marks |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Plots points correctly</li> <li>Labels axes and uses appropriate scale</li> <li>Draws appropriate line of best fit</li> </ul> | 3     |
| <ul style="list-style-type: none"> <li>One error or omission from the above</li> </ul>                                                                               | 2     |
| <ul style="list-style-type: none"> <li>More than one error or omission from the above</li> </ul>                                                                     | 1     |

**Sample answer**



**34 (b) (3 marks)**

| Criteria                                                                                                                                                                                                                                           | Marks |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <ul style="list-style-type: none"> <li>Accurately reads graph to provide concentration (between 115-135 mg L<sup>-1</sup> is acceptable)</li> <li>Correctly determines mass of Cu in sample</li> <li>Correctly determines %Cu in sample</li> </ul> | 3     |
| <ul style="list-style-type: none"> <li>Determines mass of Cu in the sample based on an incorrect reading of the graph</li> </ul>                                                                                                                   | 2     |
| <ul style="list-style-type: none"> <li>Provides some relevant calculations</li> </ul>                                                                                                                                                              | 1     |

**Sample answer**

From the graph, an absorbance of 0.150 gives a concentration of 120 mg L<sup>-1</sup>.

The brass sample was dissolved in 100 mL; hence, it contains 12.0 mg of Cu(II).

$$\begin{aligned}
 \% \text{ of Cu} &= \text{mass of Cu} / \text{mass of sample} \times 100 \\
 &= 12.0 / 19.8 \times 100 \\
 &= 60.6\%
 \end{aligned}$$

### Marker comments

A number of students struggled with the conversion of mg/L to mg/100 mL and tried using more complex routes, not always successfully.

### 34 (c) (2 marks)

| Criteria                             | Marks |
|--------------------------------------|-------|
| • Provides a detailed explanation    | 2     |
| • Provides some relevant information | 1     |

### Sample answer

The hollow cathode lamp that is used in the atomic absorption spectrometer (AAS) analysis of copper contains a copper cathode that produces wavelengths of light uniquely characteristic for copper. Zinc does not absorb light at the same wavelengths as copper.

### Marker comments

There was some confusion between the different analysis techniques – many students described colourimetry instead of AAS. AAS is more robust than colourimetry, the metal does not need to give a visible colour to be able to be analysed by AAS.

For two marks, students had to state the principle involved (unique absorption spectra for different cations) AND how AAS achieves this (use of a cathode lamp containing the analyte to produce the characteristic spectrum).

### Question 35 (5 marks)

### 35 (a) (2 marks)

| Criteria                                                                                                       | Marks |
|----------------------------------------------------------------------------------------------------------------|-------|
| • Two assumptions identified<br>• For each assumption, how the final result is affected is outlined correctly  | 2     |
| • At least one assumption identified<br>• For one assumption, how the result is affected is outlined correctly | 1     |

### Sample answer

Assumption 1: All the  $\text{SO}_4^{2-}$  ions are precipitated out of the solution. This has the effect of obtaining a result lower than the actual result for the % of  $\text{SO}_4^{2-}$  in the fertiliser.

Assumption 2: All the precipitate is filtered out of the solution. In effect, some precipitate may pass through the filter paper used, so the assumption again has the effect of causing the result to be lower than the actual content of  $\text{SO}_4^{2-}$  in the fertiliser.

### Marker comments

Some students were vague in the effect of the assumption – “affects the accuracy of the result” is not sufficient – would the measured value be *higher or lower* than the true value? NOTE: water fully evaporated as an assumption was marked correct, but students are advised to not use this as an example in future since it can be avoided by ‘drying to constant mass’ (this can be used as an example of a potential *source of error* in the method).

**35 (b) (3 marks)**

| Criteria                                                    | Marks |
|-------------------------------------------------------------|-------|
| • Correct answer calculated                                 | 3     |
| • Answer calculated but with an error made or an omission   | 2     |
| • A correct step or steps made towards solving the question | 1     |

***Sample answer***

$$n \text{ BaSO}_4 = \frac{3.12}{137.3 + 32.06 + 4 \times 16.00}$$
$$= 0.01337 \text{ mol}$$

$$\therefore \text{mass SO}_4^{2-} = 0.01337 \times (32.06 + 4 \times 16.00)$$
$$= 1.284 \text{ g}$$

$$\% \text{ SO}_4^{2-} = (1.284/4.27) \times 100\%$$
$$= 30.1 \% \text{ (3 sig figs)}$$

***Marker comments***

Many students calculated % in precipitate, rather than % in the original sample. Students are reminded to not round off their interim values in multistep calculations. Only round at the end. This question was not marked on sig. figs., but students should note that the final answer should not have more significant figures than the value with least significant figures used in the calculation.

## Part A

| Question | Marks | Content                                                                | Syllabus Outcomes       | Targeted Performance Bands |
|----------|-------|------------------------------------------------------------------------|-------------------------|----------------------------|
| 1        | 1     | Mod 6: Properties of Acids and Bases                                   | CH12-13                 |                            |
| 2        | 1     | Mod 5: Solution Equilibria                                             | CH12-12                 | 3-4                        |
| 3        | 1     | Mod 5: Factors that Affect Equilibrium                                 | CH12-5, CH12-12         | 4-5                        |
| 4        | 1     | Mod 7: Hydrocarbons                                                    | CH12-4, CH12-13         | 3-4                        |
| 5        | 1     | Mod 5: Calculating the Equilibrium Constant                            | CH12-4, CH12-12         |                            |
| 6        | 1     | Mod 7: Nomenclature                                                    | CH12-4, CH12-14         | 3-4                        |
| 7        | 1     | Mod 6: Properties of Acids and Bases                                   | CH12-6, CH12-13         | 4                          |
| 8        | 1     | Mod 5: Solution Equilibria                                             | CH12-6, CH12-12         |                            |
| 9        | 1     | Mod 7: Analysis of Organic Acids and Bases                             | CH12-5, CH12-14         | 3-4                        |
| 10       | 1     | Mod 5: Calculating the Equilibrium Constant                            | CH12-4, CH12-12         | 3-4                        |
| 11       | 1     | Mod 8: Analysis of Organic Substances                                  | CH12-5, CH12-15         | 3-4                        |
| 12       | 1     | Mod 6: Quantitative Analysis                                           | CH12-5, CH12-13         | 3-4                        |
| 13       | 1     | Mod 6: Quantitative Analysis                                           | CH12-5, CH12-13         | 3-4                        |
| 14       | 1     | Mod 8: Analysis of Inorganic Substances                                | CH12-6, CH12-15         | 3-4                        |
| 15       | 1     | Mod 5: Factors that Affect Equilibrium                                 | CH12-5, CH12-6, CH12-12 | 4-5                        |
| 16       | 1     | Mod 7: Polymers                                                        | CH12-6, CH12-14         | 4-5                        |
| 17       | 1     | Mod 8: Analysis of Inorganic Substances                                | CH12-6, CH12-14         | 4-5                        |
| 18       | 1     | Mod 8: Analysis of Organic Substances                                  | CH12-5, CH12-15         | 4-5                        |
| 19       | 1     | Mod 7: Nomenclature, Alcohols and Reactions of Organic Acids and Bases | CH12-6, CH12-14         | 5-6                        |
| 20       | 1     | Mod 7: Nomenclature                                                    | CH12-14                 | 4-6                        |

## Part B

| Question | Marks | Content                                                   | Syllabus Outcomes               | Targeted Performance Bands |
|----------|-------|-----------------------------------------------------------|---------------------------------|----------------------------|
| 21       | 6     | Mod 5: Factors that Affect Equilibrium                    | CH12-5, CH12-6, CH12-13         | 5                          |
| 22 (a)   | 1     | Mod 6: Properties of Acids and Bases                      | CH12-13                         | 2-3                        |
| 22 (b)   | 2     | Mod 6: Using the Brønsted-Lowry Theory                    | CH12-6, CH12-13                 | 3-4                        |
| 22 (c)   | 2     | Mod 6: Using the Brønsted-Lowry Theory                    | CH12-6, CH12-13                 | 4                          |
| 23 (a)   | 1     | Mod 5: Static and Dynamic Equilibrium                     | CH12-12                         |                            |
| 23 (b)   | 1     | Mod 5: Static and Dynamic Equilibrium                     | CH12-12                         |                            |
| 23 (c)   | 2     | Mod 5: Factors that Affect Equilibrium                    | CH12-5, CH12-12                 |                            |
| 23 (d)   | 2     | Mod 5: Calculating the Equilibrium Constant               | CH12-12                         |                            |
| 24 (a)   | 2     | Mod 6: Quantitative Analysis                              | CH12-2, CH12-3, CH12-7, CH12-13 | 3-4                        |
| 24 (b)   | 2     | Mod 6: Quantitative Analysis                              | CH12-2, CH12-3, CH12-7, CH12-13 | 3-4                        |
| 24 (c)   | 4     | Mod 6: Quantitative Analysis                              | CH12-5, CH12-6, CH12-13         | 2-5                        |
| 25 (a)   | 3     | Mod 7: Nomenclature                                       | CH12-4, CH12-7, CH12-14         | 3-5                        |
| 25 (b)   | 4     | Mod 7: Hydrocarbons, Reactions of Organic Acids and Bases | CH12-5, CH12-7, CH12-14         | 3-6                        |
| 26       | 4     | Mod 6: Properties of Acids and Bases                      | CH12-6, CH12-13                 |                            |
| 27 (a)   | 3     | Mod 6: Properties of Acids and Bases                      | CH12-4, CH12-5, CH12-13         | 3-6                        |
| 27 (b)   | 1     | Mod 6: Properties of Acids and Bases                      | CH12-4, CH12-5, CH12-13         | 3-6                        |
| 28 (a)   | 1     | Mod 6: Quantitative Analysis                              | CH12-13                         | 2-3                        |
| 28 (b)   | 1     | Mod 6: Quantitative Analysis                              | CH12-6, CH12-13                 | 3-4                        |
| 28 (c)   | 2     | Mod 6: Quantitative Analysis                              | CH12-7, CH12-13                 | 4-5                        |
| 29       | 3     | Mod 5: Solution Equilibria                                | CH12-4, CH12-6, CH12-12         | 3-6                        |
| 30       | 4     | Mod 6: Quantitative Analysis                              | CH12-2, CH12-13                 |                            |
| 31       | 4     | Mod 7: Reactions of Organic Acids and Bases               | CH12-6, CH12-7, CH12-13         | 4                          |
| 32 (a)   | 2     | Mod 7: Products of reactions Involving Hydrocarbons       | CH12-7, CH12-14                 | 2-5                        |

|        |   |                                                     |                                                 |     |
|--------|---|-----------------------------------------------------|-------------------------------------------------|-----|
| 32 (b) | 2 | Mod 7: Products of reactions Involving Hydrocarbons | CH12-7, CH12-14                                 | 2-5 |
| 33 (a) | 2 | Mod 7: Alcohols                                     | CH12-3, CH12-4, CH12-14                         | 2-6 |
| 33 (b) | 1 | Mod 7: Alcohols                                     | CH12-3, CH12-4, CH12-14                         | 2-6 |
| 33 (c) | 2 | Mod 7: Alcohols                                     | CH12-3, CH12-4, CH12-14                         | 2-6 |
| 33 (d) | 3 | Mod 7: Alcohols                                     | CH12-3, CH12-4, CH12-14                         | 2-6 |
| 34 (a) | 3 | Mod 8: Analysis of Inorganic Substances             | CH12-1, CH12-4, CH12-5, CH12-6, CH12-7, CH12-15 | 3-4 |
| 34 (b) | 3 | Mod 8: Analysis of Inorganic Substances             | CH12-1, CH12-4, CH12-5, CH12-6, CH12-7, CH12-15 | 3   |
| 34 (c) | 2 | Mod 8: Analysis of Inorganic Substances             | CH12-2, CH12-4, CH12-6, CH12-15                 | 4-5 |
| 35 (a) | 2 | Mod 8: Analysis of Inorganic Substances             | CH12-3, CH12-4, CH12-6, CH12-15                 | 2-6 |
| 35 (b) | 3 | Mod 8: Analysis of Inorganic Substances             | CH12-3, CH12-4, CH12-6, CH12-15                 | 2-6 |