



SAINT IGNATIUS' COLLEGE

Trial Higher School Certificate

2005

MATHEMATICS

8:50am – 11:55am
Monday 22nd August 2005

Directions to Students

• Reading Time : 5 minutes	• Total Marks 120
• Working Time : 3 hours	
• Write using blue or black pen. (sketches in pencil).	• Attempt Question 1 – 10
• Board approved calculators may be used	• All questions are of equal value
• A table of standard integrals is provided at the back of this paper.	
• All necessary working should be shown in every question.	
• Answer each question in the booklets provided and clearly label your name and teacher's name.	

This paper has been prepared independently of the Board of Studies NSW to provide additional exam preparation for students. Although references have been reproduced with permission of the Board of Studies NSW, the publication is in no way connected with or endorsed by the Board of Studies NSW.

Total marks (120)
Attempt Questions 1 – 10
All questions are of equal value


Answer each question in a SEPARATE writing booklet.

QUESTION 1 (12 MARKS) Use a SEPARATE writing Booklet Marks

- (a) The circumference of a circle is 9.29 metres. Find the radius of the circle to 3 significant figures, using the formula $C = 2\pi r$. 2
- (b) Differentiate $3x^2 - 4\sqrt{x}$. 2
- (c) Expand and simplify $(\sqrt{6} + \sqrt{2})^2$. 2
- (d) Solve the equation $a - \frac{a-2}{4} = 4$. 2
- (e) Find the values of c for which $|2c + 3| > 1$. 2
- (f) The roots of an equation are $\frac{2}{3}$ and -4 . Write down its equation in the form $ax^2 + bx + c = 0$ where a, b and c are integers. 2

QUESTION 2 (12 MARKS) Use a SEPARATE writing Booklet

Marks



The diagram shows the points $A(-2, 3)$ and $B(4, 1)$.
 M is the midpoint of the interval AB . l is the line through M perpendicular to AB .

- | | |
|---|---|
| (i) Find the gradient of the line AB . | 1 |
| (ii) Find the coordinates of M . | 1 |
| (iii) Show that the equation of AB is $x + 3y - 7 = 0$. | 1 |
| (iv) Show that the equation of l is $3x - y - 1 = 0$. | 1 |
| (v) Find the coordinates of C , the point of intersection of l and the y -axis. | 1 |
| (vi) Show that C is equidistant from A and B . | 2 |
| (vii) Find the perpendicular distance from C to the line AB . | 2 |
| (viii) Hence find the area of the triangle ABC . | 2 |
| (ix) State the inequality that defines the region of the half-plane under the line AB . | 1 |

QUESTION 3 (12 MARKS) Use a SEPARATE writing Booklet

Marks

(a) Differentiate with respect to x :

- | | |
|-----------------------|---|
| (i) $x^3 \cos x$ | 2 |
| (ii) $(3 + e^{2x})^4$ | 2 |

- | | |
|---|---|
| (b) (i) Find $\int \frac{6x}{x^2 + 2} dx$. | 2 |
| (ii) Find $\int_0^{\frac{\pi}{6}} \sec^2 2x dx$. | 2 |

(c) The gradient function of a curve is given by $\frac{dy}{dx} = 6x^2 - 4x + 1$.


The curve passes through the point $(1, 6)$.

- | | |
|---|---|
| (i) Find the equation of the curve. | 2 |
| (ii) What is the equation of the tangent to the curve at the point $(1, 6)$? | 2 |

QUESTION 4 (12 MARKS) Use a SEPARATE writing Booklet **Marks**

- (a) Consider the function $f(x) = x^3 - 6x^2 + 9x$.
- (i) Find the coordinates of the stationary points and determine their nature. 3
 - (ii) Find the x -coordinate of the point of inflexion. 2
 - (iii) Find where the curve meets the x -axis. 1
 - (iv) Sketch the graph of $y = f(x)$. 1
 - (v) Find the values of x for which the gradient is decreasing. 1

(b)



The diagram shows the region bounded by the curve $y^2 = 4x$ and the line $x = 4$.


- (i) Find the exact area of the shaded region. 2
- (ii) The region is rotated about the x -axis. Calculate the exact volume of the solid formed. 2

QUESTION 5 (12 MARKS) Use a SEPARATE writing Booklet **Marks**

- (a) Jim decides to start a daily exercise program, in which he increases the time spent on exercise each day by 2 minutes from the previous day. On the first day he does 10 minutes of exercise.
- (i) For how long will Jim exercise on the 30th day? 1
 - (ii) What will be the total length of time spent on exercise in the first 30 days? 2
 - (iii) On which day would Jim have achieved 40 hours of exercise in total? 3
- (b) Jill commences her working career, earning \$30 000 per year in her first year. If she stays with the same company, her annual salary will increase by 5% each year.
- (i) What will be Jill's annual salary in her 20th year (to nearest dollar)? 2
 - (ii) In what year would her annual salary first exceed \$50 000? 2
 - (iii) What will be her total earnings in the first 20 years of employment? 2

QUESTION 6 (12 MARKS) Use a SEPARATE writing Booklet **Marks**


(a)



The diagram shows the triangular cross-section of the roofline of a modern building.
 $AB = 8\text{m}$, $BC = 13\text{m}$, and $\angle ABC = 65^\circ$. BC is horizontal.

- | | | |
|-------|--|---|
| (i) | Find the length of the beam AC . | 2 |
| (ii) | Find the angle of elevation of the beam AC . | 2 |
| (iii) | Building regulations require a cross-sectional area of at least 48m^2 for adequate ventilation. Does this design satisfy this regulation? | 1 |

(b)




AOB is the sector of a circle, centre O and radius 10m .
 The area of the sector is 60m^2 . Calculate the exact perimeter of the sector AOB .

Question 6 continues on page 7

Question 6 (continued)

(c)



$ABCD$ is a trapezium, and AB is parallel to DC .
 The diagonals AC and BD intersect at X . $AX = 4\text{cm}$, $CX = 10\text{cm}$, $BD = 12\text{cm}$.

Copy the diagram neatly into your answer booklet.

- | | | |
|------|---|---|
| (i) | Prove $\triangle AXB$ is similar to $\triangle CXD$. | 2 |
| (ii) | Find the length of BX . | 2 |

End of Question 6


QUESTION 7 (12 MARKS) Use a SEPARATE writing Booklet **Marks**

- (a) Find the coordinates of the points of intersection of the line $2x - y - 2 = 0$ and the hyperbola $xy = 4$. **3**
- (b) Solve the equation for x :
 $3 \times 3^{2x} + 26 \times 3^x - 9 = 0$. **3**
- (c) For what values of c does the equation $x^2 + 5x + c = 0$ have two real and different roots? **2**
- (d) α, β are the roots of the quadratic equation $2x^2 - 5x - 8 = 0$.
 Without finding the roots, find the value of:
- (i) $\alpha^2 + \beta^2$. **2**
- (ii) $\frac{\alpha}{\beta} + \frac{\beta}{\alpha}$. **2**

QUESTION 8 (12 MARKS) Use a SEPARATE writing Booklet **Marks**

- (a) Evaluate $\sum_{r=2}^4 \frac{1}{r(r-1)}$. **2**
- (b) The equation of a parabola is $8y = x^2 - 4x - 4$.
 By expressing the equation in the form $(x-h)^2 = 4a(y-k)$, find:
- (i) the coordinates of the vertex. **2**
- (ii) the coordinates of the focus. **1**
- (iii) the equation of the directrix. **1**

(c)




$ABCD$ is a trapezium and AB is parallel to DC . $\angle BAC = 24^\circ$, $\angle ADC = 78^\circ$. **2**
 Explain why $\triangle ACD$ is isosceles.

- (d) Consider the function $f(x) = \begin{cases} 3 + 2^{-x} & \text{for } x \geq 0 \\ x + 4 & \text{for } x < 0 \end{cases}$.
- (i) Evaluate $f(-1) + f(1)$. **2**
- (ii) Find $\lim_{x \rightarrow 0} f(x)$. **1**
- (iii) Is the function continuous at $x = 0$? Give reasons. **1**

QUESTION 9 (12 MARKS) Use a SEPARATE writing Booklet **Marks**

- (a) The value of a car depreciates exponentially according to the formula $V = Ae^{-kt}$ where V dollars is the value of the car after t years. A car's new price is \$32 000, and after 6 years its value is \$14 000.
- (i) Show that $V = Ae^{-kt}$ satisfies the equation $\frac{dV}{dt} = -kV$. 1
 - (ii) What is the value of A ? 1
 - (iii) Find the value of k correct to 3 significant figures. 2
 - (iv) What will be the value of the car after 10 years? 1
 - (v) At what rate will the value of the car be decreasing when it is 10 years old? 1
- (b) A particle moves along a straight line. Its position x metres from O , t seconds after starting, is given by $x = 3 + \cos 2t$.
- (i) Find the position of the particle after $\frac{3\pi}{4}$ seconds. 1
 - (ii) In what direction does the particle start to move? 2
 - (iii) Sketch the graph of v as a function of t for $0 \leq t \leq 2\pi$. 2
 - (iv) What is the maximum acceleration of the particle? 1

QUESTION 10 (12 MARKS) Use a SEPARATE writing Booklet **Marks**



The area between the curve $y = 2^x$ and the x -axis for $0 \leq x \leq 2$, is rotated about the x -axis to form a solid.

- (i) Write down the definite integral which represents the volume of the solid. 1
 - (ii) Use Simpson's Rule with three function values to find the approximate volume of the solid giving your answer to 2 decimal places. 2
 - (iii) Using integration and the result $\int a^x dx = \frac{1}{\ln a} a^x$, evaluate the volume of the solid, giving your answer to 2 decimal places. 2
- (b) Consider the function $f(x) = xe^{-x}$.
- (i) Show that the graph of $y = f(x)$ has one stationary point at $x = 1$. 2
 - (ii) Show that the stationary point is a maximum. 2
 - (iii) Show that there is one point of inflexion and it occurs at $x = 2$. 1
 - (iv) By considering the signs of x and e^{-x} for large positive and large negative values of x , sketch the graph of $y = f(x)$. 2

End of paper


Mathematics: Question 1

Suggested Solutions	Marks Awarded	Marker's Comments
(a) $C = 2\pi r \therefore r = \frac{C}{2\pi}$ $r = \frac{9.29}{2\pi} = 1.4785 \dots$ $= 1.48 \text{ (3 s.f.)}$ (2)		
(b) $\frac{d}{dx} (3x^2 - 4\sqrt{x}) = 6x - 4 \times \frac{1}{2} x^{-\frac{1}{2}}$ $= 6x - \frac{2}{\sqrt{x}}$ (2)		
(c) $(\sqrt{6} + \sqrt{2})^2 = 6 + 2\sqrt{12} + 2$ $= 8 + 4\sqrt{3}$ (2)		
(d) $a - \frac{a-2}{4} = 4$ $4a - (a-2) = 16$ $3a + 2 = 16$ $a = 4\frac{2}{3}$ (2)		
(e) $2c+3 > 1$ $2c+3 < -1$ or $2c+3 > 1$ $2c < -4$ or $2c > -2$ $c < -2$ or $c > -1$ (2)		
(f) $(x - \frac{2}{3})(x+4) = 0$ $x^2 + 4x - \frac{2}{3}x - \frac{8}{3} = 0$ $3x^2 + 10x - 8 = 0$ (2) <u>OR</u> $(3x-2)(x+4) = 0$ $3x^2 + 10x - 8 = 0$ <u>OR</u> $x^2 - (\frac{2}{3} + 4)x + (\frac{2}{3})(-4) = 0$ $14. x^2 + 3\frac{1}{3}x - \frac{8}{3} = 0$ $15. 3x^2 + 10x - 8 = 0$		

Mathematics: Question 2

Suggested Solutions	Marks Awarded	Marker's Comments
(i) Grad. AB: $\frac{3-1}{2-4} = -\frac{1}{2}$ (1)		
(ii) M: $(\frac{-2+4}{2}, \frac{3+1}{2}) = (1, 2)$ (1)		
(iii) AB: $y-3 = -\frac{1}{2}(x+3)$ or $y-1 = -\frac{1}{2}(x-1)$ $2y-6 = -x-3$ or $2y-2 = -x+1$ $x+2y-7=0$ or $x+2y-7=0$ (1)		
(iv) L: $y-2 = 3(x-1)$ $y-2 = 3x-3$ $3x-y-1=0$ (1)		
(v) Subst. $x=0$ into L: $0-y-1=0$ $y=-1 \therefore C$ is $(0, -1)$ (1)		
(vi) $AC = \sqrt{(4-0)^2 + (2-1)^2} = \sqrt{16+1} = \sqrt{17}$ $BC = \sqrt{(4-0)^2 + (1-1)^2} = \sqrt{16+0} = 4$ $\therefore C$ is equidistant from A and B (2)		
(vii) C: $(0, -1)$ AB: $x+2y-7=0$ $d = \frac{ 0+2(-1)-7 }{\sqrt{1^2+2^2}} = \frac{9}{\sqrt{5}} = \frac{9\sqrt{5}}{5}$ (2) <u>OR</u> Dist. CM = $\sqrt{(4-0)^2 + (2-(-1))^2} = \sqrt{17}$		
(viii) $AB = \sqrt{6^2 + 2^2} = \sqrt{40}$ or $2\sqrt{10}$ Area $\triangle ABC = \frac{1}{2} \times 2\sqrt{10} \times \frac{9\sqrt{5}}{5}$ $= 18\sqrt{2}$ (2)		
(ix) Subst. $x=0, y=0$ in $x+2y-7 < 0$ LHS = $-7 < 0$ $\therefore$ Region is $x+2y-7 < 0$ (1)		

Mathematics: Question 3	Marks Awarded	Marker's Comments
Suggested Solutions (a) (i) $\frac{d}{dx} x^3 \cos x = \cos x (3x^2) + x^3 (-\sin x)$ $= 3x^2 \cos x - x^3 \sin x$ (2) (ii) $\frac{d}{dx} (3 + e^{2x})^4 = 4(3 + e^{2x})^3 \cdot 2e^{2x}$ $= 8e^{2x} (3 + e^{2x})^3$ (2) (b) (i) $\int \frac{6x}{x^2+2} dx = 3 \int \frac{2x}{x^2+2} dx$ $= 3 \log_e (x^2+2) + C$ (2) (ii) $\int_0^{\frac{\pi}{2}} \sec^2 2x dx = \left[\frac{1}{2} \tan 2x \right]_0^{\frac{\pi}{2}}$ $= \frac{1}{2} \tan \pi - \frac{1}{2} \tan 0$ $= \frac{\sqrt{2}}{2}$ (2) (c) $\frac{dy}{dx} = 6x^2 - 4x + 1$ (i) $y = 2x^3 - 2x^2 + x + C$ $x=1, y=6: 6 = 2 - 2 + 1 + C$ $C = 5$ Curve is $y = 2x^3 - 2x^2 + x + 5$ (2) (ii) When $x=1, \frac{dy}{dx} = 6 - 4 + 1$ $= 3$ Egn. of tangent is $y - 6 = 3(x - 1)$ (2) i.e. $3x - y + 3 = 0$ or $y = 3x + 3$		


Mathematics: Question 4	Marks Awarded	Marker's Comments
Suggested Solutions (a) $f(x) = x^3 - 6x^2 + 9x$ (i) $f'(x) = 3x^2 - 12x + 9$ $= 3(x^2 - 4x + 3)$ $= 3(x-1)(x-3) = 0$ when $x=1, 3$ $f''(x) = 6x - 12$ $f'(1) = 0, f''(1) = 6 - 12 < 0$ $\therefore$ Maximum at $(1, 4)$ $f'(3) = 0, f''(3) = 18 - 12 > 0$ $\therefore$ Minimum at $(3, 0)$ (3) (ii) Pt. of inflexion: $6x - 12 = 0$ $x = 2$ If $x < 2, f''(x) < 0$; If $x > 2, f''(x) > 0$ Change in concavity pt. of inflexion at $x = 2$ (2) (iii) When $y=0, x^3 - 6x^2 + 9x = 0$ $x(x-3)^2 = 0$ Meets x-axis at $0, 3$ (1) (iv)  (1) (v) Grad. decreasing for $x < 2$. (1) (b) (i) $A = 2 \int_0^4 x x^{\frac{1}{2}} dx = 4 \left[\frac{2}{3} x^{\frac{3}{2}} \right]_0^4$ $= \frac{8}{3} [4^{\frac{3}{2}} - 0]$ $\text{Area} = 21\frac{1}{3} \text{ unit}^2$ (2) (ii) $V = \pi \int_0^4 4x dx = \pi [2x^2]_0^4$ $= 2\pi [4^2 - 0]$ $\text{Volume} = 32\pi \text{ unit}^3$ (2)		


Mathematics: Question 5	Suggested Solutions	Marks Awarded	Marker's Comments
(a)	10, 12, 14, ...		
(i)	$T_{30} = a + (n-1)d = 10 + 24 \times 2$ Time = 68 minutes (1)		
(ii)	$S_{30} = \frac{a}{2}(a + l)$ or $\frac{a}{2}[2a + (n-1)d]$ $= \frac{30}{2}(10 + 68)$ Total time = 1170 minutes (2)		
(iii)	40 hours = 2400 minutes. $\frac{a}{2}[2a + (n-1)d] = 2400$ $\frac{a}{2}[20 + (n-1)2] = 2400$ $n(10 + n-1) = 2400$ $9n + n^2 - 2400 = 0$ $n = \frac{-9 \pm \sqrt{81 - 4(-2400)}}{2}$ $= 44.7$ (since $n > 0$) 40 hours reached on 45 th day (3)		
(b)(i)	$T_{20} = ar^{n-1} = 30\,000 \times (1.05)^{19}$ Salary = \$75 809 (2)		
(ii)	$30\,000 \times 1.05^{n-1} > 50\,000$ $1.05^{n-1} > 1\frac{1}{3}$ $(n-1) \log 1.05 > \log 1\frac{1}{3}$ $n-1 > \frac{\log 1\frac{1}{3}}{\log 1.05}$ $n > 11.47$ (2) Exceeds \$50 000 in 12 th year		
(iii)	$S_{20} = \frac{a(r^n - 1)}{r - 1} = \frac{30\,000(1.05^{20} - 1)}{1.05 - 1}$ Total salary = \$991 979 (2)		

Mathematics: Question 6	Suggested Solutions	Marks Awarded	Marker's Comments
(a)(i)	$AC^2 = 8^2 + 13^2 - 2 \times 8 \times 13 \times \cos 65^\circ$ $AC = 12.05 \text{ m.}$ (2)		
(ii)	$\frac{\sin \angle ACB}{8} = \frac{\sin 65^\circ}{12.05}$ (2) $\sin \angle ACB = \frac{8 \sin 65^\circ}{12.05} = 0.6017$ $\angle ACB = 37^\circ$ (nearest degree)		
(iii)	Area = $\frac{1}{2} \times 8 \times 13 \times \sin 65^\circ$ $= 47.12 \text{ m}^2$ Does <u>not</u> meet regulation. (1)		
(b)	$A = \frac{1}{2} \times 10 \times 1.2$ $B = 1.2$ Perimeter = $10 + 10 + 10 \times 1.2$ $= 32 \text{ m.}$ (3)		
(c)	 (i) In $\triangle AEB$, $\triangle CED$, $\begin{cases} \angle AEB = \angle CED \text{ (vertically opposite)} \\ \angle EAB = \angle EDC \text{ (alternate, } AB \parallel DC) \\ \angle EBA = \angle ECD \text{ (alternate, } AB \parallel DC) \end{cases}$ $\therefore \triangle AEB \parallel \triangle CED$ (2 angles equal) (2)		
(ii)	Let $Bx = x \text{ cm}$, then $Dx = (3-x) \text{ cm}$ $\frac{4}{x} = \frac{16}{3-x}$ $48 - 4x = 16x$ $48 = 20x$ $x = \frac{48}{20}$ $Bx = \frac{24}{5} \text{ cm (or equivalent)} \quad (2)$		

Mathematics Question 7		Marks Awarded	Marker's Comments
(a) $2x - y - 2 = 0$ or $y = 2x - 2$ $2x - y = 4$ $x(2x - 2) = 4$ $2x^2 - 2x - 4 = 0$ $x^2 - x - 2 = 0$ $(x - 2)(x + 1) = 0$ $x = 2, -1$ (3) Points of intersection: $(2, 2), (-1, -4)$			
(b) $3 \times 3^{2x} + 26 \times 3^x - 9 = 0$ Let $u = 3^x$, $3u^2 + 26u - 9 = 0$ $(3u - 1)(u + 9) = 0$ $u = \frac{1}{3}$ or $u = -9$ $3^x = \frac{1}{3}$ or $3^x = -9$ $x = -1$ (no solution) (3)			
(c) $x^2 + 5x + 6 = 0$ For real and different roots, $a > 0$ $25 - 4c > 0$ $c < 6\frac{1}{4}$ (2)			
(d) $2x^2 - 5x - 8 = 0$ $\alpha + \beta = \frac{5}{2}$, $\alpha\beta = \frac{-8}{2} = -4$ (1) $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$ $= \frac{25}{4} - 2(-4)$ $= 14\frac{1}{4}$ (2)			
(ii) $\frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \frac{\alpha^2 + \beta^2}{\alpha\beta}$ $= \frac{14\frac{1}{4}}{-4}$ $= -\frac{57}{16}$ (2)			

Mathematics Question 8		Marks Awarded	Marker's Comments
(a) $\sum_{r=2}^n \frac{1}{r(r-1)} = \frac{1}{2 \times 1} + \frac{1}{3 \times 2} + \frac{1}{4 \times 3}$ (2) $= \frac{3}{4}$			
(b) $2y = x^2 - 4x + 4$ $x^2 - 4x + 4 = 2y + 4 + 4$ $(x - 2)^2 = 2(y + 4)$ NB $a = 2$. (i) Vertex $(2, -4)$ (2) (ii) Focus $(2, -1)$ (1) (iii) Directrix $y = -5$ (1)			
(c)  $\angle ACD = \angle BAC$ (alternate, $AB \parallel DC$) $= 24^\circ$ $\angle DAC = 180^\circ - (78^\circ + 24^\circ)$ (angle sum of $\triangle ACD$) $= 78^\circ$ $\therefore \angle DAC = \angle ADC$ $\triangle ACD$ is isosceles (2)			
(d) $f(x) = \begin{cases} 3 + 2^{-x} & \text{for } x < 0 \\ x + 4 & \text{for } x \geq 0 \end{cases}$ (i) $f(-1) + f(1) = (-1 + 4) + (3 + 2^{-1})$ $= 3 + 3 + \frac{1}{2}$ $= 6\frac{1}{2}$ (2)			
(ii) $\lim_{x \rightarrow 0} f(x) = 3 + 0 = 3$ (1)			
(iii) As $x \rightarrow 0$ from below, $f(x) \rightarrow 0 + 4 = 4$ As $x \rightarrow 0$ from above, $f(x) \rightarrow 3 + 2^0 = 4$ $\therefore f(x)$ is continuous at $x = 0$. (1)			

Mathematics: Question 9		Marks Awarded	Marker's Comments
Suggested Solutions			
(a) $V = Ae^{-kt}$			
(i) $\frac{dV}{dt} = -kAe^{-kt}$ $= -kV$		(1)	
(ii) When $t=0$, $V=32\,000$ $32\,000 = Ae^0$ $\therefore A = 32\,000$		(1)	
(iii) $t=6$, $V=14\,000$: $14\,000 = 32\,000e^{-6k}$ $e^{-6k} = \frac{14\,000}{32\,000}$ $-6k = \log_e\left(\frac{14}{32}\right)$ $k = 0.102 \text{ (3 s.f.)}$		(2)	
(iv) When $t=10$, $V = 32\,000 \times e^{-0.102 \times 10}$ Value = \$8051		(1)	
(v) When $t=10$, $\frac{dV}{dt} = -0.102 \times 8051$ $= -1111$ Value decreasing at \$1111 per year		(1)	
(b) $x = 3 + 10s \sin t$			
(i) When $t = \frac{3\pi}{2}$, $x = 3 + \cos\left(2k \times \frac{3\pi}{2}\right)$ $= 3$		(1)	
(ii) $v = -2 \sin 2t$ It starts to move in negative direction		(2)	
(iii) 		(2)	
(iv) $a = -4 \cos 2t$ Maximum acceleration is 4 m/s^2		(1)	

Mathematics: Question 10		Marks Awarded	Marker's Comments								
Suggested Solutions											
(a) (i) $V = \pi \int_0^2 (2^x)^2 dx$		(1)									
(ii) <table border="1" data-bbox="1294 236 1572 300"> <tr> <td>x</td><td>0</td><td>1</td><td>2</td></tr> <tr> <td>$f(x)$</td><td>1</td><td>4</td><td>16</td></tr> </table> $\int_0^2 (2^x)^2 dx = \frac{1}{\ln 4} [1 + 4 + 16]$ $= 11$ Volume = $11 \times \pi = 34.56 \text{ unit}^3 \text{ (2 d.p.)}$		x	0	1	2	$f(x)$	1	4	16	(2)	
x	0	1	2								
$f(x)$	1	4	16								
(iii) $V = \pi \int_0^2 4^x dx$ $= \pi \left[\frac{1}{\ln 4} 4^x \right]_0^2$ $= \frac{\pi}{\ln 4} [16 - 1]$ Volume = $33.99 \text{ unit}^3 \text{ (2 d.p.)}$		(2)									
(b) $f(x) = xe^{-x}$											
(i) $f'(x) = e^{-x} + x(-e^{-x})$ $= e^{-x}(1-x)$ $= 0$ only when $x=1$ $\therefore$ only one stationary point		(2)									
(ii) $f''(x) = (1-x)(-e^{-x}) + e^{-x}(-1)$ $= xe^{-x} - 2e^{-x} = e^{-x}(x-2)$ $f''(1) = 1e^{-1} - 2e^{-1} < 0$ $\therefore$ Stat. point is maximum at $x=1$ OR show derivative changes sign		(2)									
(iii) For $f'(x) = 0$, $x=2$ $\therefore x=2$ is the only point of inflection		(1)									
(iv) For $x > 0$, $e^{-x} > 0$. $f(x) > 0$ For $x < 0$, $e^{-x} > 0$. $f(x) < 0$											
		(2)									