




SAINT IGNATIUS' COLLEGE

Trial Higher School Certificate

2010

MATHEMATICS

Directions to Students

• Reading Time : 5 minutes	• Total Marks 120
• Working Time : 3 hours	
• Write using blue or black pen. (sketches in pencil).	• Attempt Question 1 – 10
• Board approved calculators may be used	• All questions are of equal value
• A table of standard integrals is provided at the back of this paper.	
• All necessary working should be shown in every question.	
• Answer each question in the booklets provided and clearly label your name and teacher's name.	

This paper has been prepared independently of the Board of Studies NSW to provide additional exam preparation for students. Although references have been reproduced with permission of the Board of Studies NSW, the publication is in no way connected with or endorsed by the Board of Studies NSW.

Question 1 (Start a new Booklet)**Marks**

- (a) Calculate $e^{3.1}$ correct to 2 significant figures 2
- (b) Solve for x : $x^2 + 5x = 24$ 2
- (c) Find the primitive of $(2x + 5)^4$ 2
- (d) Calculate the exact value of $\cos \frac{\pi}{6}$ 2
- (e) Expand and simplify $(3 + 2\sqrt{2})^2$ 2
- (f) Find the sum of the first 40 terms of the series $4 + 10 + 16 + \dots$ 2

Question 2 (Start a new Booklet)**Marks**

- (a) Find the derivative of
- (i) $x \sin 3x$ 2
- (ii) $\frac{e^x}{x}$ 2
- (b) Integrate the following
- (i) $\frac{3x}{x^2 + 4}$ 2
- (ii) $3 \sec^2 \frac{x}{2}$ 1
- (c) Find the equation of the tangent to the curve $y = x^3 + 1$ at the point where $x = 1$. 3
- (d) Evaluate $\sum_{n=1}^3 n^3 + 3$ 2

Question 3 (Start a new Booklet)**Marks**

- (a) $A(1, -4)$ is a point on the line $J: 3x + 2y + 5 = 0$
- (i) Show that the point $B(-3, 2)$ lies on the line. 1
 - (ii) Find the equation of the line perpendicular to J passing through the point $C(3, 1)$. 2
 - (iii) Calculate the distance AB . 2
 - (iv) Find the perpendicular distance from C to the line J . 2
 - (v) Calculate the area of $\triangle ABC$. 1
- (b) The Pacific Star cruise ship travels 215 km on a bearing of 085° from Sydney. It then travels 112 km on a bearing of 135° .
- (i) How far is the ship from Sydney? 2
 - (ii) What is the final bearing of the ship from Sydney?
(give answer correct to the nearest degree) 2

Question 4 (Start a new Booklet)**Marks**


- (a) For the parabola $x^2 + 4x - 12y + 40 = 0$:
- (i) Use completing the square method to write the equation in the form $(x - h)^2 = 4a(y - k)$ 1
 - (ii) Find the focal length. 1
 - (iii) Write down the coordinates of the vertex. 1
 - (iii) Find the focus. 1
- (b) Solve $\log x + \log (x + 4) = \log 12$ 2
- (c) William drops a ball out of a window that is 25 m above the ground. On the first rebound, it rises to a height of 20 m. On subsequent rebounds, it rises to a height equal to $\frac{4}{5}$ of its previous height. If there is no interference with the ball, calculate the total distance through which the ball moves before coming to rest. 2
- (d) At a Primary School Sports Carnival the combined year relay race has one participant from each year group from Kinder to Year 6. The Kinder child runs 15 m to a point and returns to the start, then the year 1 student runs 20 m and returns to the start. Each child runs in turn with each year group running 5 m further than the previous year group.
- (i) How far does the year 6 child have to run? 2
 - (ii) How far is run by the students in one complete race? 2

Question 5 (Start a new Booklet)

Marks

- (a) Copy the following diagram into your answer booklet.

2



The curve represents the function $y = f(x)$. On the same set of axes draw the derivative function $y = f'(x)$.


- (b) Consider the function $y = 2x^3 - 9x^2 + 12x$.

- | | | |
|-------|--|---|
| (i) | Show that the only x -intercept exists at the origin. | 1 |
| (ii) | Find the stationary points and determine their nature. | 3 |
| (iii) | Show that a point of inflexion exists at the point $(1.5, 4.5)$. | 1 |
| (iv) | For what values of x is the curve monotonically decreasing? | 1 |
| (v) | Sketch the curve of the function $y = 2x^3 - 9x^2 + 12x$ in the domain $0 \leq x \leq 3$. | 2 |
| (vi) | Find the values of k for which $2x^3 - 9x^2 + 12x = k$ has only one solution. | 2 |

Question 6 (Start a new Booklet)

Marks

(a)



In the diagram above $\triangle XYZ$ is right angled. PQ is parallel to YZ and Q is the midpoint of XZ .

- | | | |
|-------|---|---|
| (i) | Copy the diagram into your answer booklet. | |
| (ii) | Give a reason why $\angle XPQ = 90^\circ$. | 1 |
| (iii) | Prove that $\triangle XPQ \equiv \triangle YPQ$. | 2 |
| (iv) | Prove $QZ = QY$. | 1 |
| | | |
| (b) | A cylindrical tank is filled with water. The volume of water in the tank is determined by the function $V = 7t^3 + 15t^2 - 3t$, where t is time in seconds and the volume in litres. What is the rate of change of the volume of the tank after 12 seconds have elapsed? | 2 |
| | | |
| (c) | Find the exact length of the radius of a circle in which an arc length of 10 cm subtends an angle of 50° at the centre of the circle. | 2 |
| | | |
| (d) | Given that α and β are the roots of the quadratic equation $2x^2 - 7x + 5 = 0$, find ; | |
| (i) | $\alpha + \beta$ | 1 |
| (ii) | $\alpha\beta$ | 1 |
| (iii) | $\alpha^2 + \beta^2$ | 2 |

Question 7 (Start a new Booklet)

Marks

(a) Prove that $\frac{\sin \theta}{1 + \cos \theta} = \frac{1 - \cos \theta}{\sin \theta}$

3

(b) The area bound by the curve $y = \sqrt{\sin x}$, $x = 0$, $x = \frac{\pi}{3}$ and the x -axis is rotated about the x -axis. Find the volume of the solid formed.

3


(c) A function $f(x)$ has a table of values

2

x	0	0.5	1	1.5	2
f(x)	3.24	4.16	2.25	1.15	0

Use Simpson's Rule to calculate $\int_0^2 f(x) dx$ (correct to 2 decimal places)

(d)



The graph above shows the sketch of the curves $y = \sqrt{3} \sin x$ and $y = \cos x$.

(i) Solve the curves simultaneously to show that the x -values of the points of intersection are $\frac{\pi}{6}$ and $\frac{7\pi}{6}$.

1


(ii) Find the shaded area in the diagram.

3

Question 8 (Start a new Booklet)

Marks

(a)



- | | | |
|-------|--|---|
| (i) | The curve $f(x)$ displayed above is in the form $f(x) = ae^x + 3$. Given that the curve passes through the point $(0, 3.5)$, show that $a = 0.5$. | 1 |
| (ii) | The x -value of the point P on the curve is 2. What is the y -value of the point P ? | 1 |
| (iii) | Find the area of the shaded region. | 2 |
- (b) A particle moves such that its position, x metres, from a fixed point O is given by the function $x = t^3 - 7\frac{1}{2}t^2 + 18t + 2$, where t is measured in seconds.
- | | | |
|-------|---|---|
| (i) | Find the particle's initial position and velocity. | 3 |
| (ii) | When is the particle at rest? | 1 |
| (iii) | What is the acceleration of the particle when it is first at rest? | 2 |
| (iv) | Find the distance travelled by the particle in the first three seconds. | 2 |

Question 9 (Start a new Booklet)**Marks**

- (a) Jenni invests \$30000 into an account on the 1st of March. She receives 9% p.a. interest compounded monthly. On the first day of each month after that she withdraws \$250 immediately after the interest is paid.
- (i) How much money did she have in the account immediately after making the first withdrawal? 1
- (ii) Show that after making the n th withdrawal the balance of the account is given by $\left(33\,333\frac{1}{3} - 3\,333\frac{1}{3} \times 1.0075^n\right)$ 2
- (iii) Find the number of withdrawals that Jenni can make before there is no money left in the account. 2
- (b) An ant colony has a population that is described by the function $P = P_0 e^{kt}$. If the ant colony initially had 300 ants and after 100 days the population had increased to 550 ants, find:
- (i) the value of P_0 and k . 3
- (ii) the time taken for the population of the colony to reach 1000 ants (write your answer correct to the nearest day). 2
- (c) It was found that on the 1st of June, 30 students in a school had the flu. Over the next month the number of cases of students being sick with the flu increased at decreasing rate. Draw a graph that would describe this situation. 2

Question 10 (Start a new Booklet)**Marks**

- (a) Find the derivative of the function $y = x \log_e x$ and hence find $\int \log_e x \, dx$. 2
- (b) A sector of a circle has an area of $\frac{3\pi}{4} \text{ cm}^2$, while its arc length is $\frac{\pi}{4} \text{ cm}$. Find the radius and angle of the sector. 4
- (c) A cylindrical can is to hold $20\pi \text{ m}^3$. The material for the top and bottom costs $\$10/\text{m}^2$ and material for the side costs $\$8/\text{m}^2$.
- (i) Show that the total cost of the material for the can be expressed by the formula $C = 20\pi r^2 + 16\pi r h$ 1
- (ii) Show that $h = \frac{20}{r^2}$ 1
- (iii) Find an expression for the cost in terms of r and hence find the values of r and h such that the cost of the materials is a minimum. 4

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$

2010 Riverview Mathematics Trials – Solutions

Question 1:

(a) 22 (2 significant figures)

$$\begin{aligned} \text{(b)} \quad x^2 + 5x - 24 &= 0 \\ (x + 8)(x - 3) &= 0 \\ x &= -8 \text{ or } x = 3 \end{aligned}$$

$$\text{(c)} \quad y = \frac{(2x+5)^3}{10} + C$$

$$\text{(d)} \quad \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$$

$$\text{(e)} \quad 9 + 12\sqrt{2} + 8 = 17 + 12\sqrt{2}$$

$$\begin{aligned} \text{(f)} \quad S_n &= \frac{n}{2}(2a + 6(n-1)) \\ \therefore S_{40} &= 20(8 + 6 \times 39) = 4840 \end{aligned}$$

Question 2:

(a) (i)

$$\text{Let } u = x \text{ therefore } \frac{du}{dx} = 1$$

$$\text{Let } v = \sin 3x \text{ therefore } \frac{dv}{dx} = 3 \cos 3x$$

$$\text{Since } \frac{dy}{dx} = u'v + v'u \text{ we have:}$$

$$\frac{dy}{dx} = \sin 3x + 3x \cos 3x$$

(ii)

$$\text{Let } u = e^x \text{ therefore } \frac{du}{dx} = e^x$$

$$\text{Let } v = \frac{1}{x} \text{ therefore } \frac{dv}{dx} = -\frac{1}{x^2}$$

$$\text{Since } \frac{dy}{dx} = u'v + v'u \text{ we have:}$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{e^x}{x} - \frac{e^x}{x^2} \\ &= \frac{e^x}{x^2}(x-1) \end{aligned}$$

(b) (i)

$$\begin{aligned} &\int \frac{3x}{x^2 + 4} dx \\ &= \frac{3}{2} \ln |x^2 + 4| + C \end{aligned}$$

(ii)

$$\begin{aligned} &\int 3 \sec^2 \frac{x}{2} dx \\ &= 6 \tan \frac{x}{2} + C \end{aligned}$$

$$\text{(c)} \quad \frac{dy}{dx} = 3x^2$$

$$\text{Therefore at } x = 1, m_T = 3$$

$$y - 2 = 3(x - 1)$$

$$y = 3x - 1 \text{ (or in general form: } 3x - y - 1 = 0)$$

$$\text{(d)} \quad (1 + 3) + (8 + 3) + (27 + 3) = 4 + 12 + 30$$

Q 3 a)

A(1, -4) $3x + 2y + 5 = 0 \leftarrow$

i) sub $x = -3, y = 2$ into J:

LHS:

$3(-3) + 2(2) + 5 \Rightarrow$

$-9 + 4 + 5 = 0 = RHS$

$\therefore P(-3, 2)$ lies on the line.

Let line be R

ii) $\frac{m_R}{m_J} = -1$

$\therefore m_J = -\frac{3}{2}$
C(3, 1)

$m_R = \frac{2}{3}$

$y - 1 = \frac{2}{3}(x - 3)$

$3y - 3 = 2x - 6$

$\therefore 2x - 3y - 3 = 0$

iii) C(3, 1)

$a = 3, x = 3$

$b = 2, y = 1$


$c = 5$

$P = \frac{|a^2 + b^2 + c|}{\sqrt{a^2 + b^2}} = \frac{|3(3) + (2)(1) + 5|}{\sqrt{3^2 + 2^2}}$

$= \frac{9 + 2 + 5}{\sqrt{13}}$

$P = \frac{16}{\sqrt{13}} \text{ units}$

$P = \frac{16}{\sqrt{13}} u$




$d_{AB} = 2\sqrt{13}$

$d_P = \frac{16}{\sqrt{13}}$

$A = \frac{1}{2}ab$

$= \frac{1}{2} \cdot \frac{16}{\sqrt{13}} \cdot 2\sqrt{13} = 16$

b)



Let $S' = \text{Sydney}$

T = Turn

F = Stop

Join $S'F$

$\angle FTS = 45^\circ$ (adjacent angles add up to 180°)

$\angle S'TS = 85^\circ$ (adjacent angles of parallel lines are eq. & c.)

$\angle S'TF = 85 + 45 = 130^\circ$

$S'F = ?$

$a^2 = b^2 + c^2 - 2bc \cos \phi$

$(S'F)^2 = (S'T)^2 + (TF)^2 - 2(TF)(S'T) \cos \phi$

$(S'F)^2 = (215)^2 + (112)^2 - 2(112)(215) \cos 130^\circ$

$= 58769 - 9816 \cos 130^\circ$

$S'F = \sqrt{89725.65} = 299.5424031 \text{ km}$

$S'F = 299.54 \text{ km (2 d.p.)}$

ii) Let $\angle TFS' = \phi$

$\frac{\sin \phi}{215} = \frac{\sin 130}{299.54}$

$\sin \phi = \frac{215 \sin 130}{299.54}$

$\sin \phi = \frac{215 \sin 130}{299.54}$

$\phi = 33^\circ 21'$

$\angle S'FW_2 = 45^\circ - 33^\circ 21'$

$= 11^\circ 39'$

Bearing is

$270^\circ + 11^\circ 39'$

$= 282^\circ \text{ (measured from)}$

Q 4c)

$$x^2 + 4x - (12y + 40) = 0$$


$$(x+2)^2 - 4 - 12y + 40 = 0$$

$$(x+2)^2 - 12y = -36$$

$$(x+2)^2 = -36 + 12y$$

$$(x+2)^2 = 12(y-3)$$

$$(x+2)^2 = 12(y-3)$$



$$4a = 12$$

$$|a| = 3$$

i) $4a = 12$
 $a = 3$

ii) $(-2, 3)$

iv) $(-2, 6)$

b) $\ln|x| + \ln|x+4| = \ln(12)$

$$\ln|x(x+4)| = \ln(12)$$

$$x^2 + 4x - 12 = 0$$


$$(x+6)(x-2) = 0$$

$$x = 2, x = -6$$

But $\ln|x|$, $x > 0$

$$\therefore x \neq -6$$

$$\{x = 2\}$$



$$T_n = \{40, 32, 25.6, \dots\} \rightarrow \{40, 32, 25.6, \dots\}$$

Let us define as $T_n \rightarrow 0$, it's at rest

$$T_n \text{ is a GP, } r = \frac{T_2}{T_1} = \frac{T_3}{T_2} = 0.8$$

$$T_n = ar^{n-1}$$

$$T = 40(0.8)^{n-1}$$

But window is 25m let define sum R

$$R = 25 + T_n$$

Since $|r| < 0.8$

S_{∞} exist

$$S_{\infty} = \frac{a}{1-r}$$

$$= \frac{40}{1-0.8} = 200m$$

$$R_n = 200 + 25 = 225m$$

$\therefore$ The ball travelled 225m



$$i) T_n = \{30, 40, 50, \dots, 90\}$$

$$T_n = a + (n-1)d$$

$$\text{Let } a = 30$$

$$d = 10$$

$$T_n = 30 + (n-1)10$$

$$= 30 - 10 + 10n$$

$$= 20 + 10n$$

$$\text{Let } n = 7$$

$$T_7 = 20 + 10(7) = 90m$$


$$ii) S_n = T_1 + T_2 + \dots + T_n$$

$$S_n = \frac{n(a+l)}{2}$$

$$= \frac{7}{2} (30 + 90) = 420m$$

$\therefore$ 7 students run 420m altogether

Question 5 (a)



$$b) i) y = 2x^3 - 9x^2 + 12x$$

$$y' = 6x^2 - 18x + 12$$

$$\text{Let } y' = 0 \text{ (TP)}$$

$$0 = x^2 - 3x + 2$$

$$0 = (x-2)(x-1)$$

$$x = 2, x = 1$$

$$y = 4, y = 5$$

$$i) y = x(2x^2 - 9x + 12)$$

$$\text{Let } y = 0, x = 0, 2x^2 - 9x + 12 = 0$$

$$\text{But } \Delta = -15 (< 0)$$

$$\therefore x = 0$$

$\therefore$ no real roots

continued
ii) ~~1.5~~

$$y'' = 12x - 18$$

$$\text{At } x = 1 \text{ in } y'$$

$$y'' = 12 - 18 = -6 (< 0) \text{ Maxime}$$

$$P_1(1, 5) \text{ TP (Max)}$$

$$\text{At } x = 2 \text{ in } y''$$

$$y'' = 12(2) - 18 = 6 (> 0) \text{ Minime}$$

$$P_2(2, 4) \text{ (TP) (Min)}$$

$$\therefore P_1(1, 5) \text{ (TP) (Max)}$$

$$P_2(2, 4) \text{ (TP) (Min)}$$

iii) At $y' = 0$ (IP)

$$0 = 12x - 18$$

$$x = \frac{18}{12} = 1.5$$

$$\text{At } y = 4.5$$

$$IP(1.5, 4.5)$$


iv)

$$\frac{dy}{dx} < 0$$

$$y' = 6x^2 - 18x + 12$$

$$6x^2 - 18x + 12 < 0$$


$$2 \cdot (x-2)(x-1) < 0$$



$\therefore$ graph is decreasing at $1 < x < 2$

v)

(3.9)



$$vi) 2x^3 - 9x^2 + 12x - k = 0$$

~~RMV~~
 $k = 0$ is a possible soln

$$\{ k < 4, k > 5 \}$$

Question 6

c)



ii) $\angle XPQ = 90^\circ = \angle PYZ$ (corresponding angles of parallel lines are equal)

iii) In $\triangle QXP$ and $\triangle XZY$:

$\angle QXP$ is common angle (INCLUDED)

$\angle XPQ = \angle XYZ = 90^\circ$ (angle)

$\therefore \triangle QXP \sim \triangle XZY$ (AAA)

$$\frac{XQ}{QZ} = \frac{XP}{PY} = 1 \text{ (common ratio of similar triangles)}$$

$$\therefore XP = PY$$

In $\triangle XQP$ and $\triangle YPQ$,

QP is common side (SIDE)

$XP = PY$ (SIDE)

$\angle QPY = \angle QPX = 90^\circ$ (INCLUDED ANGLE)

$\therefore \triangle XQP \cong \triangle YPQ$ (SAS)

iv) $QX = QY = QZ$ (given, equal side of congruent triangles)

$$\therefore QZ = QY$$

$$b) V = 7t^3 + 15t^2 - 3t$$

$$\frac{dV}{dt} = 21t^2 + 30t - 3$$


$$\text{At } t = 12$$

$$\frac{dV}{dt} = 21(12)^2 + 30(12) - 3$$

$$= 3381 \text{ L/s}$$

$$\therefore \frac{dV}{dt} = 3381 \text{ L/s at } t = 12 \text{ seconds}$$

c)



$$L = r\theta$$

$$10 = x \left(\frac{50\pi}{180} \right)$$

$$10 = x \left(\frac{5\pi}{18} \right)$$

$$\frac{180}{5\pi} = x$$

$$x = \frac{36}{\pi} \text{ cm}$$

$$d) i) A+B = \frac{-b}{a} = \frac{7}{2}$$

$$ii) AB = \frac{c}{a}$$

$$iii) A^2 + B^2 = (A+B)^2 - 2AB$$

$$= \left(\frac{7}{2}\right)^2 - 2\left(\frac{7}{2}\right)$$

$$= \frac{29}{4}$$

Question 7


$$c) \frac{\sin \theta}{1 + \cos \theta} = \frac{1 - \cos \theta}{\sin \theta}$$

LHS:

$$\frac{\sin \theta (1 - \cos \theta)}{(1 + \cos \theta)(1 - \cos \theta)}$$

$$= \frac{\sin \theta (1 - \cos \theta)}{1 - \cos^2 \theta} \quad 1 - \cos^2 \theta = \sin^2 \theta$$

$$= \frac{\sin \theta (1 - \cos \theta)}{\sin^2 \theta} = \frac{1 - \cos \theta}{\sin \theta} = \text{RHS QED}$$



$$V_y = \pi \int_0^{\pi/3} y^2 dx$$

$$= \pi \int_0^{\pi/3} \sin^2 x dx$$

$$= \pi \left[\cos x \right]_0^{\pi/3}$$

$$= \pi \left[\cos \frac{\pi}{3} - \cos 0 \right]$$

$$= \pi \left[\frac{1}{2} - 1 \right]$$

$$= -\frac{\pi}{2}$$

$$2) \int_0^1 f(x) dx$$

$$A = \frac{1}{3} (f(0) + f(1) + 2 \text{ odd} + 4 \text{ even})$$

$$= \frac{0.5}{3} (0 + 3.24 + 2(2.25) + 4(4.16 + 1.15))$$

$$= \frac{0.5}{3} (28.98) = 4.83 \text{ (2dp)}$$

$$d) i) y = \sqrt{3} \sin x \quad (1)$$

$$y = \cos x \quad (2)$$

$$\frac{(1)}{(2)}$$

$$1 = \sqrt{3} \tan x$$

$$\tan x = \frac{1}{\sqrt{3}} \quad \frac{S(1)}{C}$$

$$x = \frac{\pi}{6}$$

So:

$$x = \frac{\pi}{6}$$

3rd:

$$x = \frac{\pi}{6} + \frac{\pi}{6}$$

$$x = \frac{2\pi}{6}$$

$$\therefore x = \frac{\pi}{6}, \frac{2\pi}{6} \quad \text{QED}$$

$$ii) A = \int_{\pi/6}^{7\pi/6} (\sqrt{3} \sin x - \cos x) dx$$

$$= \left[-\sqrt{3} \cos x - \sin x \right]_{\pi/6}^{7\pi/6}$$

$$= - \left[\left(\sqrt{3} \cos \frac{7\pi}{6} + \sin \frac{7\pi}{6} \right) - \left(\sqrt{3} \cos \frac{\pi}{6} + \sin \frac{\pi}{6} \right) \right]$$

$$= - \left[\left(\frac{-\sqrt{3} \times \sqrt{3}}{2} + \left(-\frac{1}{2} \right) \right) - \left(\frac{\sqrt{3} \times \sqrt{3}}{2} + \frac{1}{2} \right) \right] = - \left(\frac{3}{2} - \frac{1}{2} \right) - \left(\frac{3}{2} - \frac{1}{2} \right)$$

$$= - \left[-2 - 2 \right] = 4$$

$$= 4 \text{ u}^2$$

Question 8

$$a) i) f(x) = ae^x + 3$$

$$\text{sub } (0, 3.5)$$

$$f(0) = ae^0 + 3 = 3.5$$


$$\therefore a(1) = 0.5 \quad a \leq 0$$

$$ii) \text{ sub } x=2$$

$$f(2) = 0.5e^2 + 3$$

$$\therefore \text{the y value} = 0.5e^2 + 3$$

iii)



$$A_{\text{Rectangle}} = (e^2 + 6)u^2$$

$$A_2 = A_{\text{rect}} - A_1$$

$$A_1 = \int_0^2 0.5e^x + 3 \, dx$$

$$= \left[0.5e^x + 3x \right]_0^2$$

$$= 0.5e^2 + 6 - 0.5 = 0.5e^2 + 5.5$$

$$A_2 = e^2 + 6 - (0.5e^2 + 5.5)$$

$$= 0.5e^2 + 0.5$$

$$= 0.5(e^2 + 1)u^2$$

$$\therefore \text{shaded region} = 0.5(e^2 + 1)u^2$$

$$Q9) c) K = \$30000$$

$$K = 1 + \frac{9i}{12} = 1.0075$$

$$n = n$$

$$K = 9250$$

$$i) A_1 = LK$$

$$= 30000(1.0075) = 30225$$

$$ii) A_2 = LK^2 - P$$

$$A_3 = LK^3 - PK - P$$

$$A_1 = LK^4 - P(1 + K + K^2)$$

$$A_n = LK^n - \frac{P(K^n - 1)}{K - 1}$$

$$= LK^n - \frac{PK^n}{K-1} + \frac{P}{K-1}$$

$$A_1 = K^n(L - \frac{P}{K-1}) + \frac{P}{K-1} = 3333\frac{1}{3} - 3333\frac{1}{3}(1.0075)^4$$

Q60

iii) from ii)

$$A_n = Lk^n - P \frac{1}{k-1} \quad \text{if } k \neq 1$$

$$\Rightarrow \frac{dA_n}{dn} = \frac{-Lk^n}{k-1} = \frac{-P}{k-1}$$

$$k^n = \frac{-P}{Lk - L - P}$$

$$k^n = 10$$

$$n \ln k = \ln 10$$

$$n = \frac{\ln(10)}{\ln k} = 309$$

$$\therefore n = 309 \text{ months}$$

She will have no money after 309 months

i) $P = P_0 e^{kt}$

$$\text{at } t=0, P=300$$

$$\therefore P_0 = 300$$

$$\text{at } t=100, P=550$$

$$550 = 300 e^{k(100)}$$

$$\ln\left(\frac{550}{300}\right) = k(100)$$

$$k = \frac{\ln\left(\frac{550}{300}\right)}{100} = 0.00606(35803)$$

$$\therefore P_0 = 300$$

$$k = 0.00606(35803)$$

ii) $k + P = 1000$

$$\frac{1000}{300} = \frac{300}{300} e^{kt}$$

$$\frac{10}{3} = e^{kt}$$


$$\ln\left(\frac{10}{3}\right) = kt$$

$$t = \frac{\ln\left(\frac{10}{3}\right)}{k}$$

$$t = 198.6308$$

$$\therefore t = 199 \text{ days}$$

c)



Q10 a)

$$y = x \ln|x|$$

$$\text{let } u = x \quad u' = 1$$

$$v = \ln|x| \quad v' = \frac{1}{x}$$

$$y' = \ln|x| + 1$$

$$\frac{dy}{dx} = \ln|x| + 1$$

$$\int dy = \int (\ln|x| + 1) dx$$


$$y = \int \ln|x| dx + x$$

$$y - x = \int \ln|x| dx$$

$$x \ln|x| - x = \int \ln|x| dx$$

$$\therefore \int \ln|x| dx = x \ln|x| - x + C$$

b)



$$A = \frac{1}{2} r^2 \theta \quad (2) \quad L = r\theta$$

$$\frac{\pi}{4} = r\theta$$

$$\theta = \frac{\pi}{4r} \quad (1)$$

sub (1) into (2)

$$\frac{3\pi}{4} = \frac{1}{2} r^2 \frac{\pi}{4r}$$

$$3\pi = \frac{1}{2} r\pi$$

$$6 = r$$

$$\begin{cases} r = 6 \\ \theta = \frac{\pi}{24} \end{cases}$$

$$c) i) V = 20\pi \text{ m}^3$$

$$2\pi r^2 \rightarrow 10 \text{ m}^2$$

$$2\pi r h \rightarrow 8 \text{ m}^2$$

ii) - - -

$$C = 10(2\pi r^2) + 8(2\pi r h)$$

$$C = 20\pi r^2 + 16\pi r h \quad QED$$

ii) ~~not a min~~

$$V = \pi r^2 h$$

$$20\pi = \pi r^2 h$$

$$h = \frac{20\pi}{\pi r^2}$$

$$\therefore h = \frac{20}{r^2} \quad QED$$

iii) sub h into C

$$C = 20\pi r^2 + 16\pi r \left(\frac{20}{r^2} \right)$$

$$C = 20\pi r^2 + \frac{320\pi}{r}$$

$$C = 20\pi r^2 + 320\pi r^{-1}$$

$$\frac{dC}{dr} = 40\pi r - 320\pi r^{-2}$$

$$\text{Let } \frac{dC}{dr} = 0 \quad (1P)$$

$$0 = 40\pi r - \frac{320\pi}{r^2}$$

$$40\pi$$

$$0 = r - \frac{8}{r^2}$$

$$0 = \frac{r^3 - 8}{r^2}$$

$$r \neq 0$$

$$r = 2$$

$$h = 5$$

1st Deriv test:

+A	2	2.1
\	-	/

$\therefore$ at $r = 2$, it is a minimum

$$\therefore r = 2, h = 5$$

$$\therefore \begin{cases} r = 2 \text{ m} \\ h = 5 \text{ m} \end{cases}$$