



**BAULKHAM HILLS
HIGH
SCHOOL**

2023

**YEAR 11
YEARLY
EXAMINATIONS**

Mathematics Extension

General Instructions

- Reading time – 10 minutes
- Working time – 2 hours
- Write using black pen
- Calculators approved by NESA may be used
- A reference sheet is provided at the back of this paper
- In Questions in Section II, show relevant mathematical reasoning and/or calculations
- Marks may be deducted for careless or badly arranged work

Total marks: 70

Section I – 10 marks (pages 2 – 5)

- Attempt Questions 1 – 10
- Allow about 15 minutes for this section

Section II – 60 marks (pages 6 – 9)

- Attempt Questions 11 – 14
- Allow about 1 hour and 45 minutes for this section

Section I

10 marks


Attempt Questions 1 – 10

Allow about 15 minutes for this section


Use the multiple-choice answer sheet for Questions 1 – 10

- 1 Which of the following graphs represents the solution to $|x - 2| \leq 4$?

(A)



(B)



(C)



(D)



- 2 Which of the following is the natural domain of $f(x) = \cos^{-1}\left(\frac{x-a}{b}\right)$, where $b > 0$?

(A) $[-1, 1]$

(B) $[a - b, a + b]$

(C) $[a - 1, a + 1]$

(D) $[-b, x + b]$

- 3 The population N of a town, after t years, is given by the formula, $N = N_0 e^{0.04t}$ where N_0 is the initial population.

Which expression represents the number of years it takes until the town has doubled its initial population?


(A) $\frac{2\ln 100}{5}$

(B) $4\ln 2$

(C) $2\ln 25$

(D) $25\ln 2$

- 4 In the diagram below, $OA = 3$ metres, $AB = 1$ metre, $OP = x$ metres and $\angle APB = \theta$.



Which of the following equations is correct?

- (A) $\theta = \tan^{-1}\left(\frac{1}{x}\right) - \tan^{-1}\left(\frac{3}{x}\right)$
- (B) $\theta = \tan^{-1}\left(\frac{4}{x}\right) - \tan^{-1}\left(\frac{3}{x}\right)$
- (C) $\theta = \tan^{-1}\left(\frac{1}{\sqrt{9-x^2}}\right)$
- (D) $\theta = \tan^{-1}(3-x)$
- 5 The polynomial $ax^8 + bx^7 + 2$ is divisible by $(x+1)^2$.
What are the values of a and b ?
- (A) $a = -14$ and $b = -16$
- (B) $a = 14$ and $b = -16$
- (C) $a = -14$ and $b = 16$
- (D) $a = 14$ and $b = 16$

6 Which of the following is equal to $\sin x \cos^3 x - \sin^3 x \cos x$?

(A) $\frac{1}{6} \sin 4x$

(B) $\frac{1}{4} \sin 4x$

(C) $\frac{1}{3} \sin 4x$

(D) $\frac{1}{2} \sin 4x$

7 Which of the following is the coefficient of x^n in the expansion of $(1+x)^n + (1-x)^n$?

(A) 0

(B) $\frac{1 + (-1)^n}{2}$

(C) $1 + (-1)^n$

(D) 2

8 Let α , β and γ be the roots of $2x^3 - 5x^2 + 2x - 1 = 0$.

Which equation has the roots $\frac{1}{\alpha}$, $\frac{1}{\beta}$ and $\frac{1}{\gamma}$?

(A) $x^3 - 2x^2 + 5x - 2 = 0$


(B) $x^3 + 2x^2 + 25x - 2 = 0$

(C) $x^3 - 2x^2 + 25x - 8 = 0$


(D) $-2x^3 + 5x^2 - 2x + 1 = 0$

9 Which of the following is the graph of $y = x \cot x$?


(A)




(B)




(C)



(D)



10 Moving only to the right or down, how many different routes are there from A to B ?



(A) 10

(B) 12

(C) 14

(D) 18

END OF SECTION I

Section II

60 marks

Attempt Questions 11 – 14

Allow about 1 hour and 45 minutes for this section

Answer each question on the appropriate answer sheet. Extra paper is available.

For questions in Section II, your responses should include relevant mathematical reasoning and/or calculations.

Question 11 (15 marks) Use the pages labelled Question 11 in the answer booklet

- (a) Sketch $y = 2\tan^{-1}(2x - 1)$. 2
- (b) Consider the function $f(x) = \frac{2}{x+1}$
- (i) Find the inverse function of $f(x)$. 1
- (ii) State the range of the inverse function. 1
- (c) A team of 4 players consists of 2 men and 2 women.
If a total of 7 players are available, of which 3 are men and 4 are women;
- (i) How many different teams can be selected? 1
- (ii) If 2 of the players are husband and wife, and wish to play on the same team, how many different teams can be selected if both are chosen? 1
- (d) Determine the term independent of x in the binomial expansion $\left(x^2 - \frac{1}{x}\right)^9$. 2
- (e) Show that $\frac{1 + \cos x}{\sin x} = \frac{1}{t}$ where $t = \tan \frac{x}{2}$. 2
- (f) Solve $\frac{x^2 + 5}{x} > 6$ 3
- (g) Evaluate $\sin\left[\tan^{-1}\left(\frac{2}{3}\right) - \cos^{-1}\left(\frac{12}{13}\right)\right]$. 2

Question 12 (15 marks) Use the pages labelled Question 12 in the answer booklet

- (a) A curve is defined by the parametric equations $x = 1 + 2t$ and $y = 1 - 4t^2$. 2

Find the cartesian equation of the curve.

- (b) In how many ways can the letters of the word

PAHVANITHAS


be arranged if;

- (i) all of the letters are used. 2
- (ii) all of the A's are together and the arrangement must start and finish with a vowel. 2

- (c) Show that $\frac{\cos 2\theta + \sin 2\theta - 1}{\cos 2\theta - \sin 2\theta + 1} = \tan \theta$. 2

- (d) Solve in simplest exact form, the inequality $e^{2x} - 4e^x + 3 < 0$. 2

- (e) The graph of $y = f(x)$ is shown below.



Sketch graphs of;

- (i) $y = f(|x|)$. 1
- (ii) $y = \frac{1}{f(x)}$. 2
- (iii) $y^2 = f(x)$. 2

Question 13 (15 marks) Use the pages labelled Question 13 in the answer booklet

(a) Show that $\tan\left(\frac{x+y}{2}\right)\cot\left(\frac{x-y}{2}\right) = \frac{\sin x + \sin y}{\sin x - \sin y}$. 2

(b) Without expanding, find 2

$$1 + 5(x-1) + 10(x-1)^2 + 10(x-1)^3 + 5(x-1)^4 + (x-1)^5.$$

(c) Prove $2\tan^{-1}\left(\frac{1}{3}\right) = \tan^{-1}\left(\frac{3}{4}\right)$. 2

(d) Two of the roots of the equation $4x^3 + px^2 - 20x + 12 = 0$ are reciprocals. 3
Find the value of p .

(e) The population of birds on an island changes according to the differential equation

$$\frac{dP}{dt} = k(P - 3500)$$

where P is the population of birds, t is the time in years and k is the constant of proportionality.

Initially there were 40 birds, and after 5 years the population has increased to 125 birds.

- (i) Verify that $P = 3500 - Ae^{kt}$, where A is a constant, satisfies the differential equation. 1
- (ii) Find the value of A . 1
- (iii) Find the value of k correct to 3 significant figures. 2
- (iv) Sketch the function $P = 3500 - Ae^{kt}$ for $t \geq 0$, showing any intercepts and asymptotes. 2

Question 14 (15 marks) Use the pages labelled Question 14 in the answer booklet

(a) A polynomial $P(x)$ is defined by $P(x) = x^3 - b^3$, where b is a constant.

(i) Express $P(x)$ as the product of a linear and a quadratic factor 1

(ii) Simplify $\frac{(\tan^2 \theta - 1)(\sin \theta \cos \theta + 1)}{\cos^2 \theta (\tan^3 \theta - 1)}$ 2


(b) Find the multiplicity of the root $x = 1$ of the equation $P(x) = 0$, where 2

$$P(x) = nx^{n+1} - (n+1)x^n + 1, \quad n \neq 0.$$

(c) Solve the equation $\sin^{-1}(2x^2 - 4x) + \cos^{-1}(2 - x) = \frac{\pi}{2}$ 3

(d) A large container in the shape of an inverted cone with semi-vertical angle 30° is full of water.

A hole is made at the vertex of the cone that allows the container to empty at a constant rate of 12 kL/min.



(i) Show that when the water has a depth of h metres, its volume is given 1
by $V = \frac{1}{9}\pi h^3$.

(You may assume that the volume cut from the cone when creating the hole, is negligible).

(ii) Hence find, in simplest exact form, the rate at which the depth of the water 3
is changing when its volume is 24π kL. (Note: $1 \text{ kL} = 1 \text{ m}^3$)

(e) Let $P(x) = x^2 - 2ax + a^4$ where a is a non-zero real number. 3


When $P(x)$ is divided by $x + b$, the remainder is 1.

The polynomial $Q(x) = bx^2 + x + 1$ has a factor of $(ax - 1)$.

Determine the value(s) of b .

End of paper



BLANK PAGE



Solution		Marks	Comments
QUESTION 14			
14(a) (i) $P(x) = x^3 - b^3$ $= (x-b)(x^2 + bx + b^2)$	1	1 mark • Correct answer	
14(a) (ii) $\frac{(\tan^2 \theta - 1)(\sin \theta \cos \theta + 1)}{\cos^2 \theta (\tan^3 \theta - 1)}$ $= \frac{(\tan \theta + 1)(\tan \theta - 1)(\sin \theta \cos \theta + 1)}{\cos^2 \theta (\tan \theta - 1)(\tan^2 \theta + \tan \theta + 1)}$ $= \frac{(\tan \theta + 1)(\sin \theta \cos \theta + 1)}{\cos^2 \theta \left(\sec^2 \theta + \frac{\sin \theta}{\cos \theta} \right)}$ $= \frac{(\tan \theta + 1)(\sin \theta \cos \theta + 1)}{(1 + \sin \theta \cos \theta)}$	2	2 marks • Correct solution 1 mark • Substantial progress towards the simplification	
14(b) $P(x) = nx^{n+1} - (n+1)x^n + 1$ $P'(1) = n - (n+1) + 1$ $= 0$ $P''(x) = n^2(n+1)x^{n-1} - n(n+1)(n-1)x^{n-2}$ $P''(1) = n^2(n+1) - n(n+1)(n-1)$ $= n^3 + n^2 - n^3 + n$ $= n^2 + n$ $\neq 0$ $\therefore x = 1$ is a root of multiplicity 2	2	2 marks • Correct solution 1 mark • Uses the fact that multiple roots are roots of the derivative as well as the polynomial	
14(c) $\sin^{-1}(2x^2 - 4x) + \cos^{-1}(2 - x) = \frac{\pi}{2}$ $2x^2 - 4x = 2 - x \quad \left(\because \sin^{-1}x + \cos^{-1}x = \frac{\pi}{2} \right)$ $2x^2 - 3x - 2 = 0$ $(2x+1)(x-2) = 0$ $x = -\frac{1}{2} \quad \text{or} \quad x = 2$ however $2 - \left(-\frac{1}{2}\right) = \frac{5}{2} > 1$ and so is outside the domain of the basic curve $y = \cos^{-1}x$, which is $-1 \leq x \leq 1$, thus $x = -\frac{1}{2}$ is not a solution $\therefore x = 2$	3	3 marks • Correct solution 2 marks • Finds two possibilities for solutions using a correct method 1 mark • Uses the inverse trig property in an attempt to find a solution • Finds a correct answer	
14(d) (i) 	1	1 mark • Correct solution	

	Solution	Marks	Comments
14(d) (ii)	$V = \frac{1}{9}\pi h^3$ $\frac{dV}{dh} = \frac{1}{3}\pi h^2$ $= (-12) \left(\frac{3}{\pi h^2} \right)$ $= -\frac{36}{\pi h^2}$ when $V = 24\pi$, $\frac{1}{9}\pi h^3 = 24\pi$ $h^3 = 216$ $h = 6$ $\therefore$ the depth is decreasing at a rate of $\frac{1}{\pi}$ m/min	3	3 marks • Correct solution 2 marks • Finds both the height of the water and an expression for $\frac{dV}{dh}$ 1 mark • Finds the height of the water • Finds an expression for $\frac{dV}{dh}$
14(e)	$P(-b) = 1$ $b^2 + 2ab + a^4 = 1$ $\frac{b}{a^2} + \frac{1}{a} + 1 = 0$ $b + a + a^3 = 0$ $b = -a^2 - a$ $\therefore (-a^2 - a)^2 + 2a(-a^2 - a) + a^4 = 1$ $a^4 + 2a^3 + a^2 - 2a^3 - 2a^2 + a^4 = 1$ $2a^4 - a^2 - 1 = 0$ $(2a^2 + 1)(a^2 - 1) = 0$ $a^2 = -\frac{1}{2} \quad \text{or} \quad a^2 = 1$ $a = \pm 1$ no solutions $\therefore b = -2$ or $b = 0$	3	3 marks • Correct solution 2 marks • Finds an equation in terms of a or b 1 mark • Use the factor or remainder theorem.

Solution	Marks	Comments
12(c) (iii)	2	2 marks - Correct graph 1 mark - Sideways parabola drawn for $0 \leq x \leq 2$ - Horizontal lines $y = \sqrt{2}$ and $y = -\sqrt{2}$ drawn for $2 \leq x \leq 3$ $y = \sqrt{x}$
QUESTION 13		
13(a) $\tan\left(\frac{x+y}{2}\right) \cot\left(\frac{x-y}{2}\right) = \frac{\sin\left(\frac{x+y}{2}\right) \cos\left(\frac{x-y}{2}\right)}{\cos\left(\frac{x+y}{2}\right) \sin\left(\frac{x-y}{2}\right)}$ $= \frac{1}{2} \left[\sin\left(\frac{x+y}{2} + \frac{x-y}{2}\right) + \sin\left(\frac{x+y}{2} - \frac{x-y}{2}\right) \right]$ $= \frac{1}{2} \left[\sin\left(\frac{x-y}{2} + \frac{x+y}{2}\right) + \sin\left(\frac{x-y}{2} - \frac{x+y}{2}\right) \right]$ $= \frac{\sin x + \sin y}{\sin x + \sin y}$ $= \sin x + \sin y$	2	2 marks - Correct solution 1 mark - Correctly transforms a product of two trig functions into a simplified sum of two trig functions
13(b) $1 + 5x + 10x^2 + 10x^3 + 5x^4 + x^5 = (1+x)^5$ $\therefore 1 + 5(x-1) + 10(x-1)^2 + 10(x-1)^3 + 5(x-1)^4 + (x-1)^5 = (1+x-1)^5$ $= x^5$	2	2 marks - Correct solution 1 mark - Recognises the pattern of a binomial expansion
13(c) Let $\theta = \tan^{-1}\left(\frac{1}{3}\right)$ $\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$ $= \frac{2 \times \frac{1}{3}}{1 - \frac{1}{9}}$ $= \frac{\frac{2}{3}}{\frac{8}{9}}$ $= \frac{3}{4}$ $2\theta = \tan^{-1}\left(\frac{3}{4}\right)$ $\therefore 2 \tan^{-1}\left(\frac{1}{3}\right) = \tan^{-1}\left(\frac{3}{4}\right)$	2	2 marks - Correct solution 1 mark - Correctly uses double angle result



Solution	Marks	Comments
13(d) Let the roots be $\alpha, \frac{1}{\alpha}$ and β $\alpha + \frac{1}{\alpha} + \beta = -\frac{p}{4}$ $\beta = -3$ $\alpha + \frac{1}{\alpha} - 3 = -\frac{p}{4}$ $\alpha + \frac{1}{\alpha} = 3 - \frac{p}{4}$ $\left(\alpha\right)\left(\frac{1}{\alpha}\right) + \alpha\beta + \frac{p}{\beta} = -5$ $1 - 3\alpha - \frac{3}{\alpha} = -5$ $3\left(\alpha + \frac{1}{\alpha}\right) = 6$ $3 - \frac{p}{4} = 2$ $\frac{p}{4} = 1$ $p = 4$	3	3 marks - Correct solution 2 marks - Finds an equation where p is the only unknown 1 mark - Finds β
13(e) (i) $P = 3500 + Ae^{kt}$ $\frac{dP}{dt} = -kAe^{kt}$ $= k(3500 - Ae^{kt} - 3500)$ $= k(P - 3500)$ 13(e) (ii) when $t = 0, P = 40 \Rightarrow 40 = 3500 - A$ $A = 3460$ 13(e) (iii) when $t = 5, P = 125 \Rightarrow 125 = 3500 - 3460e^{5k}$ $3460e^{5k} = 3375$ $e^{5k} = \frac{3375}{3460}$ $5k = \ln\left(\frac{3375}{3460}\right)$ $k = \frac{1}{5} \ln\left(\frac{3375}{3460}\right)$ $= -0.00497$ (to 3 sig figs)	1	1 mark - Correct solution
13(e) (iv)	2	2 marks - Correct solution 1 mark - Correct shape - Correct asymptote

Solution		Marks	Comments
11(d) $\left(x^2 - \frac{1}{x}\right)^9$; $T_{k+1} = \binom{9}{k} (x^2)^{9-k} \left(-\frac{1}{x}\right)^k$ $= \binom{9}{k} (-1)^k x^{18-3k}$ independent term has $x^0 \Rightarrow 18-3k=0$ $k=6$ $\therefore$ independent term is $T_7 = \binom{9}{6} (-1)^6 = 84$		2	2 marks • Correct solution 1 mark • Finds an expression for the general term • Recognises that the constant term is when the power of x is zero
11(e) $\frac{1+\cos x}{\sin x} = \frac{1+\frac{1-p^2}{1+p^2}}{\frac{2p}{1+p^2}}$ $= \frac{1+p^2+1-p^2}{2p}$ $= \frac{2}{2p}$ $= \frac{1}{p}$		2	2 marks • Correct solution 1 mark • Correct substitution of p results for either $\sin x$ or $\cos x$
11(f) $\frac{x^2+5}{x} > 6$ $x \neq 0$ $x^2+5=6x$ $x^2-6x+5=0$ $(x-5)(x-1)=0$ $x=5$ or $x=1$ 		3	3 marks • Correct graphical solution on number line or algebraic solution, with correct working 2 marks • Bald answer • Identifies the correct critical points via a correct method • Correct conclusion to their critical points obtained using a correct method 1 mark • Uses a correct method • Acknowledges a problem with the denominator.
11(g) Let $\alpha = \tan^{-1}\left(\frac{2}{3}\right)$ and $\beta = \cos^{-1}\left(\frac{12}{13}\right)$  $\sin\left[\tan^{-1}\left(\frac{2}{3}\right) - \cos^{-1}\left(\frac{12}{13}\right)\right] = \sin\alpha \cos\beta - \cos\alpha \sin\beta$ $= \left(\frac{2}{\sqrt{13}}\right)\left(\frac{12}{13}\right) - \left(\frac{3}{\sqrt{13}}\right)\left(\frac{5}{13}\right)$ $= \frac{9}{13\sqrt{13}} \times \frac{\sqrt{13}}{\sqrt{13}}$ $= \frac{9\sqrt{13}}{169}$		2	2 marks • Correct solution 1 mark • Recognises that the inverse trig functions represent angles and finds other trig ratios to use in the substitution

Solution		QUESTION 12	Marks	Comments
12(a) $x = 1 + 2t$ $t = \frac{x-1}{2}$ $y = 1 - 4\left(\frac{x-1}{2}\right)^2$ $y = 1 - (x-1)^2$ $y = 1 - x^2 + 2x - 1$ $y = 2x - x^2$			2	2 marks • Correct solution 1 mark • Makes t the subject of either x or y
12(b) (i) # arrangements = $\frac{11!}{3!2!}$ $= 3\,326\,400$			2	2 marks • Correct solution 1 mark • Treats at least one of the repeated letters correctly
12(b) (ii) # arrangements = $2 \times 1 \times \frac{7!}{2!}$ $= 5040$			2	2 marks • Correct solution 1 mark • Correctly arranges the consonants or the vowels
12 (c) Let $t = \tan\theta$, $\cos 2\theta + \sin 2\theta = \frac{1-t^2+2t-(1+t^2)}{1-t^2-2t+(1+t^2)}$ $= \frac{2t-2t^2}{2-2t}$ $= \frac{2t(1-t)}{2(1-t)}$ $= t$ $= \tan\theta$			2	2 marks • Correct solution 1 mark • Rewrites the expression with θ as the angle in all trig expressions
12 (d) $e^{2x} - 4e^x + 3 < 0$ $(e^x - 1)(e^x - 3) < 0$ $1 < e^x < 3$ $\ln 1 < x < \ln 3$ $0 < x < \ln 3$			2	2 marks • Correct solution 1 mark • Finds a solution in terms of e^x
12(e) (i) 			1	1 mark • Correct graph
12(e) (ii) 			2	2 marks • Correct graph 1 mark • Hyperbola drawn for $0 < x \leq 2$ • Horizontal line $y = \frac{1}{2}$ drawn for $2 \leq x \leq 3$

BAULKHAM HILLS HIGH SCHOOL
YEAR 11 EXTENSION YEARLY 2023 SOLUTIONS

Solution		Marks	Comments
SECTION I			
1. A -	$ x-2 \leq 4$ or $x-2 \leq 4$ $x \leq 6$ $-x+2 \leq 4$ $-x \leq 2$ $x \geq -2$ 	1	
2. B -	$-1 \leq \frac{x-a}{b} \leq 1$ $-b \leq x-a \leq b$ $a-b \leq x \leq a+b$	1	
3. D -	when $N = 2N_0$, $2N_0 = N_0 e^{0.04t}$ $e^{0.04t} = 2$ $0.04t = \ln 2$ $t = 25 \ln 2$	1	
4. B -	$\theta = \angle OPB - \angle OPA$ $= \tan^{-1}\left(\frac{4}{x}\right) - \tan^{-1}\left(\frac{3}{x}\right)$	1	
5. D -	$P(x) = ax^3 + bx^2 + 2$ $P'(x) = 3ax^2 + 2bx$ since -1 is a double root, then $P(-1) = P'(-1) = 0$ $a - b + 2 = 0 \Rightarrow 8a - 8b = -16$ (+) $-8a + 7b = 0$ $-b = -16$ $b = 16$ $\therefore a = 14$	1	
6. B -	$\sin^3 x - \sin^3 x \cos x = \sin^3 x \cos x (\cos^2 x - \sin^2 x)$ $= \frac{1}{2} \sin 2x \cos 2x$ $= \frac{1}{4} \sin 4x$	1	
7. C -	$(1+x)^n + (1-x)^n$ $= 1 + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots + x^n + 1 - \binom{n}{1}x + \binom{n}{2}x^2 - \binom{n}{3}x^3 + \dots + (-1)^n x^n$ $= 1 + (-1)^n$	1	
8. A -	$a + \beta + \gamma = \frac{5}{2}$ $a\beta + \alpha\gamma + \beta\gamma = 1$ $a\beta\gamma = \frac{1}{2}$ $\frac{1}{a} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{\beta\gamma + \alpha\gamma + a\beta}{a\beta\gamma}$ $= \frac{1}{\frac{1}{2}} = 2$	1	$\therefore$ it is either A or C However $\frac{1}{a\beta\gamma} = 2$, thus the answer must be A

Solution		Marks	Comments
9. D -	$y = x$ is an odd function $y = \cot x$ is an odd function odd function $\times$ odd function = even function so $y = x \cot x$ is an even function, thus D is the correct graph	1	
10. B -	 The total number of routes would be the same as the total number of routes in the above diagram minus the routes using paths ① or ② # routes = routes from A to B - routes using ① - routes using ② + routes using both ① and ② $= {}^6C_3 - 1 \times 1 \times {}^4C_2 - {}^3C_1 \times 1 \times {}^2C_2 + 1 \times {}^3C_3 \times {}^2C_3$ $= 20 - 6 - 3 + 1$ $= 12$	1	
SECTION II			
QUESTION 11			
11(a)		2	2 marks • Correct solution 1 mark • Correct shape • Correct range
11(b) (i)	$x = \frac{2}{y+1}$ $x(y+1) = 2$ $y+1 = \frac{2}{x}$ $y = \frac{2}{x} - 1$ $y = \frac{2-x}{x}$	1	1 mark • Correct solution
11(b) (ii)	Domain of $f(x)$ is all real x except $x = -1$ $\therefore$ range of $f'(x)$ is all real y except $y = -1$	1	1 mark • Correct answer
11(c) (i)	Teams $= {}^3C_3 \times {}^4C_2$ $= 18$	1	1 mark • Correct answer
11(c) (ii)	Teams $= 1 \times {}^2C_1 \times 1 \times {}^3C_1$ $= 6$	1	1 mark • Correct answer