



PENRITH HIGH SCHOOL

**2014  
HSC TRIAL EXAMINATION**

# Mathematics Extension 2

**General Instructions:**

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen  
Black pen is preferred
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In questions 11 – 16, show relevant mathematical reasoning and/or calculations
- Answer all Questions in the booklets provided

**Total marks–100**

**SECTION I**

**Pages 2–5**

**10 marks**

- Attempt Questions 1–10
- Allow about 15 minutes for this section

**SECTION II**

**Pages 6–11**

**90 marks**

- Attempt Questions 11–16
- Allow about 2 hours 45 minutes for this section

Student Name: \_\_\_\_\_

Teacher Name: \_\_\_\_\_

This paper **MUST NOT** be removed from the examination room

*Assessor: Mr Ferguson*

## Section I

10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section





Use the multiple-choice answer sheet for Questions 1–10.

---

1 What is  $-\sqrt{3} + i$  expressed in modulus-argument form?

- (A)  $\sqrt{2}(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$
- (B)  $2(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$
- (C)  $\sqrt{2}(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6})$
- (D)  $2(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6})$


2 The sketch of the locus of an equation  $|z - 3| + |z + 3| \leq 10$  where  $z = x + iy$  can best be represented by.



- 3 Which of the following expressions is equivalent to  $\int_0^2 \sqrt{4-x^2} dx$ .
- (A)  $\pi$   
 (B)  $2\pi$   
 (C)  $4\pi$   
 (D)  $8\pi$
- 4 Which expression is equal to  $\int \frac{1}{\sqrt{4x^2-8x+5}} dx$ ?
- (A)  $\frac{1}{2} \sin^{-1} 2(x-3) + C$   
 (B)  $\frac{1}{2} \cos^{-1} 2(x-3) + C$   
 (C)  $\frac{1}{2} \ln \left( x-1 + \sqrt{x^2-2x+\frac{5}{4}} \right) + C$   
 (D)  $\frac{1}{2} \ln \left( x-1 + \sqrt{x^2-2x-\frac{5}{4}} \right) + C$
- 5 If  $a, b, c, d$ , and  $e$  are real numbers and  $a \neq 0$ , then the polynomial equation  $ax^7 + bx^5 + cx^3 + dx + e = 0$  has.
- (A) only one real root.  
 (B) at least one real root.  
 (C) an odd number of nonreal roots  
 (D) no real roots
- 6 Suppose that a function  $y = f(x)$  is given with  $f(x) \geq 0$  for  $0 \leq x \leq 4$ . If the area bounded by the curves  $y = f(x)$ ,  $y = 0$ ,  $x = 0$ , and  $x = 4$  is revolved about the line  $y = -1$ , then the volume of the solid of revolution is given by.
- (A)  $\pi \int_0^4 [f(x-1)^2 - 1] dx$   
 (B)  $\pi \int_0^4 [(f(x)-1)^2 - 1] dx$   
 (C)  $\pi \int_0^4 [f(x+1)^2 - 1] dx$   
 (D)  $\pi \int_0^4 [(f(x)+1)^2 - 1] dx$

Use the following information to answer the next question.

A conic is graphed using technology, and is shown below. The distance between the  $x$ -intercepts is  $2\sqrt{11}$  units, and the distance between the  $y$ -intercepts is 10 units.







- 7 The equation of the graph shown above is.

- (A)  $\frac{x^2}{25} + \frac{y^2}{44} = 1$
- (B)  $\frac{x^2}{44} + \frac{y^2}{25} = 1$
- (C)  $\frac{x^2}{25} + \frac{y^2}{11} = 1$
- (D)  $\frac{x^2}{11} + \frac{y^2}{25} = 1$

- 8 If  $3x^2 + 2xy + y^2 = 2$ , then the value of  $\frac{dy}{dx}$  at  $x = 1$  is

- (A)  $-2$
- (B)  $0$
- (C)  $4$
- (D) not defined

9 Which diagram best represents the graph  $y^2 = (x-1)(x-3)^2$  ?



10 A person is standing on the outer edge of a circular disc that is spinning. His relative position on the disc remains unchanged. Which description below best describes the situation?

- (A) The person is experiencing a force that is pushing him away from the centre of the disc.
- (B) The person is experiencing a force that is pushing him towards the centre of the disc.
- (C) The person is experiencing a force tangential to the edge of the disk in the direction of the motion of the disk.
- (D) The person is experiencing a force tangential to the edge of the disk in the direction of the opposite direction to the motion of the disk.

## Section II

**90 marks**

**Attempt Questions 11–16**

**Allow about 2 hours and 45 minutes for this section**

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

In Questions 11–16, your responses should include relevant mathematical reasoning and/or calculations.

---

**Question 11** (15 marks) Use a SEPARATE writing booklet.

a) If  $z_1 = 3 + 4i$ ,  $z_2 = 1 - i$ , find

(i)  $\overline{z_1 z_2}$  1

(ii)  $\left| \frac{z_1}{z_2} \right|$  2

(iii)  $\sqrt{z_1}$  3

b) Sketch separately the following loci in an Argand plane and state the cartesian equations in each case.

(i)  $\operatorname{Re}\left(\frac{z-2}{2}\right) = 0$  2

(ii)  $\arg(z+2) = -\frac{\pi}{6}$  2

c) (i) Express  $\frac{1+2x^2}{(2+x^2)(1+x^2)}$  in the form  $\frac{A}{2+x^2} + \frac{B}{1+x^2}$  2

(ii) Use the substitution  $t = \tan x$  and your answer from part (i) to find  $\int \frac{(1+\sin^2 x) dx}{1+\cos^2 x}$  3

(Leave your answer in term of  $t$ )

**Question 12** (15 marks) Use a SEPARATE writing booklet.

a) If  $\arg z_1 = \theta$  and  $\arg z_2 = \phi$ , show that  $\arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2)$  3

b) The equation  $z^2 + (1+i)z + k = 0$  has root  $1-2i$ . Find the other root, and the value of  $k$ . 2

c) Let  $\alpha, \beta, \gamma$  be the roots (none of which is zero) of  $x^3 + 3px + q = 0$

(i) Find expressions for  $\alpha + \beta + \gamma$ ,  $\alpha\beta + \beta\gamma + \alpha\gamma$  and  $\alpha\beta\gamma$  1

(ii) Find an expression for  $\frac{\alpha\beta}{\gamma} + \frac{\beta\gamma}{\alpha} + \frac{\gamma\alpha}{\beta}$  2


(iii) Find an expression for  $\frac{\alpha\beta}{\gamma} \cdot \frac{\beta\gamma}{\alpha} + \frac{\alpha\beta}{\gamma} \cdot \frac{\gamma\alpha}{\beta} + \frac{\beta\gamma}{\alpha} \cdot \frac{\gamma\alpha}{\beta}$  2

(iv) Hence obtain a monic equation whose roots are  $\frac{\alpha\beta}{\gamma}, \frac{\beta\gamma}{\alpha}, \frac{\gamma\alpha}{\beta}$  2

d) Show that  $\int_0^1 \frac{dx}{9-x^2} = \frac{1}{6} \ln 2$  3

**Question 13** (15 marks) Use a SEPARATE writing booklet.

- a) The shaded region bounded by  $y = 3 - x^2$ ,  $y = x^2 - x$  and  $x = 1$  is rotated about the line  $x = 1$ . The point  $P$  is the intersection of  $y = 3 - x^2$  and  $y = x^2 - x$  in the second quadrant.



- i) Find the  $x$  coordinate of  $P$ . 1
- ii) Use the method of cylindrical shells to express the volume of the resulting solid of revolution as an integral [DO NOT SOLVE THE INTEGRAL] 3

- b) (i) If  $u_n = \frac{1}{n!} \int_0^1 x^n e^{-x} dx$ , where  $n \geq 0$ , show that  $\frac{1}{n!} = e(u_{n-1} - u_n)$  3

- (ii) Hence find the value of  $u_4$  2

- c) If  $a, b, c$  are positive real numbers;
- i) Show that  $a^2 + b^2 \geq 2ab$  1
- ii) Hence prove  $(a + b + c) \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \geq 9$  2

- d) If  $z_1, z_2$  are two complex numbers such that  $|z_1 + z_2| = |z_1 - z_2|$ , show that 3


$$\arg z_1 - \arg z_2 = \frac{\pi}{2}$$

**Question 14** (15 marks) Use a SEPARATE writing booklet.

- a)  $f(x)$  is defined by the equation  $f(x) = x^2 \left( x - \frac{3}{2} \right)$ , on the domain  $-2 \leq x \leq 2$ .

Note: each sketch should take about a third of a page.

- i) Draw a neat sketch of  $f(x)$ , labelling all intersections with coordinate axes and turning points 2
  - ii) Sketch  $y = \frac{1}{f(x)}$  2
  - iii) Sketch  $y = \sqrt{f(x)}$  2
  - iv) Sketch  $y = \ln(f(|x|))$  2
- b) The points  $A, B, C$  and  $D$  lie on the circle  $C$ . From the exterior point  $T$ , a tangent is drawn to point  $A$  on  $C$ . The line  $CT$  passes through  $D$  and  $TC$  is parallel to  $AB$ .



- i) Copy or trace the diagram onto your page.
- ii) Prove that  $\triangle ADT$  is similar to  $\triangle ABC$ . 3

The line  $BA$  is produced through  $A$  to point  $M$ , which lies on a second circle. The points  $A, D, T$  also lie on this second circle and the line  $DM$  crosses  $AT$  at  $O$ .

- iii) Show that  $\triangle OMA$  is isosceles. 2
- iv) Show that  $TM = BC$ . 2

**Question 15** (15 marks) Use a SEPARATE writing booklet.

a) Given that  $\int_0^a f(x) dx = \int_0^a f(a-x) dx$ , find  $\int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx$  2

b) The hyperbola  $H$  has an equation  $xy = 9$ .  $P\left(3p, \frac{3}{p}\right)$ , where  $p > 0$ , and  $Q\left(3q, \frac{3}{q}\right)$ , where  $q > 0$ , are two distinct arbitrary points on  $H$ .

(i) Prove that the equation of the tangent at  $P$  is  $x + p^2y = 6p$  2

(ii) The tangents at  $P$  and  $Q$  intersect at  $T$ . Find the coordinates of  $T$ . 3

(iii) The chord  $PQ$  produced passes through the point  $(0, 6)$ . Given that the equation of this chord is  $x + pqy = 3(p + q)$  find;

(a) Find the equation of the locus of  $T$  3

(b) Give a geometrical description of this locus 1

c) A light inextensible string of length  $3L$  is threaded through a smooth vertical ring which is free to turn. The string carries a particle at each end. One particle  $A$  of mass  $m$  is at rest at a distance  $L$  below the ring. The other particle  $B$  of mass  $M$  is rotating in a horizontal circle whose centre is  $A$ .

(i) Find  $m$  in terms of  $M$ . 2

(ii) Find the angular velocity of  $B$  in terms of  $g$  and  $L$  2

**Question 16** (15 marks) Use a SEPARATE writing booklet.

- a) Use mathematical induction to prove that for all  $n$  where  $n$  can be any positive integer that  $(a-b)$  is a factor of  $a^n - b^n$  **3**
- b) A car travels around a banked circular track of radius 90 metres at 54 km/h.
- (i) Draw a diagram showing all the forces acting on the car **1**
- (ii) Show that the car will have tendency to slip sideways if the angle at which the banked track is banked is  $\tan^{-1}\left(\frac{1}{4}\right)$ . **3**
- (iii) A second car of mass 1.2 tonnes travels around the same bend at 72 km/h. Find the sideways frictional force exerted by the road on the wheels of the car in Newtons. You may assume gravity =  $10 \text{ m/s}^2$ . (Answer correct to 1 decimal place) **3**
- c) (i) Using  $\tan(2\theta + \theta) = \tan 3\theta$ , show that  $\tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}$  **2**
- (ii) Find the value of  $x$  for which  $3 \tan^{-1} x = \frac{\pi}{2} - \tan^{-1} 3x$ , **3**
- where  $\tan^{-1} x$  and  $\tan^{-1} 3x$  both lie between 0 and  $\frac{\pi}{2}$

***End of Exam***

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, n \neq -1; x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, a \neq 0, -a < x < a$$


$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$

NOTE:  $\ln x = \log_e x, x > 0$

# Multiple Choice.

1.  $-\sqrt{3}+i$



$$2 \operatorname{cis} \frac{5\pi}{6}$$

= D.

2. sum of focal lengths is  
a constant  $\Rightarrow 2a$   
 $\therefore a=5$ .

B

3. 
$$= \frac{4\pi}{4}$$

$$= \pi.$$

A.

4. 
$$\int \frac{1}{\sqrt{4(x^2-2x+\frac{5}{2})}} dx$$

$$= \frac{1}{2} \int \frac{1}{\sqrt{x^2-2x+\frac{5}{2}}} dx$$

$$= \frac{1}{2} \int \frac{1}{\sqrt{(x-1)^2 + (\frac{\sqrt{3}}{2})^2}} dx$$



$$= \frac{1}{2} \ln \left[ (x-1) + \sqrt{x^2-2x+\frac{5}{2}} \right] + c$$

= C

5. B

unreal occur in conjugate  
pairs since real coefficient  
so since the degree of polynomial  
is odd there must be at least  
1 real root.

6.



$$\pi \int_0^4 [(y+1)^2 - 1] dy$$

= D.

7. D

8. when  $x=1$ .

D

$$3+2y+y^2=2$$

$$y^2+2y+1=0$$

$$(y+1)^2=0$$

$$\underline{y=-1.}$$

$$\frac{d}{dx} \Rightarrow 6x+y \cdot 2 + 2x \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$$

$$2x \frac{dy}{dx} + 2y \frac{dy}{dx} = -2y - 6x$$

$$\frac{dy}{dx} (2x+2y) = -2y-6x$$

$$\frac{dy}{dx} = \frac{-2y-6x}{2x+2y} \quad \text{when } x=1$$

$$y=-1$$

$$2x+2y=0$$

9. A

10. B.

$\therefore$  not defined

## Question 11

a(i)  $\overline{z_1 z_2}$

$$z_1 z_2 = (3+4i)(1-i)$$

$$= 3 - 3i + 4i - 4i^2$$

$$= 7+i$$

$$\therefore \overline{z_1 z_2} = 7-i$$

(ii)  $\left| \frac{z_1}{z_2} \right|$

$$\frac{z_1}{z_2} = \frac{3+4i}{1-i} \times \frac{1+i}{1+i}$$

$$= \frac{3+3i+4i+4i^2}{1+1}$$

$$= \frac{-1+7i}{2}$$

$$= -\frac{1}{2} + \frac{7}{2}i$$

$$\left| \frac{z_1}{z_2} \right| = \left| -\frac{1}{2} + \frac{7}{2}i \right|$$

$$= \sqrt{\left(-\frac{1}{2}\right)^2 + \left(\frac{7}{2}\right)^2}$$

$$= \sqrt{\frac{1}{4} + \frac{49}{4}}$$

$$= \sqrt{\frac{50}{4}}$$

$$= \frac{5\sqrt{2}}{2}$$

OR  $\frac{|z_1|}{|z_2|} = \frac{\sqrt{9+16}}{\sqrt{1+1}}$

$$= \frac{5}{\sqrt{2}}$$

$$= \frac{5\sqrt{2}}{2}$$

(iii)  $(x+iy)^2 = 3+4i$

$$x^2 + 2xyi - y^2 = 3+4i$$

$$x^2 - y^2 = 3 \quad 2xy = 4$$

$$\therefore xy = 2$$

$$x^2 - \left(\frac{2}{x}\right)^2 = 3$$

$$y = \frac{2}{x}$$

$$x^4 - 3x^2 - 4 = 0$$

$$(x^2+1)(x^2-4) = 0$$

$$x^2 = 4$$

$$x = \pm 2$$

$$\therefore x=2, y=1$$

$$x=-2, y=-1$$

$$\therefore \sqrt{z_1} = \pm(2+i)$$

b

(i)

$$\operatorname{Re}\left(\frac{z-2}{2}\right)=0$$


$$\frac{z-2}{2} = \frac{x+iy-2}{2}$$

$$= \frac{x-2}{2} + \frac{iy}{2}$$

$$\operatorname{Re}\left(\frac{z-2}{2}\right)=0 \Rightarrow \frac{x-2}{2}=0$$

$$\therefore x-2=0$$

$$\text{or } \underline{x=2}$$



$$(ii) \arg(z+2) = -\frac{\pi}{6}$$

$$m = \tan \frac{5\pi}{6} = -\frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = -\frac{\sqrt{3}}{3}$$

$$\text{For y intercept } \tan \frac{\pi}{6} = \frac{y}{2}$$


$$y = 2 \tan \frac{\pi}{6}$$

$$= \frac{2}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$$

$$= \frac{2\sqrt{3}}{3}$$

$$\therefore \text{equation is } y = -\frac{\sqrt{3}}{3}x - \frac{2\sqrt{3}}{3}$$

$$\text{for } x > -2,$$



$$C(i) \quad 1+2x^2 \equiv A(1+x^2) + B(2+x^2)$$

$$\equiv A+Ax^2+2B+Bx^2$$

$$= A+2B+(A+B)x^2$$

$$A+2B=1 \quad \text{--- (1)}$$

$$A+B=2 \quad \text{--- (2)}$$

$$\textcircled{1}-\textcircled{2} \quad B=-1$$

sub into (1)

$$A-2=1$$

$$A=3$$

$$\therefore \frac{3}{2+x^2} + \frac{-1}{1+x^2}$$

$$(ii) \quad \int \frac{(1+\sin^2 x)}{1+\cos^2 x} dx = \int \frac{1 + \frac{1-\cos 2x}{2}}{1 + \frac{1+\cos 2x}{2}} dx$$

$$= \int \frac{3-\cos 2x}{3+\cos 2x} dx$$

$$= \int \frac{3 - \frac{1-t^2}{1+t^2}}{3 + \frac{1-t^2}{1+t^2}} \cdot \frac{1}{1+t^2} dt$$

$$= \int \frac{3+3t^2-1+t^2}{(3+3t^2+1-t^2)(1+t^2)} dt$$

$$= \int \frac{1+2t^2}{(2+t^2)(1+t^2)} dt$$

$$\text{Let } t = \tan x$$

$$\frac{dt}{dx} = \sec^2 x$$

$$dt = \sec^2 x dx$$

$$dt = (1+t^2) dx$$

$$dx = \frac{1}{1+t^2} dt$$

from part (i)

$$= \int \frac{3}{2+t^2} dt - \int \frac{1}{1+t^2} dt$$

$$= \frac{3}{\sqrt{2}} \tan^{-1} \frac{t}{\sqrt{2}} - \tan^{-1} t + C$$

$$= \frac{3}{\sqrt{2}} \tan^{-1} \frac{\tan x}{\sqrt{2}} - x + C$$

### Question 12

a) Let  $z_1 = r_1(\cos \theta + i \sin \theta)$  and  $z_2 = r_2(\cos \phi + i \sin \phi)$

$$\frac{z_1}{z_2} = \frac{r_1(\cos \theta + i \sin \theta)}{r_2(\cos \phi + i \sin \phi)}$$

$$\frac{z_1}{z_2} = \frac{r_1(\cos \theta + i \sin \theta)}{r_2(\cos \phi + i \sin \phi)} \times \frac{(\cos \phi - i \sin \phi)}{\cos \phi - i \sin \phi}$$

$$= \frac{r_1(\cos \theta \cos \phi - i \sin \phi \cos \theta + i \sin \theta \cos \phi - i^2 \sin \theta \sin \phi)}{\cos^2 \phi - i^2 \sin^2 \phi}$$

$$= \frac{r_1[(\cos \theta \cos \phi + \sin \theta \sin \phi) - i(\sin \phi \cos \theta - \sin \theta \cos \phi)]}{r_2 \cdot (\cos^2 \phi + \sin^2 \phi)}$$

$$= \frac{r_1[\cos(\theta - \phi) - i \sin(\theta - \phi)]}{r_2}$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} [\cos(\theta - \phi) + i \sin(\theta - \phi)]$$

$$\arg\left(\frac{z_1}{z_2}\right) = \theta - \phi$$

$$= \arg z_1 - \arg z_2.$$

b) Let  $\alpha, \beta$  be the roots

$$\alpha = 1 - 2i$$

$$\alpha + \beta = -\frac{b}{a}.$$

$$1 - 2i + \beta = -1 - i$$

$$\beta = \underline{\underline{-2 + i}}$$

$$\alpha\beta = \frac{c}{a}$$

$$= k.$$

$$(-2 + i)(1 - 2i) = k$$

$$k = -2 + 4i + i - 2i^2$$

$$= \underline{\underline{5i}}$$

$$c) \quad x^3 + 3px + q = 0$$

$$(i) \quad \alpha + \beta + \gamma = 0$$

$$\alpha\beta + \beta\gamma + \alpha\gamma = 3p$$

$$\alpha\beta\gamma = -q$$

$$\begin{aligned} (ii) \quad \frac{\alpha\beta}{\gamma} + \frac{\beta\gamma}{\alpha} + \frac{\gamma\alpha}{\beta} &= \frac{(\alpha\beta)^2 + (\beta\gamma)^2 + (\gamma\alpha)^2}{\alpha\beta\gamma} \\ &= \frac{(\alpha\beta + \beta\gamma + \gamma\alpha)^2 - 2(\alpha\beta \cdot \beta\gamma + \alpha\beta \cdot \gamma\alpha + \beta\gamma \cdot \gamma\alpha)}{\alpha\beta\gamma} \\ &= \frac{(3p)^2 - 2[\alpha\beta\gamma(\alpha + \beta + \gamma)]}{\alpha\beta\gamma} \\ &= \frac{9p^2 - 0}{-q} \\ &= -\frac{9p^2}{q} \end{aligned}$$

$$\begin{aligned} (iii) \quad \frac{\alpha\beta^2\gamma + \alpha^2\beta\gamma + \alpha\beta\gamma^2}{\alpha\beta\gamma} &= \frac{\alpha\beta\gamma(\alpha^2 + \beta^2 + \gamma^2)}{\alpha\beta\gamma} \\ &= \alpha^2 + \beta^2 + \gamma^2 \\ &= (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma) \\ &= 0 - 2(3p) \\ &= -6p \end{aligned}$$

$$\begin{aligned} (iv) \quad (i) &\Rightarrow \text{Sum of roots} = -\frac{3p^2}{q} \\ (ii) &\Rightarrow \text{sum of root} \\ &\quad 2 \text{ at a time} = -6p \end{aligned}$$

$$\therefore x^3 - \frac{9p^2}{q}x^2 - 6px + q = 0$$

product of roots

$$\begin{aligned} \frac{\alpha\beta}{\gamma} \times \frac{\beta\gamma}{\alpha} \times \frac{\gamma\alpha}{\beta} &= \alpha\beta\gamma \\ &= -q \end{aligned}$$

$$(d) \int_0^1 \frac{dx}{9-x^2} = \frac{1}{6} \ln 2.$$

$$\int_0^1 \frac{1}{(3-x)(3+x)} dx.$$

$$\int_0^1 \frac{1}{(3-x)(3+x)} = \int \left[ \frac{a}{(3-x)} + \frac{b}{(3+x)} \right] dx$$

$$1 \equiv a(3+x) + b(3-x)$$

$$\text{Let } x = 3 \quad 1 = 6a$$

$$\text{Let } x = -3 \quad 1 = 6b$$

$$a = \frac{1}{6} \quad b = \frac{1}{6}$$

$$\int_0^1 \frac{1}{(3-x)(3+x)} = \frac{1}{6} \int_0^1 \frac{1}{3-x} + \frac{1}{6} \int_0^1 \frac{1}{3+x} dx.$$

$$= \left[ -\frac{1}{6} \ln(3-x) \right]_0^1 + \left[ \frac{1}{6} \ln(3+x) \right]_0^1$$

$$= \left[ -\frac{1}{6} \ln 2 + \frac{1}{6} \ln 3 \right] + \left[ \frac{1}{6} \ln 4 - \frac{1}{6} \ln 3 \right]$$

$$= -\frac{1}{6} \ln 2$$

$$= \frac{1}{6} [\ln 3 - \ln 2 + \ln 4 - \ln 3]$$

$$= \frac{1}{6} [\ln 4 - \ln 2]$$

$$= \frac{1}{6} \ln \frac{4}{2}$$

$$= \frac{1}{6} \ln 2.$$


Alternative approach: Let  $x = 3 \sin \theta$   
 $dx = 3 \cos \theta d\theta$

$$\int_0^1 \frac{dx}{9-x^2} = \int_{\sin^{-1}(\frac{1}{3})}^{\sin^{-1}(\frac{1}{3})} \frac{1}{3} \sec \theta d\theta$$

$$= \frac{1}{3} \int \frac{\sec \theta (\tan \theta + \sec \theta)}{\tan \theta + \sec \theta} d\theta$$

$$= \frac{1}{3} [\ln(\tan \theta + \sec \theta)]_{\sin^{-1}(\frac{1}{3})}^{\sin^{-1}(\frac{1}{3})}$$

$$= \frac{1}{3} \left[ \ln \left( \frac{1}{2\sqrt{2}} + \frac{3}{2\sqrt{3}} \right) \right] = \frac{1}{3} \left[ \ln \frac{4}{2\sqrt{2}} \right] = \frac{1}{3} \ln \sqrt{2}$$



### Question 13

a(i) P:  $3 - x^2 = x^2 - x$

$$2x^2 - x - 3 = 0$$



$$(x+1)(2x-3) = 0$$

$$\therefore x = -1, \frac{3}{2}$$

$\therefore x$  coord of P is  $-1$

(as P is in 2nd quadrant)

(ii)



$$h = (3 - x^2) - (x^2 - x)$$

$$= 3 + x - 2x^2$$

Volume  $\int V = \int Ah.$

$$\int V = 2\pi(1-x)(3+x-2x^2) \int x$$

$$V = \lim_{\delta x \rightarrow 0} \sum_{x=-1}^1 2\pi(3-2x-3x^2+2x^3) \delta x$$

$$= 2\pi \int_{-1}^1 3-2x-3x^2+2x^3 dx$$

$$b(i) \quad U_n = \frac{1}{n!} \int_0^1 x^n e^{-x} dx \quad \begin{array}{l} u = x^n \\ \frac{du}{dx} = n x^{n-1} \end{array} \quad \begin{array}{l} \frac{dv}{dx} = e^{-x} \\ v = -e^{-x} \end{array}$$

$$\begin{aligned} U_n &= \frac{1}{n!} \left\{ \left[ -e^{-x} x^n \right]_0^1 - \int_0^1 -e^{-x} n x^{n-1} dx \right\} \\ &= \frac{1}{n!} \left\{ -e^{-1} + n \int_0^1 e^{-x} x^{n-1} dx \right\} \\ &= -\frac{1}{en!} + \frac{1}{(n-1)!} \int_0^1 e^{-x} x^{n-1} dx \end{aligned}$$

$$U_n = -\frac{1}{en!} + U_{n-1}$$

$$\therefore -\frac{1}{en!} = U_{n-1} - U_n$$

$$\frac{1}{n!} = e(U_{n-1} - U_n)$$

$$(ii) \quad U_n = U_{n-1} - \frac{1}{en!}$$

$$U_4 = U_3 - \frac{1}{4!e}$$

$$= U_2 - \frac{1}{3!e} - \frac{1}{4!e}$$

$$\vdots$$

$$= U_0 - \frac{1}{1!e} - \frac{1}{2!e} - \frac{1}{3!e} - \frac{1}{4!e}$$

$$= \frac{1}{0!} \int_0^1 x^0 e^{-x} dx - \frac{1}{e} - \frac{1}{2e} - \frac{1}{6e} - \frac{1}{24e}$$

$$= \int_0^1 e^{-x} dx - \frac{41}{24e}$$

$$= -e^{-1} - (-e^0) - \frac{41}{24e}$$

$$= -\frac{1}{e} + 1 - \frac{41}{24e}$$

$$= -\frac{65}{24e} + 1$$

$$c) (i) (a-b)^2 \geq 0$$

$$a^2 - 2ab + b^2 \geq 0$$

$$a^2 + b^2 \geq 2ab$$

$$(ii) a\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) + b\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) + c\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) \geq 9.$$

$$\text{L.H.S.} \quad 1 + \frac{a}{b} + \frac{a}{c} + 1 + \frac{b}{a} + \frac{b}{c} + 1 + \frac{c}{a} + \frac{c}{b}$$


$$3 + \frac{a}{b} + \frac{b}{a} + \frac{a}{c} + \frac{c}{a} + \frac{b}{c} + \frac{c}{b}$$

$$3 + \frac{a^2+b^2}{ab} + \frac{a^2+c^2}{ac} + \frac{b^2+c^2}{cb}$$

$$\geq 3 + \frac{2ab}{ab} + \frac{2ac}{ac} + \frac{2bc}{cb}$$

$$\geq 9.$$

d)



$$\text{Since } |z_1 + z_2| = |z_1 - z_2|$$

$$|AC| = |OB|$$

OACB is a parallelogram with equal diagonals

$\therefore$  it is a rectangle


$\therefore \angle AOC$  is a right angle

$$\angle xOC - \angle xOA = 90^\circ$$

$$\text{or } \arg z_1 - \arg z_2 = \frac{\pi}{2}.$$

# Question 14

(i)



$$y = x^3 - \frac{3x^2}{2}$$

$$y' = 3x^2 - 3x$$

$$\text{Let } y' = 0$$

$$0 = 3x^2 - 3x$$


$$0 = 3x(x-1)$$

$$x = 0 \text{ or } x = 1$$


$$y = 0 \quad y = 1\left(-\frac{1}{2}\right) = -\frac{1}{2}$$

$$(0, 0) \quad (1, -\frac{1}{2})$$


(ii)




(iii)



(iv)



b(i)



(i) In  $\triangle ADT$  and  $\triangle ABC$

$$\angle TAD = \angle DCA \quad (\text{angle in the alternate segment are equal})$$

$$\angle DCA = \angle CAB \quad (\text{alternate angles on parallel lines DC and AB are equal})$$

$$\therefore \angle TAD = \angle CAB \quad (\text{angle})$$

$$\angle TDA = \angle CBA \quad (\text{exterior angle of a cyclic quadrilateral is equal to the interior opposite angle})$$

$\therefore \triangle ADT \sim \triangle ABC$  (Two angles in one triangle are equal to two angles in the other.)

(ii)

$$\text{Let } \angle TDM = x$$

$$\angle TAM = x \quad (\text{angle in the same segment are equal})$$

$$\angle OMA = x \quad (\text{alternate angles on parallel lines TD and AM are equal})$$

$$\therefore \angle OAM = \angle OMA$$

$\therefore \triangle OMA$  is isosceles.

(iv)

$$\angle TMA = \angle CDA \quad (\text{exterior opposite angles of a cyclic quadrilateral are equal})$$

$$\angle CDA = 180^\circ - \angle CBA \quad (\text{opposite angles of a cyclic quadrilateral are supplementary})$$

$$\therefore \angle CBA = 180^\circ - \angle TMA \quad \therefore CB \parallel TM \quad (\text{since co-interior angles are supplementary})$$

$$\therefore TMBC \text{ is a parallelogram} \quad \therefore TM = CB \quad (\text{opposite sides of parallelogram})$$

### Question 15

$$\begin{aligned}
 (a) \quad \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx &= \int_0^{\pi} \frac{(\pi - x) \sin(\pi - x)}{1 + \cos^2(\pi - x)} dx \\
 &= \int_0^{\pi} \frac{(\pi - x) \sin x}{1 + \cos^2 x} dx \quad \text{since } \sin(\pi - x) = \sin x. \\
 &\quad \text{and } \cos^2(\pi - x) = (-\cos x)^2 = \cos^2 x \\
 &\quad \int_0^{\pi} \frac{\pi \sin x}{1 + \cos^2 x} dx - \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx \\
 \therefore 2 \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx &= \int_0^{\pi} \frac{\pi \sin x}{1 + \cos^2 x} dx \\
 \therefore \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx &= \frac{\pi}{2} \int_0^{\pi} \frac{\sin x}{1 + \cos^2 x} dx \\
 &= -\frac{\pi}{2} \left[ \tan^{-1}(\cos x) \right]_0^{\pi} \quad \text{since } \frac{d}{dx}(\cos x) = -\sin x. \\
 &= -\frac{\pi}{2} \left[ \tan^{-1}(-1) - \tan^{-1}(1) \right] \\
 &= \frac{\pi^2}{4}.
 \end{aligned}$$

$$(b) \quad (i) \quad x = 3p \quad \frac{dx}{dp} = 3 \quad \frac{dy}{dp} = -\frac{3}{p^2}$$

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{dy}{dp} \times \frac{dp}{dx} \\
 &= -\frac{3}{p^2} \times \frac{1}{3} \\
 &= -\frac{1}{p^2}
 \end{aligned}$$

$$\begin{aligned}
 y - \frac{3}{p} &= -\frac{1}{p^2} (x - 3p) \\
 p^2 y - 3p &= -x + 3p \\
 x + p^2 y &= 6p
 \end{aligned}$$

$$(i) \quad x + p^2 y = 6p \quad \text{--- ①}$$

$$x + q^2 y = 6q \quad \text{--- ②}$$

$$\text{①} - \text{②} \quad (p^2 - q^2)y = 6(p - q)$$

$$\frac{(p-q)(p+q)y}{(p-q)} = 6$$

$$(p+q)y = 6$$

$$y = \frac{6}{p+q}$$

sub into ①

$$x + \frac{6p^2}{p+q} = 6p$$

$$x = 6p - \frac{6p^2}{p+q}$$

$$(p+q)x = 6p(p+q) - 6p^2$$

$$(p+q)x = 6p^2 + 6pq - 6p^2$$

$$x = \frac{6pq}{p+q}$$

$$\therefore T\left(\frac{6pq}{p+q}, \frac{6}{p+q}\right)$$

$$(iii) \quad x + pqy = 3(p+q)$$

sub (0, 6)

$$6pq = 3(p+q)$$

$$\frac{pq}{p+q} = \frac{1}{2}$$

$$x = \frac{6pq}{p+q} \quad y = \frac{6}{p+q}$$



$$x = 3 \quad y = \frac{6}{p+q}$$

$$T\left(3, \frac{6}{p+q}\right)$$


$$\therefore x = 3$$

vertical line passing through  $x=3$ .

C,



$$T = mg$$



vertically  $T \sin 30 = Mg$

$$\frac{1}{2}T = Mg$$

$$\text{or } \frac{1}{2}mg = Mg$$

$$m = 2M$$

horizontally

$$T \cos 30 = Mr \omega^2$$

$$\frac{\sqrt{3}}{2}mg = M(\sqrt{3}L)\omega^2$$

$$\frac{\sqrt{3}}{2}mg = \left(\frac{m}{2}\right)(\sqrt{3}L)\omega^2$$

$$\omega = \sqrt{\frac{g}{L}}$$

## Question 1b

(a)

Test for  $n=1$

$a-b$  is a factor of  $a^1-b^1$

Assume true for  $n=k$


$$a^k - b^k = (a-b)F \quad \text{where } F \text{ is an integer}$$

Test for  $n=k+1$



$$\begin{aligned} a^{k+1} - b^{k+1} &= a^{k+1} - a^k b + a^k b - b^{k+1} \\ &= a^k(a-b) + b(a^k - b^k) \\ &= a^k(a-b) + b(a-b)F \quad (\text{from assumption}) \\ &= (a-b)(a^k + bF) \end{aligned}$$

Since  $[a^k + bF]$  is another polynomial in  $a$  and  $b$ , we have shown what we set out to prove.

b)



(i)



$N$  - Normal reaction  
 $F$  - Frictional force  
 $mg$  - weight

(i) Vertically  $N \cos \alpha = F \sin \alpha + mg$

$$N \cos \alpha - F \sin \alpha = mg \quad \text{--- (1)}$$

Horizontally  $N \sin \alpha + F \cos \alpha = \frac{mv^2}{r} \quad \text{--- (2)}$

From (1)  $N \cos \alpha \sin \alpha - F \sin^2 \alpha = mg \sin \alpha \quad \text{--- (3)}$

and (2)  $N \sin \alpha \cos \alpha + F \cos^2 \alpha = \frac{mv^2}{r} \cos \alpha \quad \text{--- (4)}$

(4) - (3)  $F(\cos^2 \alpha + \sin^2 \alpha) = \frac{mv^2}{r} \cos \alpha - mg \sin \alpha$

$$F = \frac{mv^2}{r} \cos \alpha - mg \sin \alpha \quad *$$

no sideways slip when  $F=0$

$$\therefore mg \sin \alpha = \frac{mv^2}{r} \cos \alpha$$

$$\tan \alpha = \frac{v^2}{rg}$$

$$v = \frac{54 \times 1000}{60 \times 60} \text{ ms}^{-1} \quad r = 90.$$

$$\tan \alpha = \frac{15 \times 15}{90 \times 10}$$

$$= \frac{225}{900}$$

$$= \frac{1}{4}$$

$$\alpha = \tan^{-1}\left(\frac{1}{4}\right)$$

(ii) now  $F = m \cos \alpha \left( \frac{v^2}{r} - g \tan \alpha \right) *$

$$F = 1200 \cdot \frac{4}{\sqrt{17}} \left( \left( \frac{12 \times 1000}{60 \times 60} \right)^2 \cdot \frac{1}{90} - \frac{10}{4} \right) \quad \cos \alpha = \frac{4}{\sqrt{17}}$$

$$= \frac{4800}{\sqrt{17}} \left( \frac{400}{90} - \frac{10}{4} \right)$$

$$= \frac{4800}{\sqrt{17}} \left( \frac{800 - 450}{180} \right)$$

$$= \frac{4800}{\sqrt{17}} \times \frac{350}{180}$$

$$\doteq 2263.665 \dots$$

$$= 2263.7 \text{ N}$$

(C).

$$\begin{aligned} \text{(i)} \quad \tan(2Q+Q) &= \frac{\tan 2Q + \tan Q}{1 - \tan 2Q \tan Q} \\ &= \left( \frac{2 \tan Q}{1 - \tan^2 Q} + \tan Q \right) \div \left( 1 - \frac{2 \tan^2 Q}{1 - \tan^2 Q} \right) \\ &= \left( \frac{2 \tan Q + \tan Q - \tan^3 Q}{1 - \tan^2 Q} \right) \div \left( \frac{1 - \tan^2 Q - 2 \tan^2 Q}{1 - \tan^2 Q} \right) \\ &= \frac{3 \tan Q - \tan^3 Q}{1 - 3 \tan^2 Q} \end{aligned}$$

$$\text{(ii)} \quad \text{Let } Q = \tan^{-1} x$$

$$\begin{aligned} \therefore 3Q &= 3 \tan^{-1} x \\ &= \frac{\pi}{2} - \tan^{-1} 3x \end{aligned}$$

$$\tan^{-1} 3x = \frac{\pi}{2} - 3Q$$

$$\tan(\tan^{-1} 3x) = \tan\left(\frac{\pi}{2} - 3Q\right)$$

$$3x = \cot 3Q$$

$$3x = \frac{1 - 3 \tan^2 Q}{3 \tan Q - \tan^3 Q} \quad (\text{from (i)})$$

$$3x = \frac{1 - 3x^2}{3x - x^3} \quad (Q = \tan^{-1} x \Rightarrow x = \tan Q)$$

$$3x(3x - x^3) = 1 - 3x^2$$

$$9x^2 - 3x^4 = 1 - 3x^2$$

$$3x^4 - 12x^2 + 1 = 0$$

$$\therefore x^2 = \frac{6 \pm \sqrt{33}}{3}$$

$$\therefore x = \sqrt{\frac{6 \pm \sqrt{33}}{3}}$$