

**GIRRAWEE HIGH SCHOOL
MATHEMATICS**

Year 12 Extension 2 Task 3

Monday 23rd June 2002

- Instructions: a) Write all your answers on your own paper.
b) Show all necessary working.
c) Marks may be deducted for careless or badly arranged work.

Time Allowed: 80 minutes

Question 1 (27 marks)

Marks

a) Find the following integrals:

- | | |
|---|---|
| (i) $\int \tan^3 x dx$ | 4 |
| (ii) $\int \frac{dx}{x^3 \sqrt{x^2 - 4}}$ | 4 |
| (iii) $\int_0^{\frac{\pi}{2}} \frac{dx}{1 + \sin x + \cos x}$ | 4 |
| (iv) $\int_0^{\frac{\pi}{4}} x \sec^2 x dx$ | 4 |
| (v) $\int \frac{x+2}{\sqrt{x^2 - 6x + 7}} dx$ | 4 |
| (vi) $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^4 x \sin^5 x dx$ | 3 |
| (vii) $\int \frac{7x^2 - 5x + 4}{(x-1)(x^2 + 1)} dx$ | 4 |

Question 2 (9 marks)

a) The integral $I_n = \int_0^{\frac{\pi}{4}} x^n \cos 2x dx$;

(i) Prove that $I_n = \frac{1}{2} \left(\frac{\pi}{4} \right)^n - \frac{n(n-1)}{4} I_{n-2}$ 4


(ii) Using your answer to (i), evaluate $\int_0^{\frac{\pi}{4}} x^4 \cos 2x dx$ 3

b) In your own words explain the error in the following solution: 2


$$\begin{aligned}
 \int \tan x dx &= \int \frac{\sin x dx}{\cos x} \\
 &= \int \sec x \sin x dx \\
 &= -\sec x \cos x + \int \sec x \tan x \cos x dx \\
 &= -\sec x \cos x + \int \tan x dx \\
 \int \tan x dx - \int \tan x dx &= -\sec x \cos x \\
 0 &= -1
 \end{aligned}$$

Question 3 (24 marks)**Marks**

- a) Find the volume generated by rotating the area bounded by $y = \log x$, $y = 0$ and $x = 3$, about the x -axis. 5
- b) Consider the function $y = 2x - x^2$;
- (i) Show that when x is written as the subject, the resulting equation is $x = 1 \pm \sqrt{1 - y}$ 2
- (ii) By taking slices perpendicular to the y -axis, calculate the volume of the solid generated when the area bounded by $y = 2x - x^2$ and the x -axis is rotated about the y -axis. 5
- c) The solid shown has a circular base of radius 2 units. Vertical cross sections perpendicular to the y -axis are equilateral triangles.



- (i) Show that the area of one of these slices is given by $A(y) = \sqrt{3}(4 - y^2)$ 3
- (ii) Hence calculate the volume of the solid. 3
- d) A drinking glass is in the shape of a truncated cone in which the internal diameters of the top and the bottom are 8 cm and 6 cm respectively. The internal height of the glass is 12 cm.




- (i) Show that the cross section taken x cm above the base has radius $\left(3 + \frac{x}{12}\right)$ cm. 2
- (ii) Hence find the volume of fluid the glass can hold. 4

Question 1 (27)

i) $\int \tan^3 x \, dx$
 $= \int \tan x (\sec^2 x - 1) \, dx$
 $= \int \tan x \sec^2 x \, dx - \int \tan x \, dx$
 $= \int u \, du + \log |\cos x| + c \quad u = \tan x$
 $= \frac{1}{2} u^2 + \log |\cos x| + c \quad du = \sec^2 x \, dx$
 $= \frac{1}{2} \tan^2 x + \log |\cos x| + c \quad (4)$

ii) $\int \frac{dx}{x^3 \sqrt{x^2 - 4}}$ $x = 2 \sec \theta$
 $dx = 2 \sec \theta \tan \theta \, d\theta$
 $= \int \frac{2 \sec \theta \tan \theta \, d\theta}{8 \sec^3 \theta \sqrt{4 \sec^2 \theta - 4}}$
 $= \frac{1}{8} \int \frac{\sec \theta \tan \theta \, d\theta}{\sec^3 \theta \tan \theta}$
 $= \frac{1}{8} \int \cos^2 \theta \, d\theta$
 $= \frac{1}{16} \int (1 + \cos 2\theta) \, d\theta$
 $= \frac{1}{16} \theta + \frac{1}{32} \sin 2\theta + c$
 $= \frac{1}{16} \theta + \frac{1}{16} \sin \theta \cos \theta + c$
 $= \frac{1}{16} \sec^{-1} \frac{x}{2} + \frac{1}{16} \cdot \frac{\sqrt{x^2 - 4}}{x} \cdot \frac{2}{x} + c$
 $= \frac{1}{16} \sec^{-1} \frac{x}{2} + \frac{\sqrt{x^2 - 4}}{8x^2} + c \quad (4)$



iii) $\int_{-\pi/2}^{\pi/2} \frac{dx}{1 + \sin x + \cos x}$ $t = \tan \frac{x}{2}$
 $dx = \frac{2dt}{1+t^2}$
 $= \int_{-1}^1 \frac{2dt}{1+t^2}$
 $= \int_{-1}^1 \frac{2dt}{1+t^2 + \frac{1-t^2}{1+t^2}}$
 $= \int_{-1}^1 \frac{2dt}{1+t^2 + 2t + 1 - t^2}$
 $= \int_{-1}^1 \frac{2dt}{2+2t}$
 $= \int_{-1}^1 \frac{dt}{1+t}$
 $= [\log(1+t)]_{-1}^1$
 $= \log 2 - \log 1$
 $= \log 2 \quad (4)$

iv) $\int_{\pi/4}^{\pi/2} x \sec^2 x \, dx$ $u = x$ $v = \tan x$
 $du = dx$ $dv = \sec^2 x \, dx$
 $= [x \tan x]_{\pi/4}^{\pi/2} - \int_{\pi/4}^{\pi/2} \tan x \, dx$
 $= [x \tan x + \log |\cos x|]_{\pi/4}^{\pi/2}$
 $= \frac{\pi}{4} \tan \frac{\pi}{4} + \log \cos \frac{\pi}{4} - 0 + \log \cos 0$
 $= \frac{\pi}{4} + \log \frac{1}{\sqrt{2}} + \log 1$
 $= \frac{\pi}{4} + \log \left(\frac{1}{\sqrt{2}} \right) \quad (4)$

vi) $\int \frac{x+2}{\sqrt{x^2 - 6x + 7}} \, dx$ $u = x^2 - 6x + 7$
 $du = (2x - 6) \, dx$
 $= \frac{1}{2} \int \frac{2x - 6}{\sqrt{x^2 - 6x + 7}} \, dx + 5 \int \frac{dx}{\sqrt{(x-3)^2 - 2}}$
 $= \frac{1}{2} \int u^{-1/2} \, du + 5 \log(x - 3 + \sqrt{x^2 - 6x + 7})$
 $= u^{1/2} + 5 \log(x - 3 + \sqrt{x^2 - 6x + 7}) + c$
 $= \sqrt{x^2 - 6x + 7} + 5 \log(x - 3 + \sqrt{x^2 - 6x + 7})$

vii) $\int_{-\pi/2}^{\pi/2} \cos^4 x \sin^5 x \, dx$
 $\cos^4 x = \text{even function}$
 $\sin^5 x = \text{odd function}$
 $\therefore \cos^4 x \sin^5 x = \text{odd function}$
 $\int_{-\pi/2}^{\pi/2} \cos^4 x \sin^5 x \, dx = 0 \quad (3)$

viii) $\int \frac{7x^2 - 5x + 4}{(x-1)(x^2+1)} \, dx$
 $A(x^2+1) + (Bx+C)(x-1) = 7x^2 - 5x + 4$
 $x=1 \quad x=i$
 $2A = 6 \quad (B+C)(i-1) = -7-5i+4$
 $A = 3 \quad (-B-C) + i(C-B) = -3-5i$
 $B+C = 3$
 $B-C = 5$
 $2B = 8$
 $B = 4 \quad \therefore C = -1$

$\int \frac{7x^2 - 5x + 4}{(x-1)(x^2+1)} \, dx$
 $= \int \left[\frac{3}{x-1} + \frac{4x}{x^2+1} - \frac{1}{x^2+1} \right] \, dx$
 $= 3 \log(x-1) + 2 \log(x^2+1) - \tan^{-1} x + c \quad (4)$

Question 2 (9)

$$a) I_n = \int_0^{\frac{\pi}{2}} x^n \cos 2x \, dx \quad u = x^n \quad v = \frac{1}{2} \sin 2x$$

$$du = nx^{n-1} dx \quad dv = \cos 2x \, dx$$

$$= \left[\frac{1}{2} x^n \sin 2x \right]_0^{\frac{\pi}{2}} - \frac{1}{2} \int_0^{\frac{\pi}{2}} n x^{n-1} \sin 2x \, dx$$

$$u = x^{n-1} \quad v = -\frac{1}{2} \cos 2x$$

$$du = (n-1)x^{n-2} dx \quad dv = \sin 2x \, dx$$

$$= \left[\frac{1}{2} x^n \sin 2x + \frac{1}{4} n x^{n-1} \cos 2x \right]_0^{\frac{\pi}{2}} - \frac{1}{4} n(n-1) \int_0^{\frac{\pi}{2}} x^{n-2} \cos 2x \, dx$$

$$= \frac{1}{2} \left(\frac{\pi}{4} \right)^n \sin \pi + \frac{1}{4} n \left(\frac{\pi}{4} \right)^{n-1} \cos \pi - 0 - \frac{n(n-1)}{4} I_{n-2}$$

$$= \frac{1}{2} \left(\frac{\pi}{4} \right)^n - \frac{n(n-1)}{4} I_{n-2} \quad (4)$$

$$ii) \int_0^{\frac{\pi}{4}} x \cos 2x \, dx$$

$$= \frac{1}{2} \left(\frac{\pi}{4} \right)^2 - \frac{4(3)}{4} I_2$$

$$= \frac{\pi^2}{8} - \frac{3}{2} \left(\frac{\pi}{4} \right)^2 + \frac{2(1)}{4} I_0$$

$$= \frac{5\pi^2}{32} - \frac{3\pi^2}{32} + \frac{1}{4} \int_0^{\frac{\pi}{4}} \cos 2x \, dx$$

$$= \frac{\pi^2}{8} - \frac{3\pi^2}{32} + \frac{1}{4} \left[\sin 2x \right]_0^{\frac{\pi}{4}}$$

$$= \frac{\pi^2}{8} - \frac{3\pi^2}{32} + \frac{1}{4} \left(\sin \frac{\pi}{2} - \sin 0 \right)$$

$$= \frac{\pi^2}{8} - \frac{3\pi^2}{32} + \frac{1}{4} \quad (3)$$


$$b) \int \tan x = -\sec x \cos x + c + \int \tan x \, dx$$

$$\int \tan x - \int \tan x \, dx = -\sec x \cos x + c$$

$$0 = -1 + c$$

ie the error is neglecting the constant of integration. (2)

Question 3 (24)



$$A(y) = 2\pi y (3 - e^y)$$

$$\Delta V = (6\pi y - 2\pi y e^y) \Delta y$$

$$V = \lim_{\Delta y \rightarrow 0} \sum_{y=0}^{\log 3} (6\pi y - 2\pi y e^y) \Delta y$$

$$= 2\pi \int_0^{\log 3} (3y - y e^y) dy \quad u=y \quad v=e^y$$

$$= 2\pi \left[\frac{3}{2} y^2 - y e^y \right]_0^{\log 3} + 2\pi \int_0^{\log 3} e^y dy$$

$$= 2\pi \left[\frac{3}{2} (\log 3)^2 - (\log 3) 3 + 3 - 1 \right]$$

$$= \pi (3(\log 3)^2 - 6 \log 3 + 4) \text{ units}^3$$


$$b) y = 2x - x^2$$

$$x^2 - 2x + y = 0$$

$$x = \frac{2 \pm \sqrt{4 - 4y}}{2}$$

$$x = \frac{2 \pm 2\sqrt{1-y}}{2}$$

$$x = 1 \pm \sqrt{1-y} \quad (2)$$



$$A(y) = \pi (1 + \sqrt{1-y})^2 - \pi (1 - \sqrt{1-y})^2$$

$$= \pi (1 + 2\sqrt{1-y} + (1-y)^2 - 1 + 2\sqrt{1-y} - (1-y)^2)$$

$$= 4\pi \sqrt{1-y}$$

$$\Delta V = 4\pi \sqrt{1-y} \Delta y$$


$$V = \lim_{\Delta y \rightarrow 0} \sum_{y=0}^1 4\pi \sqrt{1-y} \Delta y$$

$$= 4\pi \int_0^1 (1-y)^{\frac{1}{2}} dy$$

$$= 4\pi \left[\frac{2}{3} (1-y)^{\frac{3}{2}} \right]_0^1$$

$$= \frac{8\pi}{3} (0 - 1)$$

$$= \frac{8\pi}{3} \text{ units}^3 \quad (5)$$



$$A(y) = \frac{1}{2} (2x)(\sqrt{3}x)$$

$$= \frac{\sqrt{3}}{2} x^2$$

$$= \frac{\sqrt{3}}{2} (4 - y^2) \quad (3)$$

$$ii) \Delta V = \sqrt{3} (4 - y^2) \Delta y$$


$$V = \lim_{\Delta y \rightarrow 0} \sum_{y=-2}^2 \sqrt{3} (4 - y^2) \Delta y$$

$$= 2\sqrt{3} \int_{-2}^2 (4 - y^2) dy$$

$$= 2\sqrt{3} \left[4y - \frac{1}{3} y^3 \right]_{-2}^2$$

$$= 2\sqrt{3} \left(8 - \frac{8}{3} - 0 \right)$$

$$= \frac{32\sqrt{3}}{3} \text{ units}^3 \quad (3)$$



$$\frac{r-3}{1} = \frac{x}{12}$$

$$r-3 = \frac{x}{12}$$

$$r = 3 + \frac{x}{12} \quad (2)$$



$$A(x) = \pi \left(3 + \frac{x}{12} \right)^2$$

$$= \pi \left(9 + \frac{x}{6} + \frac{x^2}{144} \right)$$

$$\Delta V = \pi \left(9 + \frac{1}{2} x + \frac{1}{144} x^2 \right) \Delta x$$

$$V = \lim_{\Delta x \rightarrow 0} \sum_{x=0}^{12} \pi \left(9 + \frac{1}{2} x + \frac{1}{144} x^2 \right) \Delta x$$

$$= \pi \int_0^{12} \left(9 + \frac{1}{2} x + \frac{1}{144} x^2 \right) dx$$

$$= \pi \left[9x + \frac{1}{4} x^2 + \frac{1}{432} x^3 \right]_0^{12}$$

$$= \pi (108 + 36 + 4 - 0)$$

$$= 148\pi \text{ cm}^3 \quad (4)$$