



KNOX GRAMMAR SCHOOL
MATHEMATICS FACULTY

Set By: EH

Teachers:

EH
RD
AJ

2005
TRIAL HSC EXAMINATION

Mathematics Extension 2 (Year 12)

General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using blue or black pen
- Board-approved calculators may be used
- A table of standard integrals is provided
- All necessary working should be shown in every question

Total marks (120)

- Attempt Questions 1–8
- All questions are of equal value
- Use a **SEPARATE** Writing Booklet for each question
- Write your **Board of Studies Student Number** and **Class Teacher's Initials** on the front cover of each of your writing booklets

Board of Studies Student Number: _____

Class Teacher's Initials: _____

udies
Class
front

udies
Class
front

BLANK PAGE

udies
Class
front

udies
Class
front

Total marks (120)
Attempt questions 1 – 8
All questions are of equal value

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

Question 1 (15 marks)

Marks

(a) Find $\int x e^{-2x} dx$. **2**

(b) Find $\int \sin^3 x dx$. **2**

(c) Evaluate $\int_0^{\frac{\pi}{4}} \frac{d\theta}{4 + 2 \sin 2\theta}$, using the substitution, $t = \tan \theta$. **4**

(d) (i) Find real numbers A , B and C such that: **3**

$$\frac{A}{x-2} + \frac{Bx+C}{x^2+4} \equiv \frac{1}{(x-2)(x^2+4)}$$

(ii) Evaluate $\int_{-2}^0 \frac{8}{(x-2)(x^2+4)} dx$. **4**

Question 2 (15 marks)

Use a SEPARATE writing booklet

Marks

- (a) If $z_1 = 2 + 3i$ and $z_2 = 4 - 5i$, find $z_1 \times \bar{z}_2$, in the form $x + yi$. **2**
- (b) A unit circle has its centre at the origin O . The point z_1 moves on this circle and $z_2 = \frac{\sqrt{2} - 3i}{z_1}$.
- (i) Calculate $|z_1 z_2|$. **1**
- (ii) Hence find the Cartesian equation of the locus of z_2 . **2**
- (c) Sketch the locus of z in the Argand plane such that:
- (i) $|z + 2| = |z - 3i|$ **1**
- (ii) $\arg(z - i) = \frac{3\pi}{4}$ **2**
- (d) (i) Determine the Cartesian equation of the locus of z , such that:
- $$\operatorname{Re}\left(z - \frac{1}{z}\right) = 0$$
- (ii) Hence or otherwise, sketch the locus of z , in the Argand plane, if $\operatorname{Re}\left(z - \frac{1}{z}\right) = 0$. **2**

Question 2 continues on the next page

Question 2 continued:

(e)




Diagram is not to scale

In the diagram, $ABCD$ is a rhombus whose diagonals meet at O , the origin. A represents the complex number $3 + 4i$ and $OD = 2$ units.

Find the complex number represented by:

(i) C


1

(ii) D

2

Question 3 commences on the next page

(a)



For this part of Question 3, your answers are to be superimposed on the appropriate sketches on the separate answer sheet and then handed in with your writing booklet for this question.

The diagram above shows the graph of $y = f(x)$, where $f(x) = \frac{4}{x} - x$.

Sketch on separate number planes the graphs of:

(i) $y = \sqrt{f(x)}$

2

(ii) $y^2 = f(x)$

2

(iii) $y = \frac{1}{f(x)}$

2

(iv) $y = f(|x|)$

2

(v) $y = e^{f(x)}$

2

(b) Evaluate $\int_{-\frac{4}{5}}^{-\frac{2}{5}} \frac{\sqrt{25x^2 - 4}}{x} dx$, using the substitution $x = \frac{2}{5} \sec \theta$.

5

2

Question 4 (15 marks) Use a SEPARATE writing booklet**Marks**

(a) The equation $x^3 - 3x + 2 = 0$ has roots α , β and γ .

(i) Form the cubic polynomial equation with roots:

1

$$\frac{1}{\alpha}, \frac{1}{\beta} \text{ and } \frac{1}{\gamma}$$

(ii) Form the cubic polynomial equation with roots:

3

$$\frac{\alpha}{\beta\gamma}, \frac{\beta}{\alpha\gamma} \text{ and } \frac{\gamma}{\alpha\beta}$$

(b) (i) Prove that if a polynomial $P(x)$ has a root α of multiplicity r then $P'(x)$ has a root α of multiplicity $(r-1)$.

2

(Hint: Start with $P(x) = (x - \alpha)^r Q(x)$)

(ii) Given $x = 1$ is a double root of the equation $x^4 - 5x^3 + 16x^2 - 21x + 9 = 0$, and using the result of b(i), or otherwise, find the other roots.

3

(c) Given $z = \cos\left(\frac{2\pi}{3}\right) + i\sin\left(\frac{2\pi}{3}\right)$ and $w = \sqrt{3}\left(\cos\frac{\pi}{3} + i\sin\frac{\pi}{3}\right)$.

(i) Express $\frac{w}{z}$ in modulus argument form.

1

(ii) Use Mathematical Induction to prove for positive integers n that:


3

$$(\cos\theta + i\sin\theta)^n = \cos n\theta + i\sin n\theta$$

(iii) Hence, using the result in c(ii), find the value of $\left(\frac{w}{z}\right)^{12}$.

2

(a)



The shaded region bounded by the curve $y = \frac{1}{\sqrt{1-x^2}}$, the coordinate axes and the line $x = \frac{1}{2}$ is rotated through one complete revolution about the line $x = 6$.

Use the method of cylindrical shells to find the volume of the solid of revolution formed in cubic units.

(b) Let $I_n = \int_0^{\frac{\pi}{2}} x^n \cos x \, dx$.

(i) Show that if n is a positive integer greater than one, then:


$$I_n = \left(\frac{\pi}{2}\right)^n - n(n-1)I_{n-2}$$

(ii) Evaluate I_4 .

Question 5 continues on the next page

Question 5 continued:

(c)



The base of a certain solid is the area bounded by the parabola $y^2 = 4ax$ (for $y \geq 0, a \geq 0$) and the chord joining $(0, 0)$ and $A(9a, 6a)$.

4

Cross-sections of this solid, determined by planes taken perpendicular to the x -axis, are semicircles with the diameter completely in the base of the solid.

By using the method of slicing, find the total volume of the solid formed.

- (d) (i) Sketch on the same number plane the graphs of $y = |x| - 2$ and $y = 4 + 3x - x^2$.

1

- (ii) Hence or otherwise solve $\frac{|x| - 2}{4 + 3x - x^2} > 0$.

2

Question 6 commences on the next page

(a) Solve for x if $\cos x = \sin\left(\frac{x}{2}\right)$.

3

(b)




Diagram is not to scale

A circular cone of semi-vertical angle $\frac{\pi}{3}$ is fixed with its vertex upwards as shown. A particle P of mass m kg is attached to the vertex at V by a light inextensible string of length 5 metres. The particle P rotates with uniform angular velocity ω rad/sec in a horizontal circle whose centre is vertically below V , on the outside surface of the cone and in contact with it. Let T be the tension in the string and N the normal reaction force at P .

(i) Draw a diagram showing all the forces acting on the particle.

1/1

(ii) Find the tension T in the string and the normal force N on P in Newtons. Leave your answers in terms of m , g and ω .

3

(iii) Show that for the particle to remain in uniform circular motion on the surface of the cone, then $\omega^2 < \frac{2g}{5}$, where g is the acceleration due to gravity.

2

Question 6 continues on the next page

Question 6 continued:

(c)




Diagram is not to scale

For this part of Question 6, your answers are to be placed on the answer sheet supplied and then handed in with your writing booklet for this question.

In the diagram, EC and ED are perpendicular to BA and AC at G and H respectively. The chords AC and BD meet at I . Let $\angle ECA = \alpha$.

- | | | |
|-------|--|----------|
| (i) | Prove that $\triangle BCD$ is isosceles. | 2 |
| (ii) | Prove that $\triangle CID \parallel \triangle CDA$. | 2 |
| (iii) | Given that $\triangle CIB \parallel \triangle CBA$ and $AB + AD = 2BC$. | 2 |

Prove that $CI = \frac{BD}{2}$.

Question 7 commences on the next page

Question 7 (15 marks) Use a SEPARATE writing booklet**Marks**

- (a) A projectile of unit mass is moving through air and experiences a resistance force R proportional to the square of its speed v . That is, $R = kv^2$, where k is a positive constant. *In this question, regard the direction of motion as positive.*

- (i) Suppose the projectile is fired vertically upwards from the ground with an initial speed of u metres per second. Prove that the maximum height H reached by the projectile, where g is the acceleration due to gravity, is given by:

3

$$H = \frac{1}{2k} \log_e \left(1 + \frac{ku^2}{g} \right)$$

marks

- (ii) Prove that the time T taken to reach this maximum height is given by:

2

$$T = \frac{1}{\sqrt{gk}} \tan^{-1} \left(\frac{u\sqrt{k}}{\sqrt{g}} \right)$$

- (b) (i) Differentiate with respect to x the function $h(x) = \frac{\log_{10} x}{x}$.

2

- (ii) Given that the only stationary point of $h(x)$ is a maximum, deduce, without calculating any numerical values, $e^\pi > \pi^e$.

3**Question 7 continues on the next page**

Question 7 continued:

- (c) (i) A vehicle of mass m (in kg) is moving with speed v (in m/s) around a curve of radius r (in metres) banked at angle α with the horizontal. The normal reaction between the road and the vehicle is N , the friction (taken to be up the slope) is F_r , and the acceleration due to gravity is g (in m/s²). **2**

Draw a diagram that represents the forces on the vehicle.

By resolving forces parallel and perpendicular to the road, show that:

$$F_r = mg \sin \alpha - \frac{mv^2}{r} \cos \alpha$$

- (ii) A train is travelling around a curve of radius 3000 metres at a speed of 180 km/h. The width of the rails is 1.5 metres. **3**

Taking the acceleration due to gravity to be 9.8 m/s², find how much higher than the inner rail must the outer rail be, in order for lateral thrust (F_r) on the rails to be avoided? Give your answer to the nearest centimetre.

Question 8 commences on the next page

- (a) Find all the roots of $z^5 - 1 = 0$ and then show that these roots can be represented as $1, \omega, \omega^2, \omega^3$ and ω^4 where $0 < \arg \omega < \frac{\pi}{2}$. 4
- (b) Prove that $(1 - \omega)(1 - \omega^2)(1 - \omega^3)(1 - \omega^4) = 5$. 2
- (c) Show that $(1 - \omega)(1 - \omega^4) = 2 - 2\cos\frac{2\pi}{5}$. 2
- (d) Hence or otherwise, show that $\sin\frac{\pi}{5}\sin\frac{2\pi}{5} = \frac{\sqrt{5}}{4}$. 3
- (e) Suppose P_0, P_1, P_2, P_3, P_4 are the corresponding points of $1, \omega, \omega^2, \omega^3$ and ω^4 in the Argand plane.
- (i) Show that $|\overline{P_0P_1}| = 2\sin\frac{\pi}{5}$. 2
- (ii) Hence, or otherwise deduce that $|\overline{P_0P_1}| \times |\overline{P_0P_2}| \times |\overline{P_0P_3}| \times |\overline{P_0P_4}| = 5$. 2

End of Paper

BLANK PAGE

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$

Note $\ln x = \log_e x, \quad x > 0$

Board of Studies Student Number: _____


Class Teacher's Initials: _____

Knox Grammar School
Year 12
Mathematics Extension 2
TRIAL HSC - 2005
ANSWER SHEET FOR QUESTION 3(a)
(10 Marks)

Hand in with your writing booklet for Question 3

(i) $y = \sqrt{f(x)}$

2




Please turn over ...

ANSWER SHEET FOR QUESTION 3(a)


(ii) $y^2 = f(x)$

2



(iii) $y = \frac{1}{f(x)}$


2



ANSWER SHEET FOR QUESTION 3(a)


(iv) $y = f(|x|)$

2



(v) $y = e^{f(x)}$

2



x

$$y = \frac{1}{2}$$

x

$$y = \frac{1}{2}$$

BLANK PAGE

x

$$y = \frac{1}{2}$$

x


$$y = \frac{1}{2}$$

Board of Studies Student Number: _____

Class Teacher's Initials: _____

Knox Grammar School
Year 12
Mathematics Extension 2
TRIAL HSC - 2005
ANSWER SHEET FOR QUESTION 6(c)
(6 Marks)

Hand in with your writing booklet for Question 6



In the diagram, EC and ED are perpendicular to BA and AC at G and H respectively. The chords AC and BD meet at I . Let $\angle ECA = \alpha$.


- | | | |
|-------|---|---|
| (i) | Prove that $\triangle BCD$ is isosceles. | 2 |
| (ii) | Prove that $\triangle CID \sim \triangle CDA$. | 2 |
| (iii) | Given that $\triangle CIB \sim \triangle CBA$ and $AB + AD = 2BC$. | 2 |

Prove that $CI = \frac{BD}{2}$.

*Q1: RD Q3: EH Q5: AJ Q7: RD
 MARKERS: *Q2: RD Q4: AJ Q6: EH Q8: EH



Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4

Suggested Solutions and Marking Scheme


Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>QUESTION 1: (15 MARKS)</u> (a) $\int x e^{-2x} dx$ using Integration by Parts: let $u = x$ $dv = e^{-2x}$ $du = 1$ $v = -\frac{1}{2}e^{-2x}$ $\therefore \int x e^{-2x} dx$ $= uv - \int v du$ $= -\frac{x}{2}e^{-2x} + \frac{1}{2} \int e^{-2x} dx$ $= -\frac{x}{2}e^{-2x} - \frac{1}{4}e^{-2x} + C$	✓ or implied ✓	$\therefore \int_0^{\frac{\pi}{4}} \frac{d\theta}{4+2\sin 2\theta}$ $= \int_0^{\frac{\pi}{4}} \frac{d\theta}{4+4\sin\theta\cos\theta}$ $= \int_0^1 \frac{\frac{1}{1+t^2}}{4+4\left(\frac{t}{\sqrt{1+t^2}}\right)\left(\frac{1}{\sqrt{1+t^2}}\right)} dt$ $= \frac{1}{4} \int_0^1 \frac{1}{t^2+t+1} dt$ $= \frac{1}{4} \int_0^1 \frac{1}{\left(t+\frac{1}{2}\right)^2+\frac{3}{4}} dt$ $= \frac{1}{2\sqrt{3}} \tan^{-1}\left(\frac{2t+1}{\sqrt{3}}\right) \Big _0^1$ $= \frac{1}{2\sqrt{3}} \left(\tan^{-1}\sqrt{3} - \tan^{-1}\frac{1}{\sqrt{3}} \right)$ $= \frac{\pi\sqrt{3}}{36} \left(\text{or } \frac{\pi}{12\sqrt{3}} \right)$	✓ ✓
(b) $\int \sin^3 x dx$ $= \int \sin^2 x \sin x dx$ $= \int \sin x - \sin x \cos^2 x dx$ $= -\cos x + \frac{\cos^3 x}{3} + C$	✓ method ✓	(d) (i) We want: $A(x^2+4) + (Bx+C)(x-2) \equiv 1$ when $x=2$: $8A=1$ $\therefore A=\frac{1}{8}$ when $x=0$: $4A-2C=1$ $\therefore 2C=-\frac{1}{2}$ $C=-\frac{1}{4}$ when $x=1$: $5A+(B+C)(-1)=1$ $\therefore \frac{5}{8}-B+\frac{1}{4}=1$	✓ ✓
(c) let $t = \tan \theta$  $\therefore \frac{dt}{d\theta} = \sec^2 \theta$ $\therefore \frac{d\theta}{dt} = \frac{1}{\tan^2 \theta + 1}$ $\frac{d\theta}{dt} = \frac{1}{t^2 + 1}$	✓	$\therefore B = -\frac{1}{8}$ $\therefore A = \frac{1}{8}, B = -\frac{1}{8}, C = -\frac{1}{4}$	✓✓✓

when $\theta = 0, t = 0$
 $\theta = \frac{\pi}{4}, t = 1$

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>Q1 ctd:</u> (d)(ii) $\int_{-2}^0 \frac{8}{(x-2)(x^2+4)} dx$ $= \int_{-2}^0 \frac{\frac{1}{8}}{x-2} - \left(\frac{\frac{1}{8}x + \frac{1}{4}}{x^2+4} \right) dx$ $= \int_{-2}^0 \frac{1}{x-2} - \frac{x+2}{x^2+4} dx$ $= \int_{-2}^0 \frac{1}{x-2} - \frac{x}{x^2+4} - \frac{2}{x^2+4} dx$ $= \left[\ln x-2 - \frac{1}{2} \ln(x^2+4) - \tan^{-1} \frac{x}{2} \right]_{-2}^0$ $= \log_e \left \frac{x-2}{\sqrt{x^2+4}} \right - \tan^{-1} \frac{x}{2} \Big _{-2}^0$ $= \ln (-1) - 0 - \ln \left \frac{-4}{2\sqrt{2}} \right - \frac{\pi}{4}$ $= -\ln\sqrt{2} - \frac{\pi}{4}$ $= -\frac{1}{2} \ln 2 - \frac{\pi}{4}$	✓ ✓ ✓ ✓	(b)(i) $z_1 = 1$ (given) $z_2 = \frac{\sqrt{2}-3i}{z_1}$ ($z_1 \neq 0$) $\therefore z_1 z_2 = \sqrt{2}-3i$ <u>we</u> $z_1 z_2 = \sqrt{2}-3i$ $z_1 z_2 = \sqrt{2+9} = \sqrt{11}$ (ii) $\therefore z_2 = \sqrt{11}$, <u>NB</u>: $z_1 =1$ $\therefore$ The locus of z_2 is $x^2+y^2=11$. (c)(i) $z+2 = z-3i$ <u>we</u> $z-(-2) = z-3i$ $\therefore z$ lies on the $\perp$ bisector joining $(-2,0)$ and $(0,3)$ 	✓ ✓ ✓ ✓
<u>QUESTION 2: (15 MARKS)</u> (a) $z_1 = 2+3i$ $z_2 = 4-5i$ $\bar{z}_2 = 4+5i$ $\therefore z_1 \times \bar{z}_2 = (2+3i)(4+5i)$ $= 8+10i+12i-15$ $= -7+22i$	✓ ✓	(ii) This is the half-ray excluding $(0,1)$ making an angle of $\frac{3\pi}{4}$ with the horizontal. $\arg(z-i) = \frac{3\pi}{4}$ 	✓

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>Q2 ctd:</u> (d)(i) There are several ways to do this problem: <u>Method 1:</u> Let $z = r \cos \theta$ $\therefore \frac{1}{z} = \frac{1}{r} \cos(-\theta)$ $\therefore z - \frac{1}{z} = r \cos \theta - \frac{1}{r} \cos(-\theta)$ $= r \cos \theta - \frac{1}{r} \cos \theta + i r \sin \theta + \frac{1}{r} \sin \theta i$ We want $\operatorname{Re}(z - \frac{1}{z}) = 0$ $\therefore r \cos \theta - \frac{1}{r} \cos \theta = 0$ $\therefore \cos \theta (r - \frac{1}{r}) = 0$ $\therefore$ either $\cos \theta = 0, r \neq 0$ $\therefore \theta = \pm \frac{\pi}{2}$ or $r - \frac{1}{r} = 0$ $\therefore r^2 = 1$ $\therefore r = \pm 1$ Since $r = z$ The locus of z is a unit circle, centred at $(0,0)$ and on the y-axis, <u>excluding</u> $(0,0)$ $\therefore x^2 + y^2 = 1 \text{ \& } x=0 \text{ (x,y) \neq (0,0)}$ <u>Method 2:</u> Let $z = x + iy$ $\therefore \operatorname{Re}(x + iy - \frac{1}{x + iy}) = 0$		$\therefore \operatorname{Re}\left[x + iy - \frac{x - iy}{x^2 + y^2}\right] = 0$ $\therefore \operatorname{Re}\left[\frac{x(x^2 + y^2) - x + iy(x^2 + y^2) + y}{x^2 + y^2}\right] = 0$ $\therefore x(x^2 + y^2) - x = 0$ $x(x^2 + y^2 - 1) = 0$ $\therefore x = 0 \text{ or } x^2 + y^2 = 1$ <u>except</u> when $x + iy = 0$. (ii)  (e) (i) $\vec{OC} = c = (-3, -4)$ $\therefore \vec{OC} = -3 - 4i$ (ii) Since the diagonals of a rhombus bisect each other at right-angles, then we first seek a constant k such that: $\vec{OD} = k i \vec{OA}$ $\therefore \vec{OD} = k(3i - 4)$ Since $\vec{OD} = 2$ We want $k(3i - 4) = 2$	✓ ✓ ✓

$$\therefore 9k^2 + 16k^2 = 4$$

$$\therefore k = \frac{2}{5} \text{ (k > 0) - can you see why? } \Rightarrow \vec{OD}$$

$$\therefore \vec{OD} = -\frac{8}{5} + \frac{6}{5}i$$


Board of Studies Student Number: _____

Class Teacher's Initials: _____

Knox Grammar School
Year 12
Mathematics Extension 2
TRIAL HSC - 2005
ANSWER SHEET FOR QUESTION 3(a)
(10 Marks)

Hand in with your writing booklet for Question 3

(i) $y = \sqrt{f(x)}$




Please turn over ...

ANSWER SHEET FOR QUESTION 3(a)


(ii) $y^2 = f(x)$

2




(iii) $y = \frac{1}{f(x)}$

2




ANSWER SHEET FOR QUESTION 3(a)

(iv) $y = f(|x|)$



(v) $y = e^{f(x)}$

2



Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>Q3(b)</u> Let $x = \frac{2}{5} \sec \theta$ $\therefore \frac{dx}{d\theta} = \frac{2}{5} \sec \theta \tan \theta$ when $x = -\frac{4}{5}$, $-2 = \sec \theta$ taking the smallest positive 'argument' $\theta = \frac{2\pi}{3}$. when $x = -\frac{2}{5}$, $-1 = \sec \theta$ as above $\theta = \pi$. $\therefore \int_{-\frac{4}{5}}^{-\frac{2}{5}} \frac{\sqrt{25x^2-4}}{x} \frac{dx}{d\theta} d\theta$ $= \int_{\frac{2\pi}{3}}^{\pi} \frac{\sqrt{25(\frac{4}{25}\sec^2\theta-4)}}{\frac{2}{5}\sec\theta} \cdot \frac{2}{5}\sec\theta\tan\theta d\theta$ ✓ $= \int_{\frac{2\pi}{3}}^{\pi} \frac{2\sqrt{\tan^2\theta}}{\frac{2}{5}\sec\theta} \cdot \frac{2}{5}\tan\theta\sec\theta d\theta$ $= \int_{\frac{2\pi}{3}}^{\pi} 2 \tan\theta \tan\theta d\theta$ since $\tan\theta < 0$ for $\theta \in [\frac{2\pi}{3}, \pi]$ then take $\tan\theta = -\tan\theta$ in this region. $\therefore$ we want: $\int_{\frac{2\pi}{3}}^{\pi} -2\tan^2\theta d\theta$	✓ } ✓ ✓ must have correct 'sign'	$= -2 \int_{\frac{2\pi}{3}}^{\pi} (\sec^2\theta - 1) d\theta$ $= -2 [\tan\theta - \theta]_{\frac{2\pi}{3}}^{\pi}$ $= -2 (\tan\pi - \pi - \tan\frac{2\pi}{3} + \frac{2\pi}{3})$ $= -2 (-\pi + \sqrt{3} + \frac{2\pi}{3})$ $= -2 [\sqrt{3} - \frac{\pi}{3}]$ $= \frac{2\pi}{3} - 2\sqrt{3}$ ✓ NB: $\tan\frac{2\pi}{3} = -\tan\frac{\pi}{3}$	
		<u>QUESTION 4: (15 MARKS)</u> (a) $x^3 - 3x + 2 = 0$ $\alpha, \beta, \gamma \Rightarrow$ roots. (i) let $y = \frac{1}{x}$ $\therefore x = \frac{1}{y}$ $\therefore (\frac{1}{y})^3 - 3(\frac{1}{y}) + 2 = 0$ $\therefore 1 - 3y^2 + 2y^3 = 0$ is the required equation. (ii) Observe that: $\frac{\alpha}{\beta\gamma} = \frac{\alpha^2}{\alpha\beta\gamma} = \frac{\alpha^2}{-2}$ $\frac{\beta}{\alpha\gamma} = \frac{\beta^2}{\alpha\beta\gamma} = \frac{\beta^2}{-2}$ $\frac{\gamma}{\alpha\beta} = \frac{\gamma^2}{\alpha\beta\gamma} = \frac{\gamma^2}{-2}$ $\therefore$ let $y = \frac{x^2}{-2}$ $\therefore x = \sqrt{-2y}$	} ✓


(-1 if (-) sign ignored).

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
Q4a) ctd: $\therefore (\sqrt{-2y})^3 - 3\sqrt{-2y} + 2 = 0$ $\therefore \sqrt{-2y} [(\sqrt{-2y})^2 - 3] = -2$ $\therefore \sqrt{-2y} (-2y - 3) = -2$ $\therefore -2y(2y+3) = 4$ $\therefore -2y(4y^2 + 12y + 9) = 4$ $\therefore -8y^3 - 24y^2 - 18y = 4$ $\therefore 8y^3 + 24y^2 + 18y + 4 = 0$ is the required equation. (b) (i) Let $P(x) = (x-\alpha)^r Q(x)$ $\therefore P'(x) = Q(x)[r(x-\alpha)^{r-1}] + (x-\alpha)^r Q'(x)$ $\therefore P'(x) = (x-\alpha)^{r-1} [rQ(x) + (x-\alpha)Q'(x)]$ $\therefore P'(x)$ has a root $x = \alpha$ of multiplicity $r-1$ since $P(\alpha) = P'(\alpha) = 0$ by substitution. (ii) Let $P(x) = x^4 - 5x^3 + 16x^2 - 21x + 9$ $P'(x) = 4x^3 - 15x^2 + 32x - 21$ we know $P(1) = P'(1) = 0$	✓ ✓	To find the other roots, there are several approaches: NB: $P(x) = (x-1)^2 Q(x)$ by inspection $Q(x) = x^2 - 3x + 9$ ✓ or Divide $(x-1)^2$ into $P(x)$ and use long division to find $Q(x)$ or NB: $1, 1, \alpha, \bar{\alpha}$ are the roots since in $Q(x)$, $\Delta < 0$ $x^2 - 3x + 9 = 0$ $\therefore x = \frac{3 \pm \sqrt{-27}}{2}$ $= \frac{3 \pm 3\sqrt{3}i}{2}$ $\therefore$ The roots are: $1, \frac{3+3\sqrt{3}i}{2}, \frac{3-3\sqrt{3}i}{2}$ (c) (i) $z = \cos \frac{2\pi}{3}$ $w = \sqrt{3} \cos \frac{\pi}{3}$ $\frac{w}{z} = \sqrt{3} \cos \left(\frac{\pi}{3} - \frac{2\pi}{3} \right)$ $\therefore \frac{w}{z} = \sqrt{3} \cos \left(-\frac{\pi}{3} \right)$ (iii) $\left(\frac{w}{z} \right)^{12} = \left[\sqrt{3} \cos \left(-\frac{\pi}{3} \right) \right]^{12}$ $= 3^6 \cos(-4\pi)$ $= 729 \times 1$ $= 729$ ✓	✓ ✓ ✓ ✓

$$\begin{aligned}
 &\downarrow \\
 P'(1) &= 4 - 15 + 32 - 21 \\
 &= 0 \\
 &= P'(1)
 \end{aligned}$$


Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>Q4(c) ctd</u> (ii) Let $S(n)$ represent the statement: $(\cos \theta)^n = \cos(n\theta)$ <u>Step 1</u>: Prove $S(1)$ is true: $\begin{aligned} \text{LHS} &= (\cos \theta)' \\ &= \cos \theta + i \sin \theta \\ \text{RHS} &= \cos(1 \times \theta) \\ &= \cos \theta \\ &= \cos \theta + i \sin \theta \end{aligned}$ $\therefore$ The statement is true for $n=1$. <u>Step 2</u>: Assume $S(k)$ is true for $1 \leq k \leq n, (k, n) \in \mathbb{Z}^+$ <u>Assume</u>: $(\cos \theta)^k = \cos(k\theta)$ <u>RTP</u>: $(\cos \theta)^{k+1} = \cos((k+1)\theta)$ — ① From ① $\begin{aligned} \text{LHS} &= (\cos \theta)^{k+1} \\ &= (\cos \theta)^k \cdot (\cos \theta)' \\ &= (\cos k\theta) \cos \theta \quad \text{from assumption} \\ &= (\cos k\theta + i \sin k\theta)(\cos \theta + i \sin \theta) \\ &= \cos k\theta \cos \theta + i \sin k\theta \cos \theta \\ &\quad + \cos k\theta \cdot i \sin \theta + i^2 \sin k\theta \sin \theta \\ &= \cos k\theta \cos \theta - \sin k\theta \sin \theta \\ &\quad + i(\sin k\theta \cos \theta + \cos k\theta \sin \theta) \\ &= \cos(k\theta + \theta) + i \sin(k\theta + \theta) \\ &= \cos \theta(k+1) + i \sin \theta(k+1) \end{aligned}$	✓	<u>Step 3</u>: The statement is true for $n=1$. Whenever it is true for $n=k$, it is also true for all positive integer values n. <u>QUESTION 5: (15 MARKS)</u> (a)  Slicing parallel to the axis of rotation: The area of a typical slice $A = 2\pi(6-x)y$ Volume of a typical shell $\delta V = 2\pi(6-x)y\delta x$ $\therefore$ Total Volume $V = \lim_{\delta x \rightarrow 0} \sum_{x=0}^{\frac{1}{2}} 2\pi(6-x)y\delta x$ $= 2\pi \int_0^{\frac{1}{2}} \frac{6-x}{\sqrt{1-x^2}} dx$	✓


Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme


Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
Q5(a) ctd: $V = 2\pi \int_0^{\frac{1}{2}} \left(\frac{6}{\sqrt{1-x^2}} - \frac{x}{\sqrt{1-x^2}} \right) dx$ $= 2\pi \left[6\sin^{-1}x + \sqrt{1-x^2} \right]_0^{\frac{1}{2}}$ $= 2\pi \left[6\sin^{-1}\frac{1}{2} + \sqrt{1-\frac{1}{4}} - \sqrt{1} \right]$ $= 2\pi \left[\pi + \frac{\sqrt{3}}{2} - 1 \right]$ $V = \pi [2\pi + \sqrt{3} - 2]$ cubic units. (b) $I_n = \int_0^{\frac{\pi}{2}} x^n \cos x dx$ (i) let $u = x^n$ $dv = \cos x$ $du = nx^{n-1}$ $v = \sin x$ $\therefore I_n = uv - \int v du$ $= x^n \sin x \Big _0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \sin x \cdot nx^{n-1} dx$ $= \left[\left(\frac{\pi}{2} \right)^n \sin \frac{\pi}{2} - 0 \right] - n \int_0^{\frac{\pi}{2}} x^{n-1} \sin x dx$ ↓ Apply integration by parts again $\therefore I_n = \left(\frac{\pi}{2} \right)^n - n \left[x^{n-1} (-\cos x) \Big _0^{\frac{\pi}{2}} + (n-1) \int_0^{\frac{\pi}{2}} x^{n-2} \cos x dx \right]$ $\therefore I_n = \left(\frac{\pi}{2} \right)^n - n(n-1) \int_0^{\frac{\pi}{2}} x^{n-2} \cos x dx$ $I_n = \left(\frac{\pi}{2} \right)^n - n(n-1) I_{n-2}$	✓ ✓ ✓	b(ii) $I_4 = \int_0^{\frac{\pi}{2}} x^4 \cos x dx$ $= \left(\frac{\pi}{2} \right)^4 - 4 \times 3 I_2$ $= \left(\frac{\pi}{2} \right)^4 - 12 \left[\left(\frac{\pi}{2} \right)^2 - 2(1) I_0 \right]$ $= \left(\frac{\pi}{2} \right)^4 - 12 \left[\left(\frac{\pi}{2} \right)^2 - 2 \int_0^{\frac{\pi}{2}} \cos x dx \right]$ $= \frac{\pi^4}{16} - 12 \left[\frac{\pi^2}{4} - [2 \sin x]_0^{\frac{\pi}{2}} \right]$ $= \frac{\pi^4}{16} - 12 \left[\frac{\pi^2}{4} - 2 \sin \frac{\pi}{2} \right]$ $= \frac{\pi^4}{16} - 3\pi^2 + 24$ (c) Length of $PQ = y_2 - y_1$ $\therefore$ Radius of semi-circle, in base is $\frac{ y_2 - y_1 }{2}$ $\therefore$ Area of typical slice is $A = \frac{1}{2} \pi \left(\frac{ y_2 - y_1 }{2} \right)^2$ $\therefore A = \frac{\pi}{8} (y_2 - y_1)^2$	✓ ✓ ✓

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
but $y_1, y_2 > 0$ (as given) $\therefore \text{let } y_2 = \sqrt{4ax}$ $y_1 = \frac{2}{3}x$ } see diagram $\therefore A \approx \frac{\pi}{8} \left[\sqrt{4ax} - \frac{2}{3}x \right]^2$ $= \frac{\pi}{8} \left(2\sqrt{ax} - \frac{2}{3}x \right)^2$ $\therefore$ Volume, δv of a typical slice is: $\delta v = \frac{\pi}{8} \left(2\sqrt{ax} - \frac{2}{3}x \right)^2 \delta x$ $\therefore$ Total Volume of solid: $V = \lim_{\delta x \rightarrow 0} \sum_{x=0}^{x=9a} \frac{\pi}{8} \left(2\sqrt{ax} - \frac{2}{3}x \right)^2 \delta x$ $V = \frac{\pi}{8} \int_0^{9a} \left(2\sqrt{ax} - \frac{2}{3}x \right)^2 dx$ $= \frac{\pi}{8} \left[2ax^{\frac{3}{2}} - \frac{16}{15}x^2\sqrt{ax} + \frac{4x^3}{27} \right]_0^{9a}$ $= \frac{\pi}{8} \left[2a(81a^{\frac{3}{2}}) - \frac{16}{15} \times 81a^2\sqrt{9a} + \frac{4(9a)^3}{27} \right]$ $V = \frac{27\pi a^3}{20}$ cubic units.	✓ ✓ ✓	(d)(i)  $y = x - 2$ $y = 4 + 3x - x^2$ $y = (4-x)(x+1)$ (ii) Solving $\frac{ x -2}{4+3x-x^2} > 0$ We want x such that $x -2$ & $4+3x-x^2$ are the same "sign": When both are (+): $2 < x < 4$ When both are (-): $-2 < x < -1$ $\therefore$ Total solution is: $\{x: -2 < x < -1 \text{ and } 2 < x < 4\}$	✓ ✓ ✓

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>QUESTION 6: (15 MARKS)</u> (a) $\cos x = \sin \frac{x}{2}$ Various methods are available - <u>Method 1:</u> $\cos x = \sin \left(\frac{\pi}{2} - \frac{x}{2} \right)$ $\cos x = \cos \left(\frac{\pi}{2} - \frac{x}{2} \right)$ $\therefore x = 2n\pi \pm \left(\frac{\pi}{2} - \frac{x}{2} \right)$ $\therefore x = 2n\pi + \left(\frac{\pi}{2} - \frac{x}{2} \right) \text{ or } x = 2n\pi - \left(\frac{\pi}{2} - \frac{x}{2} \right)$ $\text{e.g. } \frac{x}{2} = 2n\pi + \frac{\pi}{2} \text{ or } \frac{x}{2} = 2n\pi - \frac{\pi}{2}$ $\text{e.g. } x = \frac{\pi}{3}(4n+1) \text{ or } x = \pi(4n-1)$ where n is an integer. <u>Method 2:</u> $\cos x = \sin \frac{x}{2}$ $\text{e.g. } \cos 2\left(\frac{x}{2}\right) = \sin \frac{x}{2}$ $\text{e.g. } 1 - 2\sin^2 \frac{x}{2} = \sin \frac{x}{2}$ $\text{e.g. } 2\sin^2 \frac{x}{2} + \sin \frac{x}{2} - 1 = 0$ $(2\sin \frac{x}{2} - 1)(\sin \frac{x}{2} + 1) = 0$ $\therefore \sin \frac{x}{2} = \frac{1}{2} \text{ or } \sin \frac{x}{2} = -1$ $\therefore \frac{x}{2} = n\pi + (-1)^n \sin^{-1} \frac{1}{2}$ $\text{e.g. } x = 2n\pi + (-1)^n \frac{\pi}{6}$	Method 1 Method 2 Method 3 Method 4 Method 5 Method 6 Method 7 Method 8 Method 9 Method 10 Method 11 Method 12 Method 13 Method 14 Method 15 Method 16 Method 17 Method 18 Method 19 Method 20 Method 21 Method 22 Method 23 Method 24 Method 25 Method 26 Method 27 Method 28 Method 29 Method 30 Method 31 Method 32 Method 33 Method 34 Method 35 Method 36 Method 37 Method 38 Method 39 Method 40 Method 41 Method 42 Method 43 Method 44 Method 45 Method 46 Method 47 Method 48 Method 49 Method 50 Method 51 Method 52 Method 53 Method 54 Method 55 Method 56 Method 57 Method 58 Method 59 Method 60 Method 61 Method 62 Method 63 Method 64 Method 65 Method 66 Method 67 Method 68 Method 69 Method 70 Method 71 Method 72 Method 73 Method 74 Method 75 Method 76 Method 77 Method 78 Method 79 Method 80 Method 81 Method 82 Method 83 Method 84 Method 85 Method 86 Method 87 Method 88 Method 89 Method 90 Method 91 Method 92 Method 93 Method 94 Method 95 Method 96 Method 97 Method 98 Method 99 Method 100	(b) At P: (i)  $\theta = \frac{\pi}{3}$ (ii) Resolving forces at P: Vertically: $\sum F_y = 0$ $T \cos \theta + N \sin \theta - mg = 0$ $T \times \cos \frac{\pi}{3} + N \sin \frac{\pi}{3} - mg = 0$ $\text{e.g. } \frac{T}{2} + \frac{N\sqrt{3}}{2} - mg = 0 \quad (1)$ $\text{e.g. } T + N\sqrt{3} = 2mg \quad (1)$ Radially: $T \sin \theta - N \cos \theta = mrw^2$ $T \times \sin \frac{\pi}{3} - N \cos \frac{\pi}{3} = mrw^2$ $\frac{T\sqrt{3}}{2} - \frac{N}{2} = mrw^2$	Must have components Method 1 Method 2 Method 3 Method 4 Method 5 Method 6 Method 7 Method 8 Method 9 Method 10 Method 11 Method 12 Method 13 Method 14 Method 15 Method 16 Method 17 Method 18 Method 19 Method 20 Method 21 Method 22 Method 23 Method 24 Method 25 Method 26 Method 27 Method 28 Method 29 Method 30 Method 31 Method 32 Method 33 Method 34 Method 35 Method 36 Method 37 Method 38 Method 39 Method 40 Method 41 Method 42 Method 43 Method 44 Method 45 Method 46 Method 47 Method 48 Method 49 Method 50 Method 51 Method 52 Method 53 Method 54 Method 55 Method 56 Method 57 Method 58 Method 59 Method 60 Method 61 Method 62 Method 63 Method 64 Method 65 Method 66 Method 67 Method 68 Method 69 Method 70 Method 71 Method 72 Method 73 Method 74 Method 75 Method 76 Method 77 Method 78 Method 79 Method 80 Method 81 Method 82 Method 83 Method 84 Method 85 Method 86 Method 87 Method 88 Method 89 Method 90 Method 91 Method 92 Method 93 Method 94 Method 95 Method 96 Method 97 Method 98 Method 99 Method 100




$$\frac{3}{5} = \sin \theta$$

$$\therefore \theta = \sin^{-1} \frac{3}{5}$$

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>Q6 (td):</u> $\frac{T\sqrt{3}}{2} - \frac{N}{2} = m\left(\frac{5\sqrt{3}}{2}\right)w^2$ $\therefore T\sqrt{3} - N = 5\sqrt{3}mw^2$ (2) $\therefore N = T\sqrt{3} - 5\sqrt{3}mw^2$ (3) Sub (3) into (1) $T + (T\sqrt{3} - 5\sqrt{3}mw^2)\sqrt{3} = 2mg$ $\therefore T + 3T - 15mw^2 = 2mg$ $\therefore 4T = 2mg + 15mw^2$ $\therefore T = \frac{m}{4}(2g + 15w^2)$ (4) We want $T > 0, N > 0$ Since $w > 0, m > 0, g > 0$ $T > 0$. $\therefore$ We need $N > 0$. This will occur if: $N = \left[\frac{m}{4}(2g + 15w^2)\right]\sqrt{3} - 5\sqrt{3}mw^2 > 0$ $\therefore$ when: $\frac{m\sqrt{3}}{4}(2g + 15w^2 - 20w^2) > 0$ $\therefore N = \frac{m\sqrt{3}}{4}(2g - 5w^2) > 0$ $\therefore$ when $2g - 5w^2 > 0$ $\therefore w^2 < \frac{2g}{5}$ as expected.	✓ ✓	(C) (i) $\angle AGC = 90^\circ$ $\therefore$ In $\triangle AGC$, $\angle CAG = 90 - \alpha$ (complementary angle) Similarly $\angle CHE = 90^\circ$ $\therefore \angle DEC = 90 - \alpha$ $\therefore \angle BDC = 90 - \alpha$ (A's in the same segment) $\therefore \angle DBC = 90 - \alpha$ (as above) $\therefore \triangle BCD$ is isosceles $\therefore DC = BC$ NB: These are "equal chords" $\therefore \angle DAC = \angle CDB = 90 - \alpha$ (ii)	✓

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
In $\triangle CID$ & $\triangle COA$, 1. $\angle IDC = \angle DAC = 90 - \alpha$ $= \angle BDC$ (equal chords subtend equal angles) 2. $\angle DIC = \angle ADC = 90 + \beta$ $\uparrow$ $(\alpha + \beta + 90 - \alpha)$ $= 90 + \beta$ $\therefore \triangle CID \parallel \triangle CDA$ (A.A.A) (iii)  <u>Given:</u> $\triangle CIB \parallel \triangle CBA$ ① $AB + AD = 2BC$ ② <u>Proven:</u> $\triangle CIB \parallel \triangle CDA$ ③ <u>RTP:</u> $CI = \frac{BD}{2}$. From (ii) $\frac{CI}{CD} = \frac{CD}{CA} = \frac{ID}{DA}$ ④ From (iii) $\frac{CI}{CB} = \frac{CB}{CA} = \frac{IB}{BA}$ ⑤ Corresponding sides in $\parallel \Delta$'s are in proportion.	✓ ✓	<u>Observe:</u> $BD = ID + IB$ (check) From ④ $ID = \frac{IC \times AD}{CD}$ ⑥ From ⑤ $IB = \frac{CI \times AB}{CB}$ ⑦ $\therefore$ ⑥ + ⑦ $ID + IB = \frac{IC \times AD}{CD} + \frac{CI \times AB}{CB}$ $\therefore BD = IC \left[\frac{AD}{CD} + \frac{AB}{CB} \right]$ but $\triangle CDB$ is isosceles $\therefore CD = CB$ $\therefore BD = IC \left[\frac{AD}{CB} + \frac{AB}{CB} \right]$ $BD = \frac{IC}{BC} (AD + AB)$ but $AD + AB = 2BC$ $\therefore BD = \frac{IC}{BC} \times 2BC$ $BD = 2IC$ $\therefore IC = \frac{BD}{2}$ as required.	⑥ ⑦ ✓


Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>QUESTION 7: (15 MARKS)</u> (a) (i) (+) ↑ m=1 $t=0, v=u, x=0$ $m \ddot{x} = \Sigma F$ $\therefore \ddot{x} = -mg - mkv^2$ $\ddot{x} = -g - kv^2$ let $v \frac{dv}{dx} = -(g + kv^2)$ $\therefore \frac{dx}{dv} = -\left(\frac{v}{g + kv^2}\right)$ $\therefore x = -\frac{1}{2k} \log_e(g + kv^2) + C_1$ when $x=0, v=u$ $\therefore C_1 = \frac{1}{2k} \log_e(g + ku^2)$ $\therefore x = \frac{1}{2k} \log_e(g + ku^2) - \frac{1}{2k} \log_e(g + kv^2)$ when $x=H, v=0$ $\therefore H = \frac{1}{2k} \log_e\left(\frac{g + ku^2}{g}\right)$ $\therefore H = \frac{1}{2k} \log_e\left(1 + \frac{ku^2}{g}\right)$ (as expected)	✓ ✓ ✓	<u>NB:</u> we could have obtained the same result / expression if we had evaluated: $\int_0^H dx = \int_u^0 -\left(\frac{v}{g + kv^2}\right) dv$ (Can you see why?) (ii) Let $\frac{dv}{dt} = -g - kv^2$ $\therefore \frac{dv}{dt} = -(g + kv^2)$ $\therefore \frac{dt}{dv} = -\left(\frac{1}{kv^2 + g}\right)$ $\therefore t = -\frac{1}{k} \left[\frac{1}{\sqrt{\frac{g}{k}}} \tan^{-1} \left[\frac{v}{\sqrt{\frac{g}{k}}} \right] \right] + C_2$ when $t=0, v=u$ $C_2 = \frac{1}{k} \left(\frac{1}{\sqrt{\frac{g}{k}}} \right) \tan^{-1} \left[\frac{u\sqrt{k}}{\sqrt{g}} \right]$ $\therefore$ when $t=T, v=0$ $\therefore T = \frac{1}{k} \sqrt{\frac{k}{g}} \left[\tan^{-1} \frac{u\sqrt{k}}{\sqrt{g}} \right]$ $T = \frac{1}{\sqrt{k}g} \tan^{-1} \left(\frac{u\sqrt{k}}{\sqrt{g}} \right)$ $\therefore T = \frac{1}{\sqrt{kg}} \tan^{-1} \left(\frac{u\sqrt{k}}{\sqrt{g}} \right)$ seconds.	✓ ✓ ✓


or: we could have evaluated:

$$\int_0^T dt = -\frac{1}{k} \int_u^0 \frac{1}{v^2 + g/k} dv$$

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
<u>Q7(b)</u> (i) Let $h(x) = \frac{\log_{10} x}{x}$ Changing base: $h(x) = \frac{\log_e x}{\log_e 10} \times \frac{1}{x}$ <u>ie</u> $h(x) = \frac{\ln x}{x \ln 10}$ $\therefore h'(x) = \frac{1}{\ln 10} \left[\frac{x(\frac{1}{x}) - \ln x}{x^2} \right]$ <u>ie</u> $h'(x) = \frac{1}{\ln 10} \left(\frac{1}{x^2} - \frac{\ln x}{x^2} \right)$ (ii) $h(x)$ has SP's when $h'(x) = 0$ <u>ie</u> when $\frac{1}{x^2} - \frac{\ln x}{x^2} = 0$ <u>ie</u> $1 - \ln x = 0$ <u>ie</u> when $x = e$. We know this is the only SP's and the value of x that makes $h(x)$ maximum $\therefore$ for each x in its domain:	✓ ✓ Quotient Rule. ✓	$h(x) < h(e)$ <u>ie</u> $h(\pi) < h(e)$ <i>by substitution</i> $\therefore \frac{1}{\pi} \left(\frac{\ln \pi}{\ln 10} \right) < \frac{1}{e} \left(\frac{\ln e}{\ln 10} \right)$ <u>ie</u> $\frac{\ln e}{e} > \frac{\ln \pi}{\pi}$ $\therefore \pi \ln e > e \ln \pi$ <u>ie</u> $\ln e^\pi > \ln \pi^e$ $\therefore e^\pi > \pi^e$ as required. (c)  (ii) Resolving forces parallel to the slope (and downward) $mg \sin \alpha - F = \frac{mv^2}{r} \cos \alpha$ <u>ie</u> $F = mg \sin \alpha - \frac{mv^2}{r} \cos \alpha$	✓ ✓

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
Q7 (c) (iii) $r = 3000 \text{ m}$ $v = 180 \text{ km/h}$ $= \frac{180 \times 1000}{3600} \text{ m/s}$ $= 50 \text{ m/s}$ Also, $F = 0$ (no lateral thrust) *  $h \leftarrow$ required height. $\sin \alpha = \frac{h}{1.5} *$ but for small α, $\tan \alpha \approx \sin \alpha$. from (ii) $mg \sin \alpha = \frac{mv^2}{r} \cos \alpha$ $\therefore \tan \alpha = \frac{v^2}{rg}$ $\therefore \tan \alpha = \frac{50^2}{3000 \times 9.8}$ $\alpha = 4.9^\circ \dots$ from * $h \doteq 1.5 \sin \alpha$ $\therefore h \approx 1.5 \sin \tan^{-1} \left(\frac{50^2}{3000 \times 9.8} \right)$ $= 0.128 \dots \text{ m}$ $h \doteq 13 \text{ cm}$	✓ ✓	<u>QUESTION 8: (15 MARKS)</u> (a) $z^5 - 1 = 0$ let $z^5 = 1$ $\therefore z = \cos \left(\frac{2k\pi}{5} \right)$ $\therefore$ The roots are: $k=0, z_1 = \cos 0 = 1$ $k=1, z_2 = \cos \left(\frac{2\pi}{5} \right)$ $k=2, z_3 = \cos \left(\frac{4\pi}{5} \right)$ $k=3, z_4 = \cos \left(\frac{6\pi}{5} \right)$ $= \cos \left(\frac{6\pi}{5} - 2\pi \right)$ $= \cos \left(-\frac{4\pi}{5} \right)$ $k=4, z_5 = \cos \left(\frac{8\pi}{5} \right)$ $= \cos \left(-\frac{2\pi}{5} \right)$ If $0 < \arg w < \frac{\pi}{2}$ then select $w = \cos \frac{2\pi}{5}$. $\therefore w^2 = \cos \frac{4\pi}{5} = z_3$ $w^3 = \cos \frac{6\pi}{5} = z_4$ $w^4 = \cos \frac{8\pi}{5} = z_5$ $\therefore 1, w, w^2, w^3, w^4$ are the 5th roots of $z^5 - 1 = 0$.	✓ ✓ ✓ ✓ ✓

Year 12- 2005 Trial HSC Mathematics EXTENSION 2 Assessment Task 4
Suggested Solutions and Marking Scheme

Suggested Solution (s)	Comments	Suggested Solution (s)	Comments
(b) $z^5 - 1$ $= (z-1)(z-w)(z-w^2)(z-w^3)(z-w^4)$ $\quad \quad \quad \underbrace{\hspace{10em}}_{z^4+z^3+z^2+z+1}$ $\therefore (z-w)(z-w^2)(z-w^3)(z-w^4) = z^4 + z^3 + z^2 + z + 1$ let $z=1$. $\therefore (1-w)(1-w^2)(1-w^3)(1-w^4) = 1+1+1+1+1$ $= 5$.	✓ ✓	$\therefore (1-w^2)(1-w^3) = 2 - 2\cos \frac{4\pi}{5}$ ② ✓ ① $\times$ ② $(1-w)(1-w^2)(1-w^3)(1-w^4)$ $= [2 - 2\cos \frac{2\pi}{5}][2 - 2\cos \frac{4\pi}{5}]$ from (b) RHS = 5 ↓	✓
(c) $(1-w)(1-w^4)$ $= 1 - w^4 - w + w^5$ but $w^5 = 1$ $= -w^4 - w + 2$ $= -\cos \frac{8\pi}{5} - \cos \frac{2\pi}{5} + 2$ $= -\cos(-\frac{2\pi}{5}) - \cos(\frac{2\pi}{5}) + 2$ $= -[\cos \frac{4\pi}{5} + i \sin(\frac{2\pi}{5})]$ $\quad - [\cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5}] + 2$ $= -2\cos \frac{2\pi}{5} + 2$ $\therefore (1-w)(1-w^4) = 2 - 2\cos \frac{2\pi}{5}$	✓ ✓	$\therefore (1 - \cos \frac{2\pi}{5})(1 - \cos \frac{4\pi}{5}) = \frac{5}{4}$ $\therefore [2\sin^2 \frac{\pi}{5}][2\sin^2 \frac{2\pi}{5}] = \frac{5}{4}$ [using $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$] $\therefore \sin^2 \frac{\pi}{5} \sin^2 \frac{2\pi}{5} = \frac{5}{16}$ since $\sin \frac{\pi}{5}, \sin \frac{2\pi}{5} > 0$ then $\sin \frac{\pi}{5} \sin \frac{2\pi}{5} = \frac{\sqrt{5}}{4}$	✓
(d) $(1-w)(1-w^4) = 2 - 2\cos \frac{2\pi}{5}$ ① also. Consider: $(1-w^2)(1-w^3) = 1 - w^2 - w^3 + w^5$ $= 2 - \cos \frac{4\pi}{5} - \cos(-\frac{4\pi}{5})$	✓	(e) $P_0 P_1 = w - 1$ $= \cos \frac{2\pi}{5} - 1$ $= \sqrt{\cos^2 \frac{2\pi}{5} - 2\cos \frac{2\pi}{5} + 1 + \sin^2 \frac{2\pi}{5}}$ $= \sqrt{2 - 2\cos \frac{2\pi}{5}}$ $= \sqrt{2} \sqrt{1 - \cos \frac{2\pi}{5}}$ $= \sqrt{2} \sqrt{2\sin^2 \frac{\pi}{5}} = 2\sin \frac{\pi}{5}$	✓ ✓

Q8 (e) ctd.

(ii)

$$\begin{aligned} & |\vec{P_0P_1}| \times |\vec{P_0P_2}| \times |\vec{P_0P_3}| \times |\vec{P_0P_4}| \\ &= |w-1| |w^2-1| |w^3-1| |w^4-1| \checkmark \\ &= |1-w| |1-w^2| |1-w^3| |1-w^4| \\ &= |(1-w)(1-w^2)(1-w^3)(1-w^4)| \checkmark \\ &= |5| \\ &= 5 \quad \text{from (b).} \end{aligned}$$

End of Paper.

Several questions
throughout this paper
can be done in
alternative ways.