


HURLSTONE AGRICULTURAL HIGH SCHOOL



MATHEMATICS

2013

YEAR 12

TRIAL EXAMINATION (TASK 4)

EXAMINERS ~ G RAWSON, P BICZO, J DILLON, S FAULDS, S GUTESA, B MORRISON,

GENERAL INSTRUCTIONS

- | | |
|---|--|
| <ul style="list-style-type: none">• Reading Time – 5 minutes.• Working Time – 3 hours.• Attempt all questions.• This paper contains ten (10) multiple choice questions in Section I and six (6) free response questions in Section II.• Each question in Section II is worth 15 marks.• All necessary working should be shown in every question in Section II.• Board approved calculators and Math aids may be used. | <ul style="list-style-type: none">• Multiple Choice questions to be answered on Multiple Choice answer sheet which may be removed from the back of this question booklet.• Each free response question is to be started in a new answer booklet. Write the question number and your name and/or student number at the top of each answer booklet.• You must hand in the multiple choice answer sheet as well as an answer booklet for each question even if a question has not been attempted.• This examination must NOT be removed from the examination room. |
|---|--|

STUDENT NAME: _____

TEACHER: _____

Section I

10 marks

Attempt Questions 1 - 10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet provided on the back of this booklet for Questions 1-10

1. Given that $ABCD$ is a parallelogram, consider the following statements:

I: If diagonals AC and BD are perpendicular, then $ABCD$ is a rhombus.

II: If diagonals AC and BD are equal, then $ABCD$ is a square.

Which of the statements I and II are correct?

(A) I only


(B) II only

(C) Both I and II


(D) I and II are both incorrect

2. The graph of the function $y = \ln(x + 1)$ is shown in which of the graphs below:


(A)




(B)



(C)



(D)



3. The centre of a circle has co-ordinates $(-3, 4)$. If the circle passes through the point $(1, 2)$, then the **diameter** has length

- (A) $2\sqrt{2}$ units (B) $4\sqrt{2}$ units
(C) $2\sqrt{5}$ units (D) $4\sqrt{5}$ units

4. The solution to the inequality $6 - x - x^2 \leq 0$ is

- (A) $x \leq -3$ or $x \geq 2$ (B) $-3 \leq x \leq 2$
(C) $x \leq -2$ or $x \geq 3$ (D) $-2 \leq x \leq 3$

5. What are the coordinates of the focus of the parabola $x^2 = 2(y - 1)$?

- (A) $\left(0, \frac{1}{2}\right)$ (B) $\left(0, \frac{3}{2}\right)$
(C) $\left(\frac{1}{2}, 0\right)$ (D) $\left(\frac{3}{2}, 0\right)$

6. Which of the following conditions for $\frac{dP}{dt}$ and $\frac{d^2P}{dt^2}$ describe the slowing growth of a variable P ?

- (A) $\frac{dP}{dt} > 0$ and $\frac{d^2P}{dt^2} > 0$ (B) $\frac{dP}{dt} < 0$ and $\frac{d^2P}{dt^2} < 0$
(C) $\frac{dP}{dt} > 0$ and $\frac{d^2P}{dt^2} < 0$ (D) $\frac{dP}{dt} < 0$ and $\frac{d^2P}{dt^2} > 0$

7. The chance of a fisherman catching a legal length fish is 4 in 5. If three fish are caught at random, what is the probability that exactly one is of legal length?

- (A) $\frac{4}{125}$ (B) $\frac{12}{125}$
(C) $\frac{16}{125}$ (D) $\frac{48}{125}$

8. The fourth term of an arithmetic series is 27 and the seventh term is 12. What is the common difference?

(A) -5

(B) 5

(C) 13

(D) 42

9. Solve for $0^\circ \leq x \leq 360^\circ$, $\sin x = \frac{\sqrt{3}}{2}$,

(A) 60° or 240°

(B) 30° or 150°

(C) 30° or 210°

(D) 60° or 120°

10. The curve $y = \sin x$ is rotated about the x -axis between $x = 0$ and $x = \frac{\pi}{4}$.

The volume swept out by this rotation is:

(A) $\pi \int_0^{\frac{\pi}{4}} \sin x . dx$

(B) $\pi \int_0^{\frac{\pi}{4}} (\sin x)^2 . dx$

(C) $\pi \int_0^{\frac{\pi}{4}} \sin x . dy$

(D) $\pi \int_{\frac{\pi}{4}}^0 (1 - \sin x)^2 . dx$

Section II

90 marks

Attempt Questions 11 – 16

Allow about 2 hours and 45 minutes for this section

Answer each question in a separate writing booklet.

All necessary working should be shown in every question.

Question 11 (15 marks) <i>Commence a new answer booklet</i>		Marks
(a)	Simplify $\frac{5}{x-1} - \frac{3}{x+1}$	2
(b)	Solve $ 2x-5 > 3$ and graph the solution on a number line	3
(c)	Determine the value of a and b if $\frac{5}{2+\sqrt{3}} = a + b\sqrt{3}$	2
(d)	Factorise completely: $3x^2 - 12y^2$	2
(e)	Solve simultaneously : $y = x^2$ $y = 2 - x$	3
(f)	Simplify $\log_5 125$	1
(g)	Find the primitive function of $4x^3 - 7x + 5$	2

Question 12 (15 marks) *Commence a new answer booklet*


Marks

- (a) Find the equation of the line perpendicular to the line $y = -2x + 3$ and passing through the point $(3, -4)$. Give your answer in general form. 2
- (b) (i) Find the perpendicular distance of the line l with equation $3x + 4y + 10 = 0$, to the origin $(0, 0)$. 1
- (ii) **Explain** why the line l is a tangent to the circle $x^2 + y^2 = 4$. 1
- (iii) Find the gradient of the line l . 1
- (iv) Calculate to the nearest degree, the angle the line l makes with the positive direction of the x axis. 1
- (v) Sketch the line l , and the circle $x^2 + y^2 = 4$. On your diagram, show the y intercept of the line l , and call this point P . 1
- (vi) Given that the line l touches the circle at $Q\left(-\frac{6}{5}, -\frac{8}{5}\right)$, calculate the area of triangle POQ . 1
- (c) If the quadratic equation $3x^2 - (k + 1)x + 5 = 0$ has roots equal in magnitude but opposite in sign, find the value of k . 1
- (d) Prove that $3x^2 - x + 7$ is positive, for all values of x . You must justify your answer with working. 2
- (e) Find values for a , b and c if $x^2 - x \equiv a(x + 3)^2 + bx + c$. 2
- (f) By reducing the following equation to a quadratic, solve for x :
$$x^4 - x^2 = 2$$
 2

Question 13 (15 marks) *Commence a new answer booklet*

Marks

- (a) Below is the graph of the function $y = f(x)$. **2**
In your answer booklet copy, or trace the diagram.
On your diagram, draw a possible sketch of $y = f'(x)$.



- (b) A function $f(x)$ is defined by $f(x) = x^3 - 3x^2$.
- (i) Find all values of x that satisfy $f(x) = 0$. **1**
 - (ii) Find the coordinates of the stationary points and determine their nature. **3**
 - (iii) Find the coordinates of any points of inflexion, justifying your answer. **2**
 - (iv) Sketch the curve $y = f(x)$ showing all important features. **2**
 - (v) For what values of x is the graph of $y = f(x)$ concave down? **1**
- (c) For the parabola $y = 2x^2 - 8x$.
- (i) Find the coordinates of the vertex. **2**
 - (ii) Find the coordinates of the focus. **1**
 - (iii) Sketch the curve clearly labeling the vertex and focus. **1**

Question 14 (15 marks) *Commence a new answer booklet*

Marks

- (a) Jane's mother puts \$300 into an account at the beginning of each year to pay for Jane's education in 12 years' time. If 6% p.a. interest is paid quarterly, how much money will Jane's mother have at the end of the 12 years? **3**
- (b) In a certain city, the probability that the pollution level will be high is 0.3. If the pollution is monitored for 3 successive days, find the probability that the pollution levels will be:
- (i) High on each of the 3 days **1**
- (ii) High on exactly 2 days **1**
- (c) Each week Lucinda and Terry take part in a raffle at their respective workplaces. The probability that Lucinda wins a prize in her raffle is $\frac{1}{9}$. The probability that Terry wins a prize in his raffle is $\frac{1}{16}$. What is the probability that, during the next three weeks, at least one of them wins a prize? **2**
- (d) The zoom function in a software package multiplies the dimensions of an image by 1.2. In an image, the height of the building is 50 mm. After the zoom function is applied once, the height of the building in the image is 60 mm. After a second application, its height is 72 mm.
- (i) Calculate the height of the building in the image after the zoom function has been applied eight times. Give your answer to the nearest mm. **2**
- (ii) The height of the building in the image is required to be more than 400 mm. Starting from the original image, what is the least number of times the zoom function must be applied? **2**
- (e) Consider the geometric series $5 + 10x + 20x^2 + 40x^3 + \dots$
- (i) For what values of x does this series have a limiting sum? **2**
- (ii) The limiting sum of the series is 100. Find the value of x . **2**

Question 15 (15 marks) *Commence a new answer booklet*

Marks

- (a) By first writing the function $y = 7^x$ in logarithmic form, show that its derivative is $\ln 7 \times 7^x$. 2

- (b) The second derivative of a function is given by $f''(x) = -\frac{3}{x^2}$.
If, $f'\left(\frac{1}{2}\right) = 6$ and $f\left(\frac{1}{2}\right) = 1$, find $f(x)$ in simplest form. 3


- (c) Find the gradient of the curve $y = xe^x$ at the point where $x = \ln 2$.
(Leave your answer in exact form) 2

- (d) (i) Solve the equation $e^{x-1} - 2 = 0$, giving your answer correct to 2 decimal places. 1

- (ii) Evaluate the definite integral $\int_1^2 (e^{x-1} - 2) dx$. (Answer in exact form) 2

- (iii) Explain why your answer in (ii) does not give the area under the curve $y = e^{x-1} - 2$ between the lines $x = 1$ and $x = 2$.
(Note: it is not necessary to calculate the area). 1

(e)



- (i) In the above diagram, $AB \parallel CD$ and BD bisects AC . Prove that $\triangle ABE \equiv \triangle CED$. 2

- (ii) Hence, or otherwise, prove that $AD \parallel BC$. 2

Question 16 (15 marks) *Commence a new answer booklet*

Marks

(a) Express 200° in radians, to 3 decimal places. **1**


(b) Solve $\tan \theta = \frac{1}{\sqrt{2}}$, for $-\pi \leq \theta \leq \pi$, **2**

(c) Two ships leave the Port of Newcastle at 6 am on the 30th of June.
The first ship travels on a bearing of 060° from the port at a speed of 10 km/hr.
The second ship travels on a bearing of 120° from the port at 15 km/hr

(i) Draw a neat diagram to clearly show this information. **1**

(ii) Determine the distance between the ships after 1 hour of sailing. **2**

(d) In the diagram below, $\angle B = 90^\circ$, $\angle A = 60^\circ$ and $AB = AD = 10\text{m}$.
 BD is an arc of the circle with centre A .



Calculate the shaded area in exact form. **3**

(e) Differentiate $x^2 \tan^2(x^2)$ with respect to x . **3**

(f) (i) Differentiate $\ln(\sin x)$ with respect to x **1**

(ii) Hence, find the exact value of $\int_{\frac{\pi}{4}}^{\frac{\pi}{3}} \frac{2}{3} \cot x \, dx$ **2**

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$

This is a blank page

MATHEMATICS
2013
TRIAL EXAMINATION (TASK 4)

PART A ANSWER SHEET

NAME (or number) : _____

Teacher: _____

- Use this page to mark the answers to the questions in Part A.
- Mark the answer by shading in the circle that matches with the correct answer.
- If you make a mistake, draw a cross through the incorrect answer.

YEAR 12

MATHEMATICS

PART A Answers

- | | | | | |
|-----------|-------------------------|-------------------------|-------------------------|-------------------------|
| 1 | ☐ a | ☐ b | ☐ c | ☐ d |
| 2 | ☐ a | ☐ b | ☐ c | ☐ d |
| 3 | ☐ a | ☐ b | ☐ c | ☐ d |
| 4 | ☐ a | ☐ b | ☐ c | ☐ d |
| 5 | ☐ a | ☐ b | ☐ c | ☐ d |
| 6 | ☐ a | ☐ b | ☐ c | ☐ d |
| 7 | ☐ a | ☐ b | ☐ c | ☐ d |
| 8 | ☐ a | ☐ b | ☐ c | ☐ d |
| 9 | ☐ a | ☐ b | ☐ c | ☐ d |
| 10 | ☐ a | ☐ b | ☐ c | ☐ d |

Year 12 Trial Question No.11	Mathematics Solutions and Marking Guidelines	Examination 2013
Outcomes Addressed in this Question		
P3 - performs routine arithmetic and algebraic manipulation involving surds, simple rational expressions and trigonometric identities		
P4 - chooses and applies appropriate arithmetic, algebraic, graphical, trigonometric and geometric techniques		
Outcome	Solutions	Marking Guidelines
P3	(a) $\frac{5}{x-1} - \frac{3}{x+1} = \frac{5(x+1) - 3(x-1)}{(x-1)(x+1)}$ $= \frac{5x + 5 - 3x + 1}{x^2 - 1}$ $= \frac{2x + 8}{x^2 - 1} = \frac{2(x+4)}{(x-1)(x+1)}$	<u>2 marks</u> : correct solution <u>1 mark</u> : substantial progress towards correct solution
	(b) $ 2x - 5 > 3$ $2x - 5 < -3$ or $2x - 5 > 3$ $2x < 2$ $2x > 8$ $x < 1$ $x > 4$ 	<u>3 marks</u> : correct solution <u>2 marks</u> : substantially correct solution <u>1 mark</u> : partially correct solution (eg, finding one side of the inequality)
P3, P4	(c) $\frac{5}{2 + \sqrt{3}} \times \frac{2 - \sqrt{3}}{2 - \sqrt{3}} = \frac{10 - 5\sqrt{3}}{4 - 3}$ $= 10 - 5\sqrt{3}$ $\therefore a = 10, b = -5$	<u>2 marks</u> : correct solution <u>1 mark</u> : substantial progress towards correct solution
	(d) $3x^2 - 12y^2 = 3(x^2 - 4y^2)$ $= 3(x - 2y)(x + 2y)$	<u>2 marks</u> : correct solution <u>1 mark</u> : substantial progress towards correct solution
P4	(e) $y = x^2$... (1) $y = 2 - x$... (2) $x^2 = 2 - x$ (sub (1) $\rightarrow$ (2)) $x^2 + x - 2 = 0$ $(x + 2)(x - 1) = 0$ $x = -1, 2$ subbing into (1) ... when $x = 1$, $y = 1$ when $x = -2$, $y = 4$ $\therefore$ solutions are $(1, 1)$ & $(-2, 4)$	<u>3 marks</u> : correct solution <u>2 marks</u> : substantially correct solution <u>1 mark</u> : partially progress towards correct solution <i>NB: it is important that you explicitly state which x value corresponds to which y value. It is these <u>pairs</u> which are the solutions</i>

P4	<i>Question 11 continued...</i> (f) $\log_5 125 = \log_5 5^3$ $= 3\log_5 5$ $= 3$ (g) $\int (4x^3 - 7x + 5) dx = x^4 - \frac{7x^2}{2} + 5x + C$	1 mark : correct answer 2 marks : correct solution 1 mark : substantial progress towards correct solution (leaving out the C)
------------------	--	---

M/C answers

1. A

2. B

3. D

4. A

5. B

6. C

7. B

8. A


9. D


10. B


Outcomes Addressed in this Question



H5 applies appropriate techniques from the study of calculus, **geometry**, probability, trigonometry and series to solve problems

P4 Chooses and applies appropriate **arithmetic**, algebraic, **graphical**, trigonometric and **geometric techniques**

Outcome	Solutions	Marking Guidelines
P4	a) $y = -2x + 3$ has gradient -2, $\therefore$ line perpendicular has $m = \frac{1}{2}.$ Equation is $y + 4 = \frac{1}{2}(x - 3)$ $2y + 8 = x - 3$ $\therefore x - 2y - 11 = 0$	2 marks : correct solution 1 mark : substantial progress towards correct solution
H5	b) (i) $d = \frac{ 0 + 0 + 10 }{\sqrt{3^2 + 4^2}} = \frac{ 10 }{5} = 2$ units	1 mark: correct answer
H5	(ii) Since the perpendicular distance from the centre of the circle $x^2 + y^2 = 4$ to the line is equal to the radius of the circle, the line is a tangent to the circle.	1 mark: correct explanation
P4	(iii) Rearranging $3x + 4y + 10 = 0$, $y = \frac{-3x - 10}{4}$ $\therefore m = \frac{-3}{4}$	1 mark: correct answer
P4	(iv) Using $m = \tan \theta$, $\tan \theta = \frac{-3}{4}$ Angle of inclination is 143°	1 mark: correct answer
H5	(v) l cuts y axis when $x = 0$. $4y + 10 = 0$ $y = \frac{-5}{2}$ 	1 mark: correct graph, showing y intercept

H5	(vi)  $h^2 = 2.5^2 - 2^2$ $h = 1.5$ $\text{Area of } \Delta = \frac{1}{2} \times 1.5 \times 2 = 1.5 \text{ u}^2$	1 mark : correct answer
H5	(c) Let the roots of $3x^2 - (k+1)x + 5 = 0$ be α and $-\alpha$. Then the sum of the roots = 0. Using sum of roots = $-\frac{b}{a}$, $\frac{k+1}{3} = 0$ $\therefore k = -1.$	1 mark : correct answer
P4	(d) Using $\Delta = b^2 - 4ac$, discriminant of $3x^2 - x + 7$ is $(-1)^2 - 4 \times 3 \times 7 = -83$ $\therefore$ since the discriminant is negative and the coefficient of x^2 is positive (as = 3), $3x^2 - x + 7$ is positive for all values of x. (e) $x^2 - x \equiv a(x+3)^2 + bx + c$ $x^2 - x \equiv ax^2 + 6ax + 9a + bx + c$ $x^2 - x \equiv ax^2 + (6a+b)x + (9a+c)$ Equating like co-efficients $a = 1$ $6a + b = -1 \quad \therefore 6 + b = -1 \quad \therefore 6 + b = -1$ $9a + c = 0 \quad \therefore 9 + c = 0$ $\therefore a = 1, \quad b = -7, \quad c = -9$	2 marks : correct solution 1 mark: significant progress towards correct solution
P4	(f) $x^4 - x^2 = 2$ $(x^2)^2 - x^2 - 2 = 0$ Let $y = x^2$, $y^2 - y - 2 = 0$ $\therefore (y-2)(y+1) = 0$ $\therefore y = -1, 2$ $\therefore x^2 = -1, \quad x^2 = 2$ No solution for $x^2 = -1$ $\therefore x = \pm\sqrt{2}$ $\begin{array}{r l} -2 & 1 \\ & -2 \\ & -1 \end{array}$	2 marks : correct solution 1 mark : substantial progress towards correct solution

Year 12		Mathematics	Task 4 (Trial HSC) 2013
Question No. 13		Solutions and Marking Guidelines	
Outcomes Addressed in this Question			
H6	uses the derivative to determine the features of the graph of a function		
H7	uses the features of a graph to deduce information about the derivative		
P5	understands the concept of a function and the relationship between a function and its graph		
Part	Solutions	Marking Guidelines	
(a) H7		Award 2 for correct graph Award 1 for graph indicating correct location of stationary point (or similar)	
(b) (i) P5	$f(x) = 0$ $x^3 - 3x^2 = 0$ $x^2(x - 3) = 0$ $\therefore x = 0 \text{ or } 3.$	Award 1 for correct solution	
(ii) H6	$f'(x) = 3x^2 - 6x$ $f''(x) = 6x - 6$ Possible stationary points occur @ $f'(x) = 0$ $3x^2 - 6x = 0$ $3x(x - 2) = 0$ $\therefore x = 0 \text{ or } 2$ Test $x = 0$, $f''(0) = -6 < 0$ $\therefore$ Relative maximum @ $(0, 0)$ Test $x = 2$, $f''(2) = 6 > 0$ $\therefore$ Relative minimum @ $(2, -4)$	Award 3 for correct solution Award 2 for substantial progress towards solution Award 1 for limited progress towards solution	
(iii) H6	$f''(x) = 0$ $6x - 6 = 0$ $\therefore x = 1$ Test $x = 0$, $f''(0) = -6$ Test $x = 2$, $f''(2) = 12$ $\therefore$ Change in concavity either side of $x = 1$ $\therefore$ Point of inflexion @ $(1, -2)$	Award 2 for correct solution Award 1 for substantial progress towards solution	

(iv) H6		Award 2 for correct graph Award 1 correct graph, but with insufficient detail
(v) H6	Concave down $f''(x) < 0$ $6x - 6 < 0$ $\therefore x < 1$	Award 1 for correct solution
(c) (i) P5	$y = 2x^2 - 8x$ $= 2(x^2 - 4x)$ $= 2(x^2 - 4x + 4) - 8$ $= 2(x - 2)^2 - 8$ $\therefore 2(x - 2)^2 - 8 = y$ $\therefore 2(x - 2)^2 = y + 8$ $\therefore (x - 2)^2 = \frac{1}{2}(y + 8)$ Vertex = $(2, -8)$	Award 2 for correct solution Award 1 for substantial progress towards solution
(ii) P5	$4a = \frac{1}{2}$ $a = \frac{1}{8}$ $\text{Focus} = \left(2, -7\frac{7}{8} \right)$	Award 1 for correct solution
(iii) P5		Award 1 for correct graph with vertex and focus shown

Year 12 Trial	Mathematics	Examination 2013
Question No. 14	Solutions and Marking Guidelines	
Outcomes Addressed in this Question		
H5 - applies appropriate techniques from the study of calculus, geometry, <i>probability</i> , trigonometry and <i>series</i> to solve problems		
Outcome	Solutions	Marking Guidelines
H5	a) $r = 0.015$ $A_1 = 300(1.015)^4$ $A_2 = (A_1 + 300)(1.015)^4$ $= 300(1.015)^8 + 300(1.015)^4$ $A_3 = (A_2 + 300)(1.015)^4$ $= 300(1.015)^{12} + 300(1.015)^8 + 300(1.015)^4$... $A_{12} = 300(1.015)^{48} + 300(1.015)^{44} + 300(1.015)^{40} + \dots + 300(1.015)^4$ $= 300(1.015)^4 (1 + 1.015^4 + 1.015^8 + \dots + 1.015^{44})$ G.P. with $a=1, r=1.015^4, n=12$ $S_n = \frac{a(r^n - 1)}{r - 1}$ $= 300(1.015)^4 \left(\frac{1(1.015^{48} - 1)}{1.015^4 - 1} \right)$ $= \$5414.50$ (b) $P(\text{high}) = 0.3$ (i) $P(H, H, H) = 0.3^3$ $= 0.027$ (ii) $P(H, H, _) + P(H, _, H) + P(_, H, H)$ $= 3 \times (0.3^2 \times 0.7)$ $= 0.189$ (c) $P(\text{atleast one}) = 1 - P(\text{none win})$ $= 1 - \left(\frac{8}{9} \right)^3 \times \left(\frac{15}{16} \right)^3$ $= \frac{91}{216}$ (d) (i) $60, 72, 86.4, 103.68, \dots$ $T_n = ar^{n-1}$ $T_8 = 60(1.2)^7$ $= 214.99$ $= 215 \text{ mm (nearest mm)}$	(3 marks) correct solution (2 marks) substantial progress towards correct solution (1 mark) some progress towards correct solution (1 mark) correct solution (1 mark) correct solution (2 marks) correct solution (1 mark) some progress towards correct solution (2 marks) correct solution (1 mark) some progress towards correct solution

Year 12 Mathematics Trial (Task 4) Examination 2013		
Question No. 15		Solutions and Marking Guidelines
Outcomes Addressed in this Question		
H2	constructs arguments to prove and justify results	
H3	manipulates algebraic expressions involving logarithmic and exponential functions	
H5	applies appropriate techniques from the study of calculus to solve problems	
H8	uses techniques of integration to calculate areas and volumes	
Outcome	Solutions	Marking Guidelines
H3, H5	(a) $y = 7^x$ $x = \log_7 y \quad (\text{written in equivalent logarithmic form})$ Changing the base to 'e' $x = \frac{\ln y}{\ln 7}$ $\frac{dx}{dy} = \frac{1}{\ln 7} \cdot \frac{1}{y}$ $\frac{dy}{dx} = \frac{1}{\frac{1}{\ln 7} \cdot \frac{1}{y}}$ $= \ln 7 \cdot y$ $= \ln 7 \cdot 7^x \quad (\text{as required})$	2 marks Correct solution showing each step 1 mark Demonstrates ability to express exponential equation in logarithmic form.
H3, H5	(b) $f''(x) = -\frac{3}{x^2}$ $f'(x) = \frac{3}{x} + c$ But $f'\left(\frac{1}{2}\right) = 6$ Substituting gives $6 = \frac{3}{\frac{1}{2}} + c$ $c = 0$ $\therefore f'(x) = \frac{3}{x}$ $f(x) = 3 \ln x + c$ But $f\left(\frac{1}{2}\right) = 1$ Substituting gives $1 = 3 \ln 2^{-1} + c$ $c = 1 + 3 \ln 2$ $\therefore f(x) = 3 \ln x + 3 \ln 2 + 1$ $= 3 \ln 2x + 1$	3 marks Correct solution showing each step 2 mark Substantial progress towards correct solution demonstrating the ability to find a primitive function and evaluate the constant as required. 1 mark Some progress towards correct solution.
H3, H5	(c) $y = xe^x$ $y' = xe^x + e^x$ $= e^x(x + 1)$ Gradient of curve when $x = \ln 2$ $m = e^{\ln 2}(\ln 2 + 1)$ $= 2 \ln 2 + 2$ $= 2(\ln 2 + 1)$	2 marks Correct solution showing each step 1 mark Differentiates the function correctly.
H3	(d) (i) $e^{x-1} - 2 = 0$ $e^{x-1} = 2$ $x - 1 = \ln 2$ $x = \ln 2 + 1$ $= 1.69 \quad (2 \text{ dec. pl.})$	1 mark Correct solution

H3, H5	(ii) $\int_1^2 e^{x-1} - 2 \, dx = [e^{x-1} - 2x]_1^2$ $= (e - 4) - (1 - 2)$ $= e - 4 + 1$ $= e - 3$	2 marks Correct solution showing each step 1 mark Correctly finds the integral but does not proceed to correct solution.
H8	(iii) The curve $y = e^{x-1} - 2$ cuts the x-axis at 1.69. Hence, some of the curve will be above the x-axis, a “positive” area, and some below the x-axis, a “negative” area. Area under the curve should be the magnitude of these areas summed. The integral is the sum of the “signed” area.	1 mark Statement demonstrates a clear understanding of the relationship between an integral and an area.
H2	(e) (i) In Δ's ABE and CED $\angle ABE = \angle CDE$ (alternate angles equal, $AB \parallel CD$) $\angle AEB = \angle CED$ (vertically opposite angles equal) $AE = CE$ (given) $\therefore \Delta ABE \equiv \Delta CED$ (AAS Test)	2 marks Correct solution showing all reasoning. 1 mark Substantial progress towards correct solution or correct solution with deficient reasoning.
H2	(ii) Now, $BE = DE$ (matching sides in congruent triangles) Since $AE = CE$ (given) And $BE = DE$ (proven) Diagonals of quadrilateral $ABCD$ bisect each other, $\therefore ABCD$ is a parallelogram (diagonal properties of a parallelogram.) $\therefore AD \parallel BC$ (opposite sides of a parallelogram)	2 marks Correct solution showing all reasoning. 1 mark Substantial progress towards correct solution or correct solution with deficient reasoning.

Outcomes Addressed in this Question

H3 - manipulates algebraic expressions involving logarithmic and exponential functions
P3 - performs routine arithmetic and algebraic manipulation involving surds, simple rational expressions and trigonometric identities
H5 - applies appropriate techniques from the study of calculus, geometry, probability, trigonometry and series to solve problems
H9 - communicates using mathematical language, notation, diagrams and graphs

Question 16:

H3 - manipulates algebraic expressions involving logarithmic and exponential functions
P3 - performs routine arithmetic and algebraic manipulation involving surds, simple rational expressions and trigonometric identities
H5 - applies appropriate techniques from the study of calculus, geometry, probability, trigonometry and series to solve problems
H9 - communicates using mathematical language, notation, diagrams and graphs

	(a) $\frac{200}{360} \times 2\pi = 3.491$	Correct answer only 1 mark
	(b) Quadrants 1 & 3. Related angle ≈ 0.6155 (retain full value in memory) $\theta = 0.6155$ or -2.5261	Both answers 2 marks
H5	(c) (i) By the cosine rule $AB^2 = 10^2 + 15^2 - 2 \times 10 \times 15 \times \cos(60^\circ)$ (ii) Distance = $5\sqrt{7}$ km ≈ 13.23 km	1 mark for a neat, labelled diagram 2 marks for complete solution
H5	(d) In right triangle ABC $BC = 10\sqrt{3}$ m Area of ABC = $\frac{1}{2} \times 10 \times 10\sqrt{3} \therefore 50\sqrt{3} \text{ m}^2$ Area of sector ABD = $\frac{\pi \times 10^2}{6} \therefore \frac{50\pi}{3} \text{ m}^2$ Exact area = $(50\sqrt{3} - \frac{50\pi}{3}) \text{ m}^2$	1 mark each for the area of the triangle and the sector. 3 marks for the complete solution.

H5	$(e) \frac{d(x^2 \tan^2(x^2))}{dx} = 2x \times \tan^2(x^2) + x^2 \times \frac{d(\tan^2(x^2))}{dx}$ Using the chain rule and letting $u = \tan(v)$ where $v = x^2$ $\frac{d(\tan^2(x^2))}{dx} = \frac{du^2}{du} \times \frac{du}{dv} \times \frac{dv}{dx} = 2u \times \sec^2 v \times 2x$ $= 2 \tan(x^2) \times \sec^2(x^2) \times 2x$ $= 4x \tan(x^2) \sec^2(x^2)$ Hence: $\frac{d(x^2 \tan^2(x^2))}{dx} = 2x \tan^2(x^2) + 4x^3 \tan(x^2) \sec^2(x^2)$ $= 2x \tan(x^2) \{ \tan(x^2) + 2x^2 \sec^2(x^2) \}$	1 mark for the product rule or correct use. 1 mark for substantial progress. 3 marks for the complete solution.
H3	(f) (i) $\frac{d \ln(\sin x)}{dx} = \cot x$ (ii) $\int_{\frac{\pi}{4}}^{\frac{\pi}{3}} \frac{2}{3} \cot x dx = \frac{2}{3} \{ \ln(\sin x) \}_{\frac{\pi}{4}}^{\frac{\pi}{3}}$ $= \frac{2}{3} \{ \ln \frac{\sqrt{3}}{2} - \ln \frac{1}{\sqrt{2}} \}$ $= \frac{2}{3} \{ \ln \frac{\sqrt{3}}{2} + \ln \sqrt{2} \}$ $= \frac{2}{3} \ln \frac{\sqrt{6}}{2}$ ← acceptable progress for full marks $= \frac{2}{3} \{ \ln 6^{\frac{1}{2}} - \ln 2 \}$ $= \frac{2}{3} \times \frac{1}{2} \ln 6 - \frac{2}{3} \ln 2$ $= \frac{\ln 6 - 2 \ln 2}{3} \quad (2 \ln 2 = \ln 4)$ $= \frac{1}{3} \ln \left(\frac{3}{2} \right)$	1 mark 1 mark for correct integration and 1 mark for substantial work towards the complete solution 2 marks for the complete solution.

Note: Part (e) was, in some cases, expressed as

$2x(2x^2 + 1) \tan^2(x^2) + 4x^3 \tan(x^2)$, after having replaced $\sec^2(x^2)$ by $1 + \tan^2(x^2)$ and collecting the $\tan^2(x^2)$