



--	--	--	--	--	--	--	--	--	--

Student Number

2020

Trial Higher School Examination

Mathematics Advanced

General Instructions

- Reading time – 10 minutes
- Working time – 3 hours
- Write using black pen
- Calculators approved by NESA may be used
- A reference sheet is provided at the back of this paper
- For questions in Section II, show relevant mathematical reasoning and/or calculations

Total marks: 100

Section I – 10 marks (pages 2-4)

- Attempt Questions 1-10
- Allow about 15 minutes for this section

Section II – 90 marks (pages 5-24)

- Attempt Questions 11-32
- Allow about 2 hours and 45 minutes for this section

Section I**10 marks****Attempt questions 1 - 10****Allow about 15 minutes for this section**

Use the multiple-choice answer sheet for questions 1-10

1. What is the value of $\operatorname{cosec} \frac{\pi}{3}$ to three significant figures?
(A) 1.00
(B) 1.15
(C) 1.41
(D) 2.00

2. What is the value of c for which the circle $(x-3)^2 + (y-2)^2 = c$ touches the x axis?
(A) 2
(B) 3
(C) 4
(D) 9

3. What is the equation of the tangent to $y = x^2 - 3$ at $x = -1$?
(A) $y = -2x - 4$
(B) $y = 2x - 4$
(C) $y = \frac{x}{2} - \frac{3}{2}$
(D) $y = -\frac{x}{2} - \frac{3}{2}$

4. Which statement is true for an ungrouped data set with no outliers?
- (A) The largest possible range is 2 times the interquartile range.
 - (B) The largest possible range is 3 times the interquartile range.
 - (C) The largest possible range is 4 times the interquartile range.
 - (D) The largest possible range is 5 times the interquartile range.
5. Which one of the following is the set of all solutions to $2x^2 - 5x + 2 \geq 0$?
- (A) $\left[\frac{1}{2}, 2\right]$
 - (B) $\left(\frac{1}{2}, 2\right)$
 - (C) $\left(-\infty, \frac{1}{2}\right) \cup (2, \infty)$
 - (D) $\left(-\infty, \frac{1}{2}\right] \cup [2, \infty)$
6. The graph of $y = f(x)$ has a stationary point at $(2, -3)$.
- Which one of the following is a guaranteed stationary point of $y = -f\left(\frac{x}{2}\right) - 5$?
- (A) $(1, -2)$
 - (B) $(1, 2)$
 - (C) $(4, -2)$
 - (D) $(4, 2)$

7. What is the period and amplitude for the curve $y = \sin \pi x$?
- (A) Amplitude = 1; Period = 2
 (B) Amplitude = π ; Period = 2
 (C) Amplitude = 1; Period = 2π
 (D) Amplitude = π ; Period = 2π
8. If the z scores on an examination are normally distributed and $P(z < N) = 0.6$ for some number N , what is the value of $P(-N < z < N)$?
- (A) 0.1
 (B) 0.2
 (C) 0.3
 (D) 0.4
9. Which one of the following equations is NOT correct?
- (A) $\int x(x^2 - 1)^2 dx = \frac{(x^2 - 1)^3}{6} + c$
 (B) $\int_{-3}^3 \sqrt{9 - x^2} dx = \frac{9\pi}{2}$
 (C) $\int_{-1}^1 3^x dx = \frac{1}{\ln 3} \left(3 - \frac{1}{3} \right)$
 (D) $\int_{-5}^5 4x^4 - x^3 + \cos x dx = 0$
10. Consider the series $\sqrt{5} + \sqrt{45} + \sqrt{125} + \dots + z = 225\sqrt{5}$, the value of z is:
- (A) $25\sqrt{5}$
 (B) $29\sqrt{5}$
 (C) $30\sqrt{5}$
 (D) $35\sqrt{5}$

Section II**90 marks****Attempt all questions****Allow about 2 hours and 45 minutes for this section**

Answer each question in the spaces provided.

Your responses should include relevant mathematical reasoning and/or calculations.



Extra writing space is provided at the back of the examination paper.

Question 11 (2 marks)**Marks**

What angle does the line $2x + 3y + 6 = 0$ make with the positive x -axis? Round to the nearest minute.

2**Question 12 (2 marks)**


Sketch a possible function which could have the gradient function as graphed below.

**2**

Question 13 (3 marks)

Marks

In triangle ABC , the length of $AB=5\text{cm}$, $AC=13\text{cm}$ and $\cos\angle BAC = \frac{1}{8}$



(a) Find the exact value of $\sin\angle BAC$

2

(b) Find the area of triangle ABC

1

Question 14 (3 marks)

Solve $2\log x = \log(5x + 6)$

3

Question 15 (3 marks)

Marks

Solve $|1 - 2\cos^2 x| = 1$ for $0 \leq x \leq 2\pi$

3

Question 16 (5 marks)

Differentiate the following expressions.

Marks

(a) $\log_5(\tan x)$


2

(b) $\frac{2^x}{e^x}$

3

Question 17 (5 marks)**Marks**

Consider the graph $y = f(x)$. Both arcs have a radius of four units.



Using the graph of $y = f(x)$, $x \geq 0$, evaluate exactly the following integrals.

(a) $\int_0^{12} f(x) dx$

3

(b) $\int_0^{18} f(x) dx$

2

Question 18 (5 marks)

Marks

The discrete random variable X has probability distribution shown in the table below

x	-1	0	1	2	3
$P(X=x)$	a	b	0.2	0.15	0.13

and $E(X) = 0.55$

- (a) By forming a pair of simultaneous equations, or otherwise, find the values of a and b . **3**

This image is a completely blank white document with no visible content, text, or markings.

- (b) Calculate $\text{Var}(X)$ 2

[http://www.scribd.com/doc/10678924/](#)

Question 19 (2 marks)**Marks**

The length of steel rods produced by a machine is normally distributed with a standard deviation of 3 mm. It is found that 2.5% of all rods are less than 25 mm long. Find the mean length of rods produced by the machine.

2**Question 20 (8 marks)**

Consider the function $f(x) = x^3 + 6x^2 + 9x + 4$ in the domain $-4 \leq x \leq 1$

(a) Find the coordinates of any stationary points and determine their nature.

3

(b) Determine the coordinates of its point(s) of inflexion.

2


Question 20 continued on next page

Question 20 continued

Marks

- (c) Draw a sketch of the curve $y = f(x)$ in the domain $-4 \leq x \leq 1$ clearly showing all essential features.


2



- (d) What is the global maximum value of the curve in the domain $-4 \leq x \leq 1$

1

The diagram shows the graph of $y = a \sin(bx) + c$ for $0 \leq x \leq 2\pi$, where a , b and c are positive integers.



2

Find the values of a , b and c .

Question 22 (5 marks)

Marks


A pet ownership survey resulted in the following results:

$$P(C) = \frac{3}{7}, \quad P(D|\bar{C}) = \frac{2}{5}, \quad \text{and} \quad P(\bar{D}|C) = \frac{3}{4}.$$

Where C is the event that “a person has a cat” and D is the event that “a person has a dog”

(a) Complete the probability tree by marking a probability on each branch.

2



(b) If one person is chosen at random, find the probability that the person has:

i) a cat and a dog

1

ii) at least one pet (cat or dog)

2

Question 23 (5 marks)

Marks

The function f is defined by $f(x) = 2 + \sqrt{x-3}$ for $x \geq 3$.

The function g is defined by $g(x) = \frac{12}{x} + 2$ for $x > 0$

- (a) Write the domain and range of the function f using interval notation.


3

- (b) Write an expression for the composite function $h(x) = g(f(x))$ and hence find a value for $g(f(12))$

2

Question 24 (5 marks)

The diagram shows the graphs $y = \sin x$ and $y = \cos x$, $0 \leq x \leq 2\pi$. The graphs intersect at A and B .



- (a) Show that A has coordinates $\left(\frac{\pi}{4}, \frac{1}{\sqrt{2}}\right)$.

2

Question 24 continued on next page

Question 24 continued


- (b) Find the area enclosed by the two graphs.

3

[illegible]

Question 25 (3 marks)**Marks**

In the diagram, the points $A(-5,3)$, $B(2,2)$ and $C(1,-5)$ are shown.



- (a) Calculate the gradient of AC .

1

- (b) Find the coordinates of X , the midpoint of AC .

1

- (c) Find the coordinates of D if X is also the midpoint of BD .

1

Question 26 (5 marks)**Marks**

A particle is moving in a straight line. Its velocity for $t \geq 0$ is given by $v = \frac{4}{t+1} - 2t$, where time is in seconds and displacement in metres.

(a) Find when the particle changes direction.

2

(b) Find the exact distance travelled in the first 2 seconds.

3

Question 27 (6 marks)

Marks

The continuous random variable X has probability density function $f(x)$ given by

$$f(x) = \begin{cases} k(x^2 - 2x + 2) & 1 \leq x \leq 4 \\ 0 & \text{otherwise} \end{cases}$$

Where k is a constant

- (a) Show that $k = \frac{1}{12}$

2

- (b) Fully define the cumulative distribution function $F(x)$.

3

1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525

Question 27 continued on next page

Question 27 continued

(c) Show that the median of X lies between $x = 3.2$ and $x = 3.3$

1

Question 28 (3 marks)**Marks**

Consider the geometric series $1 + \frac{4}{3}\sin^2 x + \frac{16}{9}\sin^4 x + \frac{64}{27}\sin^6 x + \dots$

- (a) When the limiting sum exists, find an expression for its value.

1

- (b) For what values of x in the interval $0 < x < \frac{\pi}{2}$ does the limiting sum of this series exist?

2

Question 29 (4 marks)

- (a) Find $\int \sec^2(2x) \tan^4(2x) dx$

2

- (b) Find $\int \frac{5x^2}{x^3 + 1} dx$

2

Question 31 (4 marks)**Marks**

The table below shows the future values of an annuity, for different rates of interest and for different numbers of compounding periods, where contributions of \$1 are made at the end of each compounding period.

Table of future value interest factors

<i>n</i>	1%	2%	3%	4%	5%	6%
1	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
2	2.0100	2.0200	2.0300	2.0400	2.0500	2.0600
3	3.0301	3.0604	3.0909	3.1216	3.1525	3.1836
4	4.0604	4.1216	4.1836	4.2465	4.3101	4.3746
5	5.1010	5.2040	5.3091	5.4163	5.5256	5.6371
6	6.1520	6.3081	6.4684	6.6330	6.8019	6.9753

- (a) An annuity account is opened and contributions of \$1200 are made at the end of each half year for 3 years at an interest rate of 4% p.a. compounding half yearly. Calculate the final amount in the account immediately after the last contribution is made.

2

- (b) Calculate the single lump sum amount that would need to be invested at the start to reach the same final amount at the end of the 3 years with the same interest rate of 4% compounding half yearly.

2

Question 32 continued

(c)

It is known that $\frac{dV}{dr} = \frac{300\pi r - \pi^2 r^3}{\sqrt{450\pi - \pi^2 r^2}}$ (do not prove this).

2

Hence or otherwise, find the value of r for the volume of the cone to be a maximum.

End of paper

Section I**10 marks****Attempt questions 1 - 10****Allow about 15 minutes for this section**

Use the multiple-choice answer sheet for questions 1-10

1. What is the value of $\operatorname{cosec} \frac{\pi}{3}$ to three significant figures?
- (A) 1.00
(B) 1.15
(C) 1.41
(D) 2.00
2. What is the value of c for which the circle $(x-3)^2 + (y-2)^2 = c$ touches the x axis?
- (A) 2
(B) 3
(C) 4
(D) 9
3. What is the equation of the tangent to $y = x^2 - 3$ at $x = -1$?
- (A) $y = -2x - 4$
(B) $y = 2x - 4$
(C) $y = \frac{x}{2} - \frac{3}{2}$
(D) $y = -\frac{x}{2} - \frac{3}{2}$

4. Which statement is true for an ungrouped data set with no outliers?

- (A) The largest possible range is 2 times the interquartile range.
- (B) The largest possible range is 3 times the interquartile range.
- ☒ (C) The largest possible range is 4 times the interquartile range.
- (D) The largest possible range is 5 times the interquartile range.

5. Which one of the following is the set of all solutions to $2x^2 - 5x + 2 \geq 0$?

- (A) $\left[\frac{1}{2}, 2\right]$
- (B) $\left(\frac{1}{2}, 2\right)$
- (C) $\left(-\infty, \frac{1}{2}\right) \cup (2, \infty)$
- ☒ (D) $\left(-\infty, \frac{1}{2}\right] \cup [2, \infty)$

6. The graph of $y = f(x)$ has a stationary point at $(2, -3)$.

Which one of the following is a guaranteed stationary point of $y = -f\left(\frac{x}{2}\right) - 5$?

- (A) $(1, -2)$
- (B) $(1, 2)$
- ☒ (C) $(4, -2)$
- (D) $(4, 2)$

7. What is the period and amplitude for the curve $y = \sin \pi x$?

- (A) Amplitude = 1; Period = 2
- (B) Amplitude = π ; Period = 2
- (C) Amplitude = 1; Period = 2π
- (D) Amplitude = π ; Period = 2π

8. If the z scores on an examination are normally distributed and $P(z < N) = 0.6$ for some number N , what is the value of $P(-N < z < N)$?

- (A) 0.1
- (B) 0.2
- (C) 0.3
- (D) 0.4

9. Which one of the following equations is NOT correct?

(A) $\int x(x^2 - 1)^2 dx = \frac{(x^2 - 1)^3}{6} + c$

(B) $\int_{-3}^3 \sqrt{9 - x^2} dx = \frac{9\pi}{2}$

(C) $\int_{-1}^1 3^x dx = \frac{1}{\ln 3} \left(3 - \frac{1}{3} \right)$

(D) $\int_{-5}^5 4x^4 - x^3 + \cos x dx = 0$

10. Consider the series $\sqrt{5} + \sqrt{45} + \sqrt{125} + \dots + z = 225\sqrt{5}$, the value of z is:

- (A) $25\sqrt{5}$
- (B) $29\sqrt{5}$
- (C) $30\sqrt{5}$
- (D) $35\sqrt{5}$

Section II

90 marks

Attempt all questions

Allow about 2 hours and 45 minutes for this section

Answer each question in the spaces provided.

Your responses should include relevant mathematical reasoning and/or calculations.

Extra writing space is provided at the back of the examination paper.

Question 11 (2 marks)

Marks

What angle does the line $2x + 3y + 6 = 0$ make with the positive x -axis? Round to the nearest minute.



2

$$m = -\frac{2}{3} \quad \therefore \tan \alpha = -\frac{2}{3}$$

$$\alpha = 146^\circ 19'$$

Question 12 (2 marks)


Sketch a possible function which could have the gradient function as graphed below.



2

Question 13 (3 marks)**Marks**

In triangle ABC , the length of $AB=5\text{cm}$, $AC=13\text{cm}$ and $\cos\angle BAC = \frac{1}{8}$



- (a) Find the exact value of $\sin\angle BAC$

2

$$\sin \angle BAC = \frac{3\sqrt{7}}{5}$$

- (b) Find the area of triangle ABC

1

$$\begin{aligned} A &= \frac{1}{2} \times 5 \times 13 \times \sin \angle BAC \\ &= \frac{1}{2} \times 5 \times 13 \times \frac{3\sqrt{7}}{5} \\ &= \frac{195\sqrt{7}}{16} \\ &\approx 32.2 \text{ u}^2 \end{aligned}$$

Question 14 (3 marks)

Solve $2\log x = \log(5x+6)$

3

$$\begin{aligned} \log x^2 &= \log(5x+6) \\ \therefore x^2 &= 5x+6 \\ x^2 - 5x - 6 &= 0 \\ (x-6)(x+1) &= 0 \\ x &= 6, -1 \quad \text{but } x > 0 \\ \therefore x &= 6 \end{aligned}$$

Question 15 (3 marks)

Marks

Solve $|1 - 2\cos^2 x| = 1$ for $0 \leq x \leq 2\pi$

3

$$1 - 2\cos^2 x = 1$$

$$2\cos^2 x = 0$$

$$\cos^2 x = 0$$

$$x = \frac{\pi}{2}, \frac{3\pi}{2}$$

$$1 - 2\cos^2 x = -1$$

$$2\cos^2 x = 2$$

$$\cos^2 x = 1$$

$$\cos x = \pm 1$$

$$x = 0, \pi, 2\pi$$

$$\therefore x = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi$$

Question 16 (5 marks)

Differentiate the following expressions.

Marks

(a) $\log_5(\tan x)$

2

$$\frac{d}{dx}(\log_5(\tan x)) = \frac{\sec^2 x}{\ln 5 \tan x}$$

(b) $\frac{2^x}{e^x}$

3

$$u = 2^x$$

$$v = e^x$$

$$u' = \ln 2 \cdot 2^x$$

$$v' = e^x$$

$$\frac{d}{dx} \left(\frac{2^x}{e^x} \right) = \frac{\ln 2 \cdot 2^x \cdot e^x - 2^x \cdot e^x}{(e^x)^2}$$


$$= \frac{e^x 2^x (\ln 2 - 1)}{e^{2x}}$$

$$= \frac{2^x (\ln 2 - 1)}{e^x}$$

Question 17 (5 marks)

Marks

Consider the graph $y = f(x)$. Both arcs have a radius of four units.



Using the graph of $y = f(x)$, $x \geq 0$, evaluate exactly the following integrals.

(a)

$$\int_0^{12} f(x) dx$$

3

$$= (5 \times 6) + (6 + 8) + \frac{1}{4} \times \pi \times 4^2 + \frac{1}{2} \times 2 \times 2$$

$$= 46 + 4\pi$$

(b)

$$\int_0^{18} f(x) dx$$

2

$$= \int_0^{12} f(x) dx + \int_{12}^{18} f(x) dx$$

$$= 46 + 4\pi - \left(\frac{1}{2} \times 2 \times 2 + (4 \times 6 - \frac{1}{4} \times \pi \times 4^2) \right)$$

$$= 46 + 4\pi - (26 - 4\pi)$$

$$= 20 + 8\pi$$

Question 18 (5 marks)**Marks**

The discrete random variable X has probability distribution shown in the table below

x	-1	0	1	2	3
$P(X=x)$	a	b	0.2	0.15	0.13

and $E(X) = 0.55$

- (a) By forming a pair of simultaneous equations, or otherwise, find the values of a and b . 3

$$\begin{aligned}
 E(X) &= \sum x_i p_i \\
 &= -1(a) + 0(b) + 1(0.2) + 2(0.15) + 3(0.13) \\
 \therefore 0.55 &= -a + 0.89 \\
 a &= 0.34 \\
 \downarrow 0.34 + b + 0.2 + 0.15 + 0.13 &= 1 \\
 \therefore b &= 0.18
 \end{aligned}$$

- (b) Calculate $\text{Var}(X)$ 2

$$\begin{aligned}
 \text{Var}(X) &= E(X^2) - \mu^2 \\
 E(X^2) &= 1(a) + 0(b) + 1(0.2) + 4(0.15) + 9(0.13) \\
 &= 2.31 \\
 \therefore \text{Var}(X) &= 2.31 - 0.55^2 \\
 &= 2.0075
 \end{aligned}$$

Question 19 (2 marks)**Marks**

The length of steel rods produced by a machine is normally distributed with a standard deviation of 3 mm. It is found that 2.5% of all rods are less than 25 mm long. Find the mean length of rods produced by the machine.

2

$$\sigma = 3 \quad x = 25 \quad z = -2$$

$$\therefore -2 = \frac{25 - \mu}{3}$$

$$25 - \mu = -6$$

$$\mu = 31$$

Question 20 (8 marks)

Consider the function $f(x) = x^3 + 6x^2 + 9x + 4$ in the domain $-4 \leq x \leq 1$

- (a) Find the coordinates of any stationary points and determine their nature.

3

$$f'(x) = 3x^2 + 12x + 9$$

$$f''(x) = 6x + 12$$

stat. pts at $f'(x) = 0$

$$3(x^2 + 4x + 3) = 0$$

$$(x+3)(x+1) = 0$$

$$x = -3$$

$$x = -1$$

$$y = 4$$

$$y = 0$$

$$f''(-3) = -18 + 12 < 0$$

$$f''(-1) = -6 + 12 > 0$$

$\therefore$ max turning pt at

$\therefore$ min turning pt at

$$(-3, 4)$$

$$(-1, 0)$$

- (b) Determine the coordinates of its point(s) of inflexion.

2

poss. pt. inflex at $y'' = 0$

$$\therefore 6x + 12 = 0$$

$$x = -2$$

$$y = 2$$

x	-1	-2	-3
$f''(x)$	6	0	-6

$\therefore$ change in concavity
 $\therefore$ pt. inflex at $(-2, 2)$


Question 20 continued on next page

Question 20 continued

Marks

- (c) Draw a sketch of the curve $y = f(x)$ in the domain $-4 \leq x \leq 1$ clearly showing all essential features.

2



endpoints $(-4, 0)$ & $(1, 20)$
 $y\text{-int} = 4$


- (d) What is the global maximum value of the curve in the domain $-4 \leq x \leq 1$

1

20

Question 21 (2 marks)

The diagram shows the graph of $y = a \sin(bx) + c$ for $0 \leq x \leq 2\pi$, where a , b and c are positive integers.



2

Find the values of a , b and c .

$$a = 3$$

$$T = \pi$$

$$c = 1$$

$$\therefore \frac{2\pi}{b} = \pi$$

$$\therefore b = 2$$

Question 22 (5 marks)

Marks


A pet ownership survey resulted in the following results:

$$P(C) = \frac{3}{7}, \quad P(D|\bar{C}) = \frac{2}{5}, \quad \text{and} \quad P(\bar{D}|C) = \frac{3}{4}.$$

Where C is the event that "a person has a cat" and D is the event that "a person has a dog"

- (a) Complete the probability tree by marking a probability on each branch.

2



- (b) If one person is chosen at random, find the probability that the person has:

- i) a cat and a dog

1

$$P(C \cap D) = \frac{3}{7} \times \frac{1}{4} = \frac{3}{28}$$

- ii) at least one pet (cat or dog)

2

$$P(\text{at least 1 pet}) = 1 - P(\bar{D}\bar{C}) = 1 - \frac{4}{7} \times \frac{3}{5} = \frac{23}{35}$$

Question 23 (5 marks)**Marks**

The function f is defined by $f(x) = 2 + \sqrt{x-3}$ for $x \geq 3$.

The function g is defined by $g(x) = \frac{12}{x} + 2$ for $x > 0$

- (a) Write the domain and range of the function f using interval notation.

3

domain: $x \geq 3$ $[3, \infty)$

range: $y \geq 2$ $[2, \infty)$

- (b) Write an expression for the composite function $h(x) = g(f(x))$ and hence find a value for $g(f(12))$

2


$$g(f(x)) = \frac{12}{2 + \sqrt{x-3}} + 2$$

$$g(f(12)) = \frac{12}{2 + \sqrt{12-3}} + 2$$

$$= \frac{22}{5}$$

Question 24 (5 marks)

The diagram shows the graphs $y = \sin x$ and $y = \cos x$, $0 \leq x \leq 2\pi$. The graphs intersect at A and B .



- (a) Show that A has coordinates $\left(\frac{\pi}{4}, \frac{1}{\sqrt{2}}\right)$.

2

pt. int at $\sin x = \cos x$ ($\div \cos x$)
 $\tan x = 1$

$$x = \frac{\pi}{4} \quad 0 < x < \frac{\pi}{2}$$

sub in $y = \sin x$

$$\therefore y = \sin \frac{\pi}{4}$$

$$y = \frac{1}{\sqrt{2}}$$

$$\therefore A\left(\frac{\pi}{4}, \frac{1}{\sqrt{2}}\right)$$

Question 24 continued on next page

Question 24 continued

- (b) Find the area enclosed by the two graphs.

3

$$\text{solve for } B. \quad \tan x = 1 \quad \pi < x < \frac{3\pi}{2}$$

$$\therefore x = \frac{5\pi}{4}$$

$$A = \int_{\frac{\pi}{4}}^{\frac{5\pi}{4}} (\sin x - \cos x) dx$$

$$= [-\cos x - \sin x]_{\frac{\pi}{4}}^{\frac{5\pi}{4}}$$

$$= -[\cos x + \sin x]_{\frac{\pi}{4}}^{\frac{5\pi}{4}}$$


$$= -\left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} - \left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}\right)\right)$$

$$= -\left(-\frac{2}{\sqrt{2}}\right)$$

$$= 2\sqrt{2} \quad u^2$$

Question 25 (3 marks)**Marks**

In the diagram, the points $A(-5,3)$, $B(2,2)$ and $C(1,-5)$ are shown.



- (a) Calculate the gradient of AC .

1

$$\begin{aligned} m(AC) &= \frac{-5-3}{1+5} \\ &= -\frac{8}{6} \\ &= -\frac{4}{3} \end{aligned}$$

- (b) Find the coordinates of X , the midpoint of AC .

1

$$\begin{aligned} M &= \left(\frac{-5+1}{2}, \frac{3-5}{2} \right) \\ &= (-2, -1) \end{aligned}$$

- (c) Find the coordinates of D if X is also the midpoint of BD .

1

$$\begin{aligned} -2 &= \frac{2+x}{2} & -1 &= \frac{2+y}{2} \\ -4 &= 2+x & 2+y &= -2 \\ x &= -6 & y &= -4 \\ \therefore D &= (-6, -4) \end{aligned}$$

Question 26 (5 marks)**Marks**

A particle is moving in a straight line. Its velocity for $t \geq 0$ is given by $v = \frac{4}{t+1} - 2t$, where time is in seconds and displacement in metres.

- (a) Find when the particle changes direction.

2

$$\begin{array}{l|l} \text{change direction at } v=0 & \\ \hline \therefore \frac{4}{t+1} - 2t = 0 & (t+2)(t-1) = 0 \\ & t = -2, 1, \quad t > 0 \\ & \therefore t = 1 \\ 4 - 2t^2 - 2t = 0 & \\ t^2 + t - 2 = 0 & \end{array}$$

- (b) Find the exact distance travelled in the first 2 seconds.

3

$$\begin{aligned} d &= \left| \int_0^1 \left(\frac{4}{t+1} - 2t \right) dt \right| + \left| \int_1^2 \left(\frac{4}{t+1} - 2t \right) dt \right| \\ &= \left| [4\ln|t+1| - t^2]_0^1 \right| + \left| [4\ln|t+1| - t^2]_1^2 \right| \\ &= |4\ln 2 - 1| + |4\ln 3 - 3 - 4\ln 2| \\ &= 4\ln 2 - 1 - (4\ln 3 - 3 - 4\ln 2) \\ &= 8\ln 2 + 2 - 4\ln 3 \end{aligned}$$

Question 27 (6 marks)**Marks**

The continuous random variable X has probability density function $f(x)$ given by

$$f(x) = \begin{cases} k(x^2 - 2x + 2) & 1 \leq x \leq 4 \\ 0 & \text{otherwise} \end{cases}$$

Where k is a constant

- (a) Show that $k = \frac{1}{12}$

2

$$\begin{aligned} F(x) &= k \int_1^4 (x^2 - 2x + 2) dx \\ \therefore k \left[\frac{x^3}{3} - x^2 + 2x \right]_1^4 &= 1 \\ \therefore k \left(\left(\frac{64}{3} - 16 + 8 \right) - \left(\frac{1}{3} - 1 + 2 \right) \right) &= 1 \\ \therefore 12k &= 1, \quad k = \frac{1}{12} \end{aligned}$$

- (b) Fully define the cumulative distribution function $F(x)$.

3

$$\begin{aligned} \text{For } 1 \leq x \leq 4 \\ F(x) &= \frac{1}{12} \int_1^x (x^2 - 2x + 2) dx \\ &= \frac{1}{12} \left[\frac{x^3}{3} - x^2 + 2x \right]_1^x \\ &= \frac{1}{12} \left[\frac{x^3}{3} - x^2 + 2x - \left(\frac{1}{3} - 1 + 2 \right) \right] \\ &= \frac{1}{12} \left[\frac{x^3}{3} - x^2 + 2x - \frac{4}{3} \right] \\ &= \frac{x^3}{36} - \frac{x^2}{12} + \frac{x}{6} - \frac{1}{9} \\ \therefore F(x) &= \begin{cases} 0 & x < 1 \\ \frac{x^3}{36} - \frac{x^2}{12} + \frac{x}{6} - \frac{1}{9} & 1 \leq x \leq 4 \\ 1 & x > 4 \end{cases} \end{aligned}$$

Question 27 continued on next page

Question 27 continued

(c) Show that the median of X lies between $x = 3.2$ and $x = 3.3$

1

$$F(3,2) = \frac{3 \cdot 2^3}{36} - \frac{3 \cdot 2^2}{12} + \frac{3 \cdot 2}{6} - \frac{1}{2}$$

$$= 0.479 < 0.5$$

$$F(3.3) = \frac{3.3^3}{36} - \frac{3.3^2}{12} + \frac{3.3}{6} - \frac{1}{9}$$
$$= 0.5296 > 0.5$$

$$F(\text{median}) = 0.5$$

$\therefore$ median lies between $x = 3.2$
+ $x = 3.3$

Question 28 (3 marks)

Marks

Consider the geometric series $1 + \frac{4}{3}\sin^2 x + \frac{16}{9}\sin^4 x + \frac{64}{27}\sin^6 x + \dots$

- (a) When the limiting sum exists, find an expression for its value.

1

$$r = \frac{4}{3} \sin^2 x$$

$$S_{\infty} = \frac{1}{1 - \frac{4}{3} \sin^2 x}$$

$$a = 1$$

- (b) For what values of x in the interval $0 < x < \frac{\pi}{2}$ does the limiting sum of this series exist?

2

$$S_{\infty} \text{ exists for } |r| < 1 \quad \text{ie } -1 < r < 1$$

$$\therefore \left| \frac{4}{3} \sin^2 x \right| < 1 \quad \text{ie } -1 < \frac{4}{3} \sin^2 x < 1$$

$$0 < \frac{4}{3} \sin^2 x < 1 \quad \text{in Q1}$$

$$0 < \sin^2 x < \frac{3}{4}$$

$$0 < \sin x < \frac{\sqrt{3}}{2}$$

$$\therefore 0 < x < \frac{\pi}{3}$$

Question 29 (4 marks)

- (a) Find $\int \sec^2(2x) \tan^4(2x) dx$

2

$$= \frac{1}{2} \int 2 \sec^2(2x) \tan^4(2x) dx$$

$$= \frac{1}{2} \frac{\tan^5(2x)}{5} + C$$

$$= \frac{\tan^5(2x)}{10} + C$$

- (b) Find $\int \frac{5x^2}{x^3+1} dx$

2

$$= \frac{5}{3} \int \frac{3x^2}{x^3+1} dx$$

$$= \frac{5}{3} \ln|x^3+1| + C$$

Question 30 (4 marks)

Marks

Luke suspects that the rate at which he spends cash is affected by the amount of cash he withdrew at his previous visit to an ATM.

The table below shows the amount of cash withdrawn, \$ x , from an ATM, and the time, y hours, until Luke's next withdrawal from an ATM.

Withdrawal	1	2	3	4	5	6	7	8	9	10
x	40	10	100	110	120	150	20	90	80	130
y	56	62	195	330	94	270	48	196	214	286

- (a) Find the equation of the least squares regression line for y in terms of x , for the withdrawals 1 to 10 and hence estimate how much cash (to the nearest \$10) Luke would need to withdraw from the ATM at his previous visit in order to not need to visit an ATM again for 120 hours.

2

from calculator:

$$A = 30.26$$

$$B = 1.7$$

$$y = A + Bx$$

$$y = 1.7x + 30.26$$

Find x at $y = 120$

$$120 = 1.7x + 30.26$$

$$\therefore x = 52.79$$

$\therefore$ withdrawal = \$60

- (b) Calculate the correlation coefficient between x and y for the withdrawals 1 to 10. Describe the nature of the correlation.

2

$$r = 0.777$$

{strong} positive correlation

Question 31 (4 marks)

Marks

The table below shows the future values of an annuity, for different rates of interest and for different numbers of compounding periods, where contributions of \$1 are made at the end of each compounding period.

Table of future value interest factors

n	1%	2%	3%	4%	5%	6%
1	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
2	2.0100	2.0200	2.0300	2.0400	2.0500	2.0600
3	3.0301	3.0604	3.0909	3.1216	3.1525	3.1836
4	4.0604	4.1216	4.1836	4.2465	4.3101	4.3746
5	5.1010	5.2040	5.3091	5.4163	5.5256	5.6371
6	6.1520	6.3081	6.4684	6.6330	6.8019	6.9753

- (a) An annuity account is opened and contributions of \$1200 are made at the end of each half year for 3 years at an interest rate of 4% p.a. compounding half yearly. Calculate the final amount in the account immediately after the last contribution is made.

2

$$FV = 1200 \times 6.3081$$

$$= \$7569.72$$

$$r = 2\% \text{ half year}$$

$$n = 6$$

- (b) Calculate the single lump sum amount that would need to be invested at the start to reach the same final amount at the end of the 3 years with the same interest rate of 4% compounding half yearly.

2

$$7569.72 = P(1.02)^6$$

$$\therefore P = \$6721.69$$

Question 32 (6 marks)**Marks**

An open cone, of radius r cm and height h cm is made from a sector of a circle. The area of the sector used is 300 cm^2




Figure 1




Figure 2

- (a) Show from Figure 1 that the slant height l is given by $l^2 = \frac{450}{\pi}$

2

$$\begin{aligned}
 A &= \frac{1}{2} r^2 \theta \\
 A &= \frac{1}{2} \times l^2 \times \left(2\pi - \frac{2\pi}{3}\right) \\
 &= \frac{1}{2} l^2 \times \frac{4\pi}{3} \\
 300 &= \frac{2}{3} \pi l^2 \\
 \therefore l^2 &= \frac{900}{2\pi} \\
 l^2 &= \frac{450}{\pi}
 \end{aligned}$$

- (b) In Figure 2 it is given that $h = \sqrt{l^2 - r^2}$ (do not prove this).

2

Show that the volume of the cone is given by $V = \frac{1}{3} r^2 \sqrt{450\pi - \pi^2 r^2}$

$$V = \frac{1}{3} \pi r^2 h$$

$$V = \frac{1}{3} \pi r^2 \sqrt{l^2 - r^2}$$

$$= \frac{1}{3} \pi r^2 \sqrt{\frac{450}{\pi} - r^2}$$

$$= \frac{1}{3} r^2 \sqrt{\pi^2 \left(\frac{450 - \pi r^2}{\pi} \right)}$$

$$= \frac{1}{3} r^2 \sqrt{450\pi - \pi^2 r^2}$$

Question 32 continued on next page

Question 32 continued

(c)

It is known that $\frac{dV}{dr} = \frac{300\pi r - \pi^2 r^3}{\sqrt{450\pi - \pi^2 r^2}}$ (do not prove this).

2

Hence or otherwise, find the value of r for the volume of the cone to be a maximum.

... stat pt at $\frac{dV}{dr} = 0$

$$\therefore 300\pi r - \pi^2 r^3 = 0$$

$$\therefore \pi r (300 - \pi r^2) = 0 \quad r > 0$$

$$\pi r^2 = 300$$

$$r^2 = \frac{300}{\pi}$$

$$r = \sqrt{\frac{300}{\pi}} = \frac{10\sqrt{3}}{\sqrt{\pi}}$$

r	$\sqrt{\frac{290}{\pi}}$	$\sqrt{\frac{300}{\pi}}$	$\sqrt{\frac{310}{\pi}}$
$\frac{dV}{dr}$	13.46	0	-31.38

$\therefore$ max vol at

$$r = \sqrt{\frac{300}{\pi}}$$

End of paper