

STUDENT NAME: _____

TEACHER: _____



HURLSTONE
AGRICULTURAL
HIGH
SCHOOL

2021

Year 11
Assessment Task 1

Mathematics Advanced

Examiners

- **Ms T Tarannum**, Ms M Sabah, Mr S Faulds.

**General
Instructions**

- Working time – 40 minutes
- Reading time – 3 minutes
- Write using blue or black pen
- NESA-approved calculator may be used
- A Reference sheet is provided
- In Questions 4 – 6, show relevant mathematical reasoning and/or calculations

**Total marks:
30****Section I – 3 marks**

- Attempt Questions 1 – 3
- Allow about 3 minutes for this section

Section II – 27 marks

- Attempt Questions 4 – 6
- Allow about 37 minutes for this section

SECTION I

3 marks

Attempt Questions 1 – 3

Allow about 3 minutes for this section.

Use the multiple-choice answer sheet for Questions 1 – 3

Question 1

What is the value of $x + y$ if $5x + 8y = 67$ and $2x - y = 31$?

- A. 14
- B. 16
- C. 15
- D. 0

Question 2

If $\sin x = \frac{3}{4}$, then which of the following is the value of $\cos x$ for $0^\circ \leq x \leq 90^\circ$?

- A. $\frac{2}{3}$
- B. $\frac{\sqrt{3}}{2}$
- C. $\frac{\sqrt{7}}{4}$
- D. $\frac{1}{2}$

Question 3

Two functions are defined as follows:

$$g(x) = 5 - 4x$$

$$h(x) = (x - 1)^2$$

What is the correct expression for the composite function $g(h(x))$?

- A. $1 - 4x^2$
- B. $x^2 - x + 1$
- C. $1 + 8x - 4x^2$
- D. $16x^2 - 16x + 16$

**Mathematics Advanced**
Mathematics Extension 1
Mathematics Extension 2**REFERENCE SHEET****Measurement****Length**

$$l = \frac{\theta}{360} \times 2\pi r$$

Area

$$A = \frac{\theta}{360} \times \pi r^2$$

$$A = \frac{h}{2}(a + b)$$

Surface area

$$A = 2\pi r^2 + 2\pi rh$$

$$A = 4\pi r^2$$

Volume

$$V = \frac{1}{3}Ah$$

$$V = \frac{4}{3}\pi r^3$$

Functions

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For $ax^3 + bx^2 + cx + d = 0$:

$$\alpha + \beta + \gamma = -\frac{b}{a}$$

$$\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a}$$

$$\text{and } \alpha\beta\gamma = -\frac{d}{a}$$

Relations

$$(x - h)^2 + (y - k)^2 = r^2$$

Financial Mathematics

$$A = P(1 + r)^n$$

Sequences and series

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2}[2a + (n - 1)d] = \frac{n}{2}(a + l)$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(1 - r^n)}{1 - r} = \frac{a(r^n - 1)}{r - 1}, r \neq 1$$

$$S = \frac{a}{1 - r}, |r| < 1$$

Logarithmic and Exponential Functions

$$\log_a a^x = x = a^{\log_a x}$$

$$\log_a x = \frac{\log_b x}{\log_b a}$$

$$a^x = e^{x \ln a}$$

Trigonometric Functions

$$\sin A = \frac{\text{opp}}{\text{hyp}}, \quad \cos A = \frac{\text{adj}}{\text{hyp}}, \quad \tan A = \frac{\text{opp}}{\text{adj}}$$

$$A = \frac{1}{2}ab \sin C$$



$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$l = r\theta$$

$$A = \frac{1}{2}r^2\theta$$



Trigonometric identities

$$\sec A = \frac{1}{\cos A}, \quad \cos A \neq 0$$

$$\operatorname{cosec} A = \frac{1}{\sin A}, \quad \sin A \neq 0$$

$$\cot A = \frac{\cos A}{\sin A}, \quad \sin A \neq 0$$

$$\cos^2 x + \sin^2 x = 1$$

Compound angles

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\text{If } t = \tan \frac{A}{2} \text{ then } \sin A = \frac{2t}{1 + t^2}$$

$$\cos A = \frac{1 - t^2}{1 + t^2}$$

$$\tan A = \frac{2t}{1 - t^2}$$

$$\cos A \cos B = \frac{1}{2} [\cos(A - B) + \cos(A + B)]$$

$$\sin A \sin B = \frac{1}{2} [\cos(A - B) - \cos(A + B)]$$

$$\sin A \cos B = \frac{1}{2} [\sin(A + B) + \sin(A - B)]$$

$$\cos A \sin B = \frac{1}{2} [\sin(A + B) - \sin(A - B)]$$

$$\sin^2 nx = \frac{1}{2} (1 - \cos 2nx)$$


$$\cos^2 nx = \frac{1}{2} (1 + \cos 2nx)$$

Statistical Analysis

$$z = \frac{x - \mu}{\sigma}$$

An outlier is a score
less than $Q_1 - 1.5 \times IQR$
or
more than $Q_3 + 1.5 \times IQR$

Normal distribution



- approximately 68% of scores have z -scores between -1 and 1
- approximately 95% of scores have z -scores between -2 and 2
- approximately 99.7% of scores have z -scores between -3 and 3

$$E(X) = \mu$$

$$\operatorname{Var}(X) = E[(X - \mu)^2] = E(X^2) - \mu^2$$

Probability

$$P(A \cap B) = P(A)P(B)$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad P(B) \neq 0$$

Continuous random variables

$$P(X \leq x) = \int_a^x f(x) dx$$

$$P(a < X < b) = \int_a^b f(x) dx$$

Binomial distribution

$$P(X = r) = {}^nC_r p^r (1 - p)^{n-r}$$

$$X \sim \operatorname{Bin}(n, p)$$

$$\Rightarrow P(X = x)$$

$$= \binom{n}{x} p^x (1 - p)^{n-x}, \quad x = 0, 1, \dots, n$$

$$E(X) = np$$

$$\operatorname{Var}(X) = np(1 - p)$$

Differential Calculus

Function

Derivative

$$y = f(x)^n$$

$$\frac{dy}{dx} = n f'(x) [f(x)]^{n-1}$$

$$y = uv$$

$$\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$y = g(u) \text{ where } u = f(x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$y = \frac{u}{v}$$

$$\frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

$$y = \sin f(x)$$

$$\frac{dy}{dx} = f'(x) \cos f(x)$$

$$y = \cos f(x)$$

$$\frac{dy}{dx} = -f'(x) \sin f(x)$$

$$y = \tan f(x)$$

$$\frac{dy}{dx} = f'(x) \sec^2 f(x)$$

$$y = e^{f(x)}$$

$$\frac{dy}{dx} = f'(x) e^{f(x)}$$

$$y = \ln f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{f(x)}$$

$$y = a^{f(x)}$$

$$\frac{dy}{dx} = (\ln a) f'(x) a^{f(x)}$$

$$y = \log_a f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{(\ln a) f(x)}$$

$$y = \sin^{-1} f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{\sqrt{1 - [f(x)]^2}}$$

$$y = \cos^{-1} f(x)$$

$$\frac{dy}{dx} = -\frac{f'(x)}{\sqrt{1 - [f(x)]^2}}$$

$$y = \tan^{-1} f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{1 + [f(x)]^2}$$

Integral Calculus

$$\int f'(x) [f(x)]^n dx = \frac{1}{n+1} [f(x)]^{n+1} + c$$

where $n \neq -1$

$$\int f'(x) \sin f(x) dx = -\cos f(x) + c$$

$$\int f'(x) \cos f(x) dx = \sin f(x) + c$$

$$\int f'(x) \sec^2 f(x) dx = \tan f(x) + c$$

$$\int f'(x) e^{f(x)} dx = e^{f(x)} + c$$

$$\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c$$

$$\int f'(x) a^{f(x)} dx = \frac{a^{f(x)}}{\ln a} + c$$

$$\int \frac{f'(x)}{\sqrt{a^2 - [f(x)]^2}} dx = \sin^{-1} \frac{f(x)}{a} + c$$

$$\int \frac{f'(x)}{a^2 + [f(x)]^2} dx = \frac{1}{a} \tan^{-1} \frac{f(x)}{a} + c$$

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

$$\int_a^b f(x) dx$$

$$\approx \frac{b-a}{2n} \left\{ f(a) + f(b) + 2[f(x_1) + \dots + f(x_{n-1})] \right\}$$

where $a = x_0$ and $b = x_n$

Combinatorics

$${}^nP_r = \frac{n!}{(n-r)!}$$

$$\binom{n}{r} = {}^nC_r = \frac{n!}{r!(n-r)!}$$

$$(x+a)^n = x^n + \binom{n}{1}x^{n-1}a + \cdots + \binom{n}{r}x^{n-r}a^r + \cdots + a^n$$

Vectors

$$|\underline{u}| = |x\underline{i} + y\underline{j}| = \sqrt{x^2 + y^2}$$

$$\underline{u} \cdot \underline{v} = |\underline{u}| |\underline{v}| \cos \theta = x_1x_2 + y_1y_2,$$

$$\text{where } \underline{u} = x_1\underline{i} + y_1\underline{j}$$

$$\text{and } \underline{v} = x_2\underline{i} + y_2\underline{j}$$

$$\underline{r} = \underline{a} + \lambda \underline{b}$$

Complex Numbers

$$\begin{aligned} z &= a + ib = r(\cos \theta + i \sin \theta) \\ &= re^{i\theta} \end{aligned}$$

$$\begin{aligned} [r(\cos \theta + i \sin \theta)]^n &= r^n(\cos n\theta + i \sin n\theta) \\ &= r^n e^{in\theta} \end{aligned}$$

Mechanics

$$\frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx} \left(\frac{1}{2} v^2 \right)$$

$$x = a \cos(nt + \alpha) + c$$

$$x = a \sin(nt + \alpha) + c$$

$$\ddot{x} = -n^2(x - c)$$

Student Name _____

Teacher _____

Select the letter A, B, C or D that best answers the question. Fill in the response oval completely.

Sample $3 + 5 =$ (A) 7 (B) 8 (C) 9 (D) 10

A B C D

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

A B C D

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word '*correct*' and drawing an arrow as follows:

A  B  C  D 

Question

1. **A**  **B**  **C**  **D** 
2. **A**  **B**  **C**  **D** 
3. **A**  **B**  **C**  **D** 

Question	Marks
MC	
4	
5	
6	
Total	30

STUDENT NAME: _____

TEACHER: _____

Q 4

SECTION II

27 marks

Attempt Questions 4 – 6

Allow about 37 minutes for this section.

In Questions 4 – 6, your responses should include relevant mathematical reasoning and/or calculations.

- **Write your name at the top of each booklet.** Answer the questions in the space provided. These spaces provide guidance for the expected length of response.
- Extra writing space is provided at the back of each question booklet. If you use this space, clearly indicate which part of the question you are answering.

Question 4 (8 marks)

(a) Rationalise the denominator $\frac{3}{2-\sqrt{2}}$.

2

(b) Factorise completely $4x^2 - 12xy + 9y^2 - 49z^2$

2

Question 4 continues on page 10

(c) Solve by completing the square $2x^2 + 7x + 6 = 0$

3

(d) Given that $9^p \times 27^q = 3^n$. Express n in terms of p and q .

1

End of Question 4

Extra writing space.

If you use this space, clearly indicate which Question/part you are answering.

[illegible]

Extra writing space.

If you use this space, clearly indicate which Question/part you are answering.

[illegible]

STUDENT NAME: _____

Q 5


TEACHER: _____

- **Write your name at the top of each booklet.** Answer the questions in the space provided. These spaces provide guidance for the expected length of response.
- Extra writing space is provided at the back of each question booklet. If you use this space, clearly indicate which part of the question you are answering.

Question 5 (5 marks)

- (a) A radio station tower was built in two sections. From a point 87 m from the base of the tower, the angle of elevation of the top of the first section is 25° , and the angle of elevation of the top of the second section is 40° . To the nearest metre, what is the height of the top section of the tower?

3



Question 5 continues on page 14

(b) What is the **exact** value of $\operatorname{cosec}60^\circ - \cot 60^\circ$?

2

End of Question 5

Extra writing space.

If you use this space, clearly indicate which Question/part you are answering.

[illegible]

Extra writing space.

If you use this space, clearly indicate which Question/part you are answering.

This image shows a full page of primary-ruled paper. It features multiple sets of horizontal lines designed to guide handwriting. Each set consists of three lines: two dashed outer lines and one solid middle line. These sets are repeated down the entire page, providing ample space for practicing letter formation and alignment. The paper is otherwise blank, with no margins or additional markings.

STUDENT NAME: _____

TEACHER: _____

Q 6

- **Write your name at the top of each booklet.** Answer the questions in the space provided. These spaces provide guidance for the expected length of response.
- Extra writing space is provided at the back of each question booklet. If you use this space, clearly indicate which part of the question you are answering.

Question 6 (14 marks)

- (a) Express the quadratic function $y = x^2 - 8x + 10$ in the form $y = (x - a)^2 + b$. **2**


- (b) Neatly sketch the parabola from part (a) in the space below, clearly showing the coordinates of the vertex and y intercept. **2**

Question 6 continues on page 18

(c) Each of the diagrams below show an incomplete graph of a function. Complete each graph if the function is:


(i) odd

1



(ii) Even

1




Question 6 continues on page 19

(d) Draw a neat sketch of each graph in the space provided, stating it's domain and range.

(i) $x^2 + y^2 = 4$

2




Domain:

Range:

(ii) $y = -\frac{2}{x}$

2



Domain:

Range:

Question 6 continues on page 20

- (iii) With reference to the graphs in (i) and (ii) above, explain why the graph in (i) represents a relation and the graph in (ii) represents a function.

2

- (e) Solve the equation $|2x - 8| = 6$.

2

End of Question 6

End of Paper

Extra writing space.

If you use this space, clearly indicate which Question/part you are answering.

[illegible]

Extra writing space.

If you use this space, clearly indicate which Question/part you are answering.

[illegible]

Outcomes Addressed in this Question

MA11-1

uses algebraic and graphical techniques to solve, and where appropriate, compare alternative solutions to problems

Solutions	Marking Guidelines
a) $\frac{3}{2-\sqrt{2}}$ $= \frac{3}{2-\sqrt{2}} \times \frac{2+\sqrt{2}}{2+\sqrt{2}}$ $= \frac{3(2+\sqrt{2})}{4-2}$ $= \frac{3(2+\sqrt{2})}{2}$ b) $4x^2 - 12xy + 9y^2 - 49z^2$ $= (4x^2 - 12xy + 9y^2) - 49z^2$ $= (2x - 3y)^2 - 49z^2$ $= (2x - 3y)^2 - (7z)^2$ $= (2x - 3y - 7z)(2x - 3y + 7z)$ c) $2x^2 + 7x + 6 = 0$ $x^2 + \frac{7}{2}x = -3$ $x^2 + \frac{7}{2}x + \left(\frac{7}{4}\right)^2 = -3 + \left(\frac{7}{4}\right)^2$ $\left(x + \frac{7}{4}\right)^2 = \frac{1}{16}$ $x + \frac{7}{4} = \pm \sqrt{\frac{1}{16}}$ $x + \frac{7}{4} = \pm \frac{1}{4}$ $x = -\frac{7}{4} \pm \frac{1}{4}$ $x = -\frac{3}{2} \text{ or } x = -2$	Award 2 marks for the correct solution. Award 1 mark for substantial progress towards the solution Award 2 marks for the correct solution. Award 1 mark for substantial progress towards the solution Award 3 marks for the correct solution. Award 2 mark for substantial progress towards the correct solution. Award 1 mark for some progress towards the correct solution.

d) $9^p \times 27^q = 3^n$ $(3^2)^p \times (3^3)^q = 3^n$ $3^{2p} \times 3^{3q} = 3^n$ $3^{2p+3q} = 3^n$ $\therefore n = 2p+3q$	Award 1 mark for the correct solution.
--	---

Multiple Choice Answers

1	A
2	C
3	C

Outcomes Addressed in this Question


MA11-3

uses the concepts and techniques of trigonometry in the solution of equations and problems involving geometric shapes

Solutions

Marking Guidelines

a)



Award 3 marks for the correct solution.

Award 2 mark for substantial progress towards the correct solution.

Award 1 mark for some progress towards the correct solution.

$$\frac{y}{87} = \tan 25^\circ$$

$$y = 87 \tan 25^\circ$$

$$\frac{x+y}{87} = \tan 40^\circ$$

$$x + y = 87 \tan 40^\circ$$

$$x = 87 \tan 40^\circ - y$$

$$x = 87 \tan 40^\circ - 87 \tan 25^\circ$$

$$x = 32.43290165$$

$$x = 32m \text{ (nearest metre)}$$

b)

$$\operatorname{cosec} 60^\circ - \cot 60^\circ$$




$$= \frac{1}{\sin 60^\circ} - \frac{1}{\tan 60^\circ}$$



$$= \frac{1}{\frac{\sqrt{3}}{2}} - \frac{1}{\sqrt{3}}$$

$$= \frac{2}{\sqrt{3}} - \frac{1}{\sqrt{3}} = \frac{1}{\sqrt{3}} \text{ or } \frac{\sqrt{3}}{3}$$

Award 2 marks for the correct solution.

Award 1 mark for substantial progress towards the solution

Year 11 Mathematics Half Yearly Assessment (Task 1) 2021		
Question No. 9		Solutions and Marking Guidelines
Outcomes Addressed in this Question		
MA11-2 uses the concepts of functions and relations to model, analyse and solve practical problems		
Outcome	Solutions	Marking Guidelines
MA11-2	(a) $y = x^2 - 8x + 10$ $= (x^2 - 8x + 16) - 16 + 10$ $= (x - 4)^2 - 6$	2 marks Correct solution. 1 mark Attempts to complete the square or equivalent merit.
MA11-2	(b) 	2 marks Correct solution indicating y-intercept and vertex. 1 mark Correct graph only OR incorrect graph with correct vertex and y intercept.
MA11-2	(c) (i) 	1 mark Correct solution.
MA11-2	(c) (i) 	1 mark Correct solution.

MA11-2	(d) (i)  $-2 \leq x \leq 2$ $-2 \leq y \leq 2$ Domain: <i>OR</i> Range: <i>OR</i> $x : x \in [-2, 2]$ $y : y \in [-2, 2]$	2 marks Correct graph AND correct domain and range. 1 mark Incorrect graph but correct domain and range for drawn graph OR correct graph but incorrect domain and range.
MA11-2	(ii)  $x : x \neq 0$ $y : y \neq 0$ Domain: <i>OR</i> Range: <i>OR</i> $(-\infty, 0) \cup (0, \infty)$ $(-\infty, 0) \cup (0, \infty)$	2 marks Correct graph AND correct domain and range. 1 mark Incorrect graph but correct domain and range for drawn graph OR correct graph but incorrect domain and range.
MA11-2	(iii) Graph (i) is a relation because: * it fails the vertical line test * it is one-many * it is many-many Graph (ii) is a function because: * it satisfies the requirements of the vertical line test * it is one -one	2 marks Correct explanation for both. 1 mark Correct explanation for only one.
MA11-2	(f) $2x - 8 = 6$ $2x - 8 = 6$ $2x = 14$ $x = 7$ OR $2x - 8 = -6$ $2x = 2$ $x = 1$	2 marks Correct solution (both equations). 1 mark Only one correct solution OR attempts to apply correct process but makes error.