


- 1 The diagram shows the volume of water displaced when an insoluble solid is placed in a measuring cylinder containing water. The mass of the solid is 5.70 g. ► Bottom of cylinder is truncated.



What is the density of the solid?

- (A) 0.308 g mL⁻¹
(B) 1.90 g mL⁻¹
(C) 2.10 g mL⁻¹
(D) 28.0 g mL⁻¹
- 2 What does 'm' represent in the equation... $\Delta H = -m C \Delta T$?
- (A) molar mass of the reactant chemical
(B) molarity of the final solution
(C) mass of the substance being heated or cooled
(D) moles of the reactant chemical
- 3 A solution is prepared by dissolving 1.50 g of Ca(OH)₂ in enough water to produce 1.00 L of solution. What is the concentration of the hydroxide ions in the solution?
- (A) 3.00 g L⁻¹
(B) 2.00×10^{-2} mol L⁻¹
(C) 4.00×10^{-2} mol L⁻¹
(D) 1.50×10^{-2} g L⁻¹





4 Which of the following is both an empirical formula and a molecular formula?

- (A) CO
- (B) C₃H₆
- (C) CuSO₄ · 5H₂O
- (D) NaCl

5 Which of the statements is a consequence of Avogadro's Law?

- (A) The volume of one mole of CO and one mole of He are both 24.79 L at 25°C and 100 kPa.
- (B) One mole of a compound contains 6.022×10^{23} molecules of that compound.
- (C) Mass divided by the molar mass gives the number of moles.
- (D) Gases expand when heated.

6 Which diagram shows the bonding between a water molecule and an ammonia molecule?







7 When one mole of sodium hydroxide is completely dissolved in water 41.6 kJ of heat energy is released. A chemistry class performed a simple calorimetry experiment to determine this value and *all* of the students got results *lower* than the accepted value.


Which statement best explains the reason for the lower values obtained?

- (A) The thermometers used were inaccurate.
- (B) The mass of water in the cup was not accurately measured.
- (C) The students didn't wait long enough for the final temperature to stabilise.
- (D) Not all the heat produced by the dissolution was absorbed by the calorimeter.

- 8 100 mL of salt water was heated in a flask from 20°C to boiling. Which graph best represents the heating curve?



Questions 9 and 10 refer to the energy level diagram for the reaction...



9 Which statement is *incorrect*?

- (A) The activation energy per mole of hydrogen fluoride formed is 10 kJ mol⁻¹.
- (B) The activation energy of the reverse reaction is 74 kJ mol⁻¹.
- (C) The heat of reaction per mole of hydrogen fluoride is represented by the value of X.
- (D) The heat of reaction is 64 kJ mol⁻¹ of fluorine reacted.

10 What effect would adding a catalyst have on the reaction?

- (A) X would stay the same and Y would become smaller.
- (B) X would become smaller and Y would become larger.
- (C) Both X and Y would become smaller.
- (D) Z would become larger.

11 Which physical property of water can be used to explain its relatively high values for surface tension and boiling point?

- (A) strong intermolecular forces
- (B) low molecular mass
- (C) low viscosity
- (D) high solvency

- 12 Hydrogen chloride has a very high solubility in water = 70 g per 100 g H₂O at 20°C.

Which intermolecular force best explains this high aqueous solubility?

- (A) dispersion force
- (B) dipole – dipole force
- (C) hydrogen bonding
- (D) all of the above

- 13 To which of these reactions can Guy–Lussac’s law be applied?

- (A) $\text{Fe}_{(s)} + \text{S}_{(s)} \rightarrow \text{FeS}_{(s)}$
- (B) $2\text{H}_{2(g)} + \text{O}_{2(g)} \rightarrow 2\text{H}_2\text{O}_{(g)}$
- (C) $\text{Ba}^{2+}_{(aq)} + \text{SO}_4^{2-}_{(aq)} \rightarrow \text{BaSO}_{4(s)}$
- (D) $\text{HCl}_{(aq)} + \text{NaOH}_{(aq)} \rightarrow \text{NaCl}_{(aq)} + \text{H}_2\text{O}_{(l)}$

- 14 One mole samples of metals are completely reacted in oxygen gas to form metal oxides. The table shows the formulae results...

METAL	METAL OXIDE FORMED
Na	Na ₂ O
Mg	MgO
Al	Al ₂ O ₃
Ni	NiO ₂

Which metal requires the most moles of oxygen to form an oxide?

- (A) Na
- (B) Mg
- (C) Al
- (D) Ni

- 15 Which of these solute types is insoluble in water?

- (A) non–polar covalent
- (B) covalent network
- (C) large molecular
- (D) all of the above

Part B – 48 marks**Attempt Questions 16 – 27****Allow about 90 minutes for this part****► Show all relevant working in questions involving calculations.**

Question 16 (2 marks)

Construct Lewis electron dot structures for water and hydrogen sulfide...

WATER	HYDROGEN SULFIDE

Question 17 (3 marks)

Identify whether the compound is soluble (s) or insoluble (i) in water...

Compound	NaCl	sucrose	NaOH	HCl	cellulose	SiO ₂
s or i ?						

Question 18 (4 marks)

- (a) Identify the physical property of water which explains its ability to moderate the temperature of aquatic environments. **(1 mark)**


- (b) Thermal pollution can cause serious environmental problems in waterways.

- (i) Identify a source of thermal pollution. **(1 mark)**

- (ii) Discuss the implications for life if a body of water is affected by thermal pollution. **(2 marks)**

Question 19 (2 marks)

The diagrams show the crystal structures of diamond and ice...



Describe how these crystals are held together.

Question 20 (6 marks)

When nickel(II) chloride (NiCl_2) solution is added to sodium hydroxide solution, a green precipitate forms.

- (a) Write the net ionic equation for the reaction. **(1 mark)**

- (b) In one reaction, 60 mL of 0.20 mol L^{-1} sodium hydroxide is added to 40 mL of 0.50 mol L^{-1} solution of NiCl_2 .

- (i) Calculate the mass of precipitate formed. **(3 marks)**

Question 20 (continued)

- (ii) Calculate the concentration of chloride ions in the final solution. **(2 marks)**

Question 21 (6 marks)

Weighed samples of aluminium and magnesium were reacted with excess dilute hydrochloric acid solution, each *metal–acid* reaction producing hydrogen gas. The table shows the masses of metal reacted and the resultant volume of hydrogen gas produced at 25°C and 100 kPa.

METAL	MASS OF METAL (g)	VOLUME OF H ₂ (mL)
Al	0.108	150
Mg	0.0973	100

- (a) From the data in the table above, calculate these quantities... **(2 marks)**

METAL	MOLES OF METAL REACTED	MOLES OF H ₂ PRODUCED
Al		
Mg		

- (b) What is the mole ratio between each of the metal reacted and the hydrogen gas produced? **(1 mark)**

Al : H ₂	Mg : H ₂
:	:

- (c) Write the balanced formulae equations for the reactions of each metal with dilute HCl. **(2 marks)**

Question 21 (continued)

(d) Do the equations in (c) support the results of the experiment? Explain. **(1 mark)**

Question 22 (2 marks)

15.9 grams of potassium dichromate ($\text{K}_2\text{Cr}_2\text{O}_7$) are dissolved in a sufficient amount of water to make up exactly 250 mL of solution.

What is the concentration of the solution? **(2 marks)**

Question 23 (1 mark)

How many hydrogen atoms are present in 25.6 g of urea, $(\text{NH}_2)_2\text{CO}$?

Question 24 (6 marks)

Nikita performs a simple calorimetry experiment to determine the molar $\Delta H_{\text{solution}}$ for KOH. The table shows her experimental data ...

Mass of water in calorimeter	150 g
Mass of KOH dissolved	5.6 g
Initial temperature of water	18°C
Final temperature of water	27°C

- (a) Calculate the molar $\Delta H_{\text{solution}}$ for KOH using Nikita's data. **(3 marks)**

- (b) KOH is also soluble in ethanol. Nikita plans to repeat the experiment using ethanol as the solvent instead of water and she thinks about the changes to the equation... $\Delta H = -m C \Delta T$

- (i) Compare (relatively) the specific heat capacity (C) of ethanol with the value of water. **(1 mark)**

- (ii) Considering your answer in (i), predict the effect on the ΔT , *i.e. will it decrease, increase or remain the same compared to the ΔT of water?* **(1 mark)**

- (c) Identify a salt where $\Delta H_{\text{solution}}$ is endothermic. **(1 mark)**

Question 25 (5 marks)

- (a) Kiely conducts a series of investigations on factors which effect reaction rate. In trial 1, she reacts a one centimetre cube (1 cm^3) of magnesium weighing 1.74 g with 200 mL of $3\text{ mol L}^{-1}\text{ HCl}$ in a beaker. In trial 2, she reacts eight $\frac{1}{2}$ centimetre cubes (total weight = 1.74 g) simultaneously in a beaker.



TRIAL 1



TRIAL 2

Identify which trial has the fastest rate and identify the factor which causes the effect. **(1 mark)**

- (b) Kiely broadens her investigation and explores the effects of temperature and concentration on reaction rate.

- (i) What effect would you predict for temperature variations and how is this effect explained on a particle level? **(2 marks)**

- (ii) What effect would you predict for concentration variations and how is this effect explained on a particle level? **(2 marks)**


Question 26 (4 marks)

Propanone ($\text{C}_3\text{H}_6\text{O}$) is a flammable liquid used extensively as a solvent.

- (a) Write a balanced chemical equation showing the complete combustion of propanone. **(1 mark)**
-
- (b) Write a balanced chemical equation showing the incomplete combustion of propanone. **(1 mark)**
-
- (c) Identify a pollutant produced in equation (b) and explain how its production can be avoided. **(2 marks)**
-
-

Question 27 (7 marks)

- (a) Complete the energy level diagram by drawing in a curve for an endothermic reaction... **(1 mark)**







- (b) In the diagram, draw a dashed (---) line showing the effect of employing a catalyst. **(1 mark)**

Question 27 (continued)


- (c) Explain how a catalyst speeds up reaction rate. **(1 mark)**

- (d) Identify a catalyst used in industry. **(1 mark)**

- (e) Study the four energy level diagrams (I – IV) ...



- (i) Which diagram best represents the most easily ignitable fuel? _____ **(1 mark)**
- (ii) Which diagram best represents a fuel with the highest ignition temperature? _____ **(1 mark)**
- (f) Complete the graph by drawing in a line or curve showing the relative relationship between the kinetic energy of particles and the temperature as they are heated up... **(1 mark)**



HIGHER SCHOOL CERTIFICATE EXAMINATION
Chemistry

DATA SHEET

Avogadro constant, N_A	$6.022 \times 10^{23} \text{ mol}^{-1}$
Volume of 1 mole ideal gas: at 100 kPa and	
at 0°C (273.15 K)	22.71 L
at 25°C (298.15 K)	24.79 L
Ionisation constant for water at 25°C (298.15 K), K_w	1.0×10^{-14}
Specific heat capacity of water	$4.18 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$

Some useful formulae

$$\text{pH} = -\log_{10}[\text{H}^+] \qquad \Delta H = -m C \Delta T$$

Some standard potentials

$\text{K}^+ + \text{e}^-$	$\rightleftharpoons$	K(s)	-2.94 V
$\text{Ba}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ba(s)	-2.91 V
$\text{Ca}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ca(s)	-2.87 V
$\text{Na}^+ + \text{e}^-$	$\rightleftharpoons$	Na(s)	-2.71 V
$\text{Mg}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Mg(s)	-2.36 V
$\text{Al}^{3+} + 3\text{e}^-$	$\rightleftharpoons$	Al(s)	-1.68 V
$\text{Mn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Mn(s)	-1.18 V
$\text{H}_2\text{O} + \text{e}^-$	$\rightleftharpoons$	$\frac{1}{2}\text{H}_2(\text{g}) + \text{OH}^-$	-0.83 V
$\text{Zn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Zn(s)	-0.76 V
$\text{Fe}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Fe(s)	-0.44 V
$\text{Ni}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ni(s)	-0.24 V
$\text{Sn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Sn(s)	-0.14 V
$\text{Pb}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Pb(s)	-0.13 V
$\text{H}^+ + \text{e}^-$	$\rightleftharpoons$	$\frac{1}{2}\text{H}_2(\text{g})$	0.00 V
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	$\text{SO}_2(\text{aq}) + 2\text{H}_2\text{O}$	0.16 V
$\text{Cu}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Cu(s)	0.34 V
$\frac{1}{2}\text{O}_2(\text{g}) + \text{H}_2\text{O} + 2\text{e}^-$	$\rightleftharpoons$	2OH^-	0.40 V
$\text{Cu}^+ + \text{e}^-$	$\rightleftharpoons$	Cu(s)	0.52 V
$\frac{1}{2}\text{I}_2(\text{s}) + \text{e}^-$	$\rightleftharpoons$	I^-	0.54 V
$\frac{1}{2}\text{I}_2(\text{aq}) + \text{e}^-$	$\rightleftharpoons$	I^-	0.62 V
$\text{Fe}^{3+} + \text{e}^-$	$\rightleftharpoons$	Fe^{2+}	0.77 V
$\text{Ag}^+ + \text{e}^-$	$\rightleftharpoons$	Ag(s)	0.80 V
$\frac{1}{2}\text{Br}_2(\text{l}) + \text{e}^-$	$\rightleftharpoons$	Br^-	1.08 V
$\frac{1}{2}\text{Br}_2(\text{aq}) + \text{e}^-$	$\rightleftharpoons$	Br^-	1.10 V
$\frac{1}{2}\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	H_2O	1.23 V
$\frac{1}{2}\text{Cl}_2(\text{g}) + \text{e}^-$	$\rightleftharpoons$	Cl^-	1.36 V
$\frac{1}{2}\text{Cr}_2\text{O}_7^{2-} + 7\text{H}^+ + 3\text{e}^-$	$\rightleftharpoons$	$\text{Cr}^{3+} + \frac{7}{2}\text{H}_2\text{O}$	1.36 V
$\frac{1}{2}\text{Cl}_2(\text{aq}) + \text{e}^-$	$\rightleftharpoons$	Cl^-	1.40 V
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^-$	$\rightleftharpoons$	$\text{Mn}^{2+} + 4\text{H}_2\text{O}$	1.51 V
$\frac{1}{2}\text{F}_2(\text{g}) + \text{e}^-$	$\rightleftharpoons$	F^-	2.89 V

Aylward and Findlay, *SI Chemical Data* (5th Edition) is the principal source of data for this examination paper. Some data may have been modified for examination purposes.

KEY										2		
1		4		79		Symbol of element		5		10		
H 1.008 Hydrogen		Li 6.941 Lithium		Au 197.0 Gold		Name of element		B 10.81 Boron		He 4.003 Helium		
Atomic Number		Atomic Weight		Atomic Weight		Atomic Weight		Atomic Weight		Atomic Weight		
3	Be	4	Be	79	Au	197.0	Gold	5	B	10.81	He	4.003
6.941	9.012	9.012	9.012	197.0	197.0	197.0	197.0	10.81	12.01	14.01	16.00	19.00
11	Na	12	Mg	24.31	24.31	24.31	24.31	13	Al	26.98	28.09	30.97
22.99	22.99	24.31	24.31	24.31	24.31	24.31	24.31	14	Si	28.09	30.97	32.07
Sodium	Magnesium	Sodium	Magnesium	Sodium	Magnesium	Sodium	Magnesium	Aluminum	Silicon	Phosphorus	Sulfur	Chlorine
19	K	20	Ca	40.08	40.08	40.08	40.08	31	Ga	69.72	72.61	74.92
39.10	40.08	40.08	40.08	40.08	40.08	40.08	40.08	32	Ge	72.61	74.92	78.96
Potassium	Calcium	Potassium	Calcium	Potassium	Calcium	Potassium	Calcium	Cerium	Germanium	Arsenic	Selenium	Bromine
37	Rb	38	Sr	87.62	87.62	87.62	87.62	49	In	114.8	118.7	126.9
85.47	87.62	87.62	87.62	87.62	87.62	87.62	87.62	50	Sn	118.7	126.9	131.3
Rubidium	Strontium	Rubidium	Strontium	Rubidium	Strontium	Rubidium	Strontium	Indium	Antimony	Tellurium	Iodine	Xenon
55	Cs	56	Ba	137.3	137.3	137.3	137.3	81	Tl	204.4	207.2	209.0
132.9	137.3	137.3	137.3	137.3	137.3	137.3	137.3	82	Pb	207.2	209.0	210.0
Cesium	Barium	Cesium	Barium	Cesium	Barium	Cesium	Barium	Lead	Bismuth	Polonium	Astatine	Radon
87	Fr	88	Ra	226.0	226.0	226.0	226.0	113	Uuq	114	115	116
[223.0]	[226.0]	[226.0]	[226.0]	[226.0]	[226.0]	[226.0]	[226.0]	114	Uuq	115	116	117
Francium	Radium	Francium	Radium	Francium	Radium	Francium	Radium	Ununquadium	Ununhexium	Ununseptium	Ununoctium	Ununnilium

57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
138.9	140.1	140.9	144.2	[146.9]	150.4	152.0	157.3	158.9	162.5	164.9	167.3	168.9	173.0	175.0
Lanthanum	Cerium	Praseodymium	Neodymium	Promethium	Samarium	Europium	Gadolinium	Terbium	Dysprosium	Holmium	Erbium	Thulium	Ytterbium	Lutetium

89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr
[227.0]	232.0	231.0	238.0	[237.0]	[239.1]	[241.1]	[244.1]	[249.1]	[252.1]	[252.1]	[257.1]	[258.1]	[259.1]	[262.1]
Actinium	Thorium	Protactinium	Uranium	Neptunium	Plutonium	Americium	Curium	Berkelium	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Lawrencium

JRAHS Year 11 Chemistry Final Exam – 2004

Multiple Choice Answers

Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Q13	Q14	Q15
B	C	C	A	A	C	D	C	D	A	A	B	B	D	D

Part B – 48 marks

Attempt Questions 16 – 27

Allow about 90 minutes for this part

► Show all relevant working in questions involving calculations.

Question 16 (2 marks)

Construct Lewis electron dot structures for water and hydrogen sulfide...

WATER	HYDROGEN SULFIDE
$\begin{array}{c} \text{H} : \ddot{\text{O}} : \text{H} \\ \text{ } \end{array}$	$\begin{array}{c} \text{H} : \ddot{\text{S}} : \text{H} \\ \text{ } \end{array}$

Question 17 (3 marks)

Identify whether the compound is soluble (s) or insoluble (i) in water...

Compound	NaCl	sucrose	NaOH	HCl	cellulose	SiO ₂
s or i ?	s	s	s	s	i	i

Question 18 (4 marks)

- (a) Identify the physical property of water which explains its ability to moderate the temperature of aquatic environments. **(1 mark)**

specific heat capacity

- (b) Thermal pollution can cause serious environmental problems in waterways.

- (i) Identify a source of thermal pollution. **(1 mark)**


power generating stations or industrial plants

- (ii) Discuss the implications for life if a body of water is affected by thermal pollution. **(2 marks)**

Thermal pollution causes heat stress in aquatic organisms. Heat stress has diverse consequences including increased metabolic rate, increased heart rate and decreased ability of haemoglobin (and other blood pigments, e.g. hemovanadin) to carry oxygen. Thermal pollution also decreases the amount of dissolved oxygen in water and some species of fish (e.g. trout) will not breed if the water is too warm. ► Any one point mentioned (1 mark) and elucidated (1 mark).

Question 19 (2 marks)

The diagrams show the crystal structures of diamond and ice...



Describe how these crystals are held together.

Diamond is held together by covalent bonds. (1 mark)

Ice is held together by hydrogen bonding. (1 mark)

Question 20 (6 marks)

When nickel(II) chloride (NiCl_2) solution is added to sodium hydroxide solution, a green precipitate forms.

(a) Write the net ionic equation for the reaction.



(b) In one reaction, 60 mL of 0.20 mol L^{-1} sodium hydroxide is added to 40 mL of 0.50 mol L^{-1} solution of NiCl_2 .

(i) Calculate the mass of precipitate formed. **(3 marks)**

$$\text{moles NiCl}_2 = c \times V = 0.50 \text{ mol L}^{-1} \times 0.040 \text{ L} = 0.020 \text{ mol}$$

$$\text{moles NaOH} = c \times V = 0.20 \text{ mol L}^{-1} \times 0.060 \text{ L} = 0.012 \text{ mol} \quad \therefore \text{NaOH is the limiting reactant}$$

$$\text{mol Ni}(\text{OH})_2 = \frac{1}{2} \text{ mol NaOH} = \frac{1}{2} \times 0.012 \text{ mol} = 0.0060 \text{ mol}$$

$$\text{mass Ni}(\text{OH})_2 = n \times M = 0.0060 \text{ mol} \times 92.706 \text{ g mol}^{-1} = 0.5562 = 0.56 \text{ g}$$

► **Correct answer with relevant working. (3 marks)**

► **Incorrect answer with two of correct moles of NiCl_2 and NaOH (2 marks)**
Correct mass from incorrect mol OH^{-}

► **Incorrect answer with correct mol NiCl_2 and NaOH (1 mark)**
or correct mass from incorrect mol OH^{-}
or correct answer with no working

Question 20 (continued)

- (ii) Calculate the concentration of chloride ions in the final solution.
- (2 marks)**

$$\text{mol Cl}^- = 2 \times \text{mol NiCl}_2 = 2 \times 0.020 \text{ mol} = 0.040 \text{ mol} \quad (1 \text{ mark})$$

$$[\text{Cl}^-] = n \div V = 0.040 \text{ mol} \div 0.10 \text{ L} = 0.40 \text{ mol L}^{-1} \quad (1 \text{ mark})$$

Question 21 (6 marks)

Weighed samples of aluminium and magnesium were reacted with excess dilute hydrochloric acid solution, each *metal–acid* reaction producing hydrogen gas. The table shows the masses of metal reacted and the resultant volume of hydrogen gas produced at 25°C and 100 kPa.

METAL	MASS OF METAL (g)	VOLUME OF H ₂ (mL)
Al	0.108	150
Mg	0.0973	100



- (a) From the data in the table above, calculate these quantities...
- (2 marks)**

METAL	MOLES OF METAL REACTED	MOLES OF H ₂ PRODUCED
Al	$n = m \div M$ $n = 0.108 \text{ g} \div 26.98 \text{ g mol}^{-1}$ $n = 0.00400 \text{ mol}$	$n = V \div 24.79 \text{ L mol}^{-1}$ $n = 0.150 \text{ L} \div 24.79 \text{ L mol}^{-1}$ $n = 0.00605 \text{ mol}$
Mg	$n = m \div M$ $n = 0.0973 \text{ g} \div 24.31 \text{ g mol}^{-1}$ $n = 0.00400 \text{ mol}$	$n = V \div 24.79 \text{ L mol}^{-1}$ $n = 0.100 \text{ L} \div 24.79 \text{ L mol}^{-1}$ $n = 0.00403 \text{ mol}$

- (b) What is the mole ratio between each of the metal reacted and the hydrogen gas produced?
- (1 mark)**

Al : H ₂	Mg : H ₂
1:1.51 or 2:3	1:1

- (c) Write the balanced formulae equations for the reactions of each metal with dilute HCl.
- (2 marks)**



Question 21 (continued)

- (d) Do the equations in (c) support the results of the experiment? Explain. **(1 mark)**

Yes, the results of the experiment are in concordance with the results shown by the metal to hydrogen gas ratio as expressed in the balanced equations.

Question 22 (2 marks)

15.9 grams of potassium dichromate ($\text{K}_2\text{Cr}_2\text{O}_7$) are dissolved in a sufficient amount of water to make up exactly 250 mL of solution.

What is the concentration of the solution? **(2 marks)**

$$\text{moles K}_2\text{Cr}_2\text{O}_7 = m \div M = 15.9 \text{ g} \div 294.2 \text{ g mol}^{-1} = 0.0540 \text{ mol} \quad (1 \text{ mark})$$

$$[\text{K}_2\text{Cr}_2\text{O}_7] = n \div V = 0.0540 \text{ mol} \div 0.250 \text{ L} = 0.216 \text{ mol L}^{-1} \quad (1 \text{ mark})$$

Question 23 (1 mark)

How many hydrogen atoms are present in 25.6 g of urea, $(\text{NH}_2)_2\text{CO}$?

$$\text{moles urea} = m \div M = 25.6 \text{ g} \div 60.062 \text{ g mol}^{-1} = 0.426 \text{ mol}$$

$$\text{number of hydrogen atoms} = 0.426 \text{ moles urea} \times 4 \times 6.022 \times 10^{23} \text{ atoms mol}^{-1}$$

$$\text{number of hydrogen atoms} = 1.02 \times 10^{24} \text{ atoms} \quad (1 \text{ mark})$$

Question 24 (6 marks)

Nikita performs a simple calorimetry experiment to determine the molar $\Delta H_{\text{solution}}$ for KOH. The table shows her experimental data ...

Mass of water in calorimeter	150 g
Mass of KOH dissolved	5.6 g
Initial temperature of water	18°C
Final temperature of water	27°C

- (a) Calculate the molar $\Delta H_{\text{solution}}$ for KOH using Nikita's data. **(3 marks)**

$$\Delta H = -m C \Delta T = -150 \text{ g} \times 4.18 \text{ J g}^{-1} \text{ }^{\circ}\text{C} \times (27 - 18^{\circ}\text{C}) = 5643 \text{ J} = 5.643 \text{ kJ} \quad (1 \text{ mark})$$

$$\Delta H \text{ g}^{-1} = \Delta H \div g_{\text{KOH}} = 5.643 \text{ kJ} \div 5.6 \text{ g} = 1.008 \text{ kJ g}^{-1} \quad (1 \text{ mark})$$

$$\text{molar } \Delta H = \Delta H \text{ g}^{-1} \times M_{\text{KOH}} = 1.008 \text{ kJ g}^{-1} \times 56.108 \text{ g mol}^{-1} = 56.54 = 60 \text{ kJ mol}^{-1} \quad (1 \text{ mark})$$

► *The final answer is expressed in one significant figure due to the ΔT subtraction calculation.*

- (b) KOH is also soluble in ethanol. Nikita plans to repeat the experiment using ethanol as the solvent instead of water and she thinks about the changes to the equation... $\Delta H = -m C \Delta T$

- (i) Compare (relatively) the specific heat capacity (C) of ethanol with the value of water. **(1 mark)**

$$C_{\text{ethanol}} < C_{\text{water}}$$

- (ii) Considering your answer in (i), predict the effect on the ΔT , i.e. will it decrease, increase or remain the same compared to the ΔT of water? **(1 mark)**

Ethanol's lower specific heat capacity would cause the ΔT to be greater than water's.

- (c) Identify a salt where $\Delta H_{\text{solution}}$ is endothermic. **(1 mark)**

AgNO₃ CuSO₄ · 5H₂O KCl KI KNO₃ MgSO₄ · 7H₂O NaCl NaNO₃ Na₂SO₄ · 10H₂O NH₄Cl NH₄NO₃

Question 25 (5 marks)

- (a) Kiely conducts a series of investigations on factors which effect reaction rate. In trial 1, she reacts a one centimetre cube (1 cm^3) of magnesium weighing 1.74 g with 200 mL of $3\text{ mol L}^{-1}\text{ HCl}$ in a beaker. In trial 2, she reacts eight $\frac{1}{2}$ centimetre cubes (total weight = 1.74 g) simultaneously in a beaker.



TRIAL 1



TRIAL 2

Identify which trial has the fastest rate and identify the factor which causes the effect. **(1 mark)**

Trial 2 has the fastest rate because the magnesium cubes have greater surface area.

- (b) Kiely broadens her investigation and explores the effects of temperature and concentration on reaction rate.

- (i) What effect would you predict for temperature variations and how is this effect explained on a particle level? **(2 marks)**

An increase in temperature would increase reaction rate. (1 mark)

This effect is primarily explained by the fact that more reactant molecules will have more energy to react properly, i.e. exceed the activation energy requirement. (1 mark)

► A minor effect is the increase in collision frequency of the reactant molecules... an answer grudgingly accepted!

- (ii) What effect would you predict for concentration variations and how is this effect explained on a particle level? **(2 marks)**

An increase in reactant concentration would increase reaction rate. (1 mark)


This effect is explained by the fact that the collision frequency of the reactant molecules will increase. (1 mark)

► An increase in the concentration of only one of the reactants will cause this effect.


Question 26 (4 marks)

Propanone ($\text{C}_3\text{H}_6\text{O}$) is a flammable liquid used extensively as a solvent.

- (a) Write a balanced chemical equation showing the complete combustion of propanone. (1 mark)



- (b) Write a balanced chemical equation showing the incomplete combustion of propanone. (1 mark)



► *Several other equations are possible.*


- (c) Identify a pollutant produced in equation (b) and explain how its production can be avoided. (2 marks)

C or CO is a pollutant. (1 mark)

Pollution can be avoided by providing sufficient oxygen to the combusting fuel. (1 mark)

Question 27 (7 marks)

- (a) Complete the energy level diagram by drawing in a curve for an endothermic reaction... (1 mark)



- (b) In the diagram, draw a dashed (----) line showing the effect of employing a catalyst. (1 mark)

Question 27 (continued)





- (c) Explain how a catalyst speeds up reaction rate. **(1 mark)**

Catalysts speed up reaction rates tremendously by providing an alternative reaction mechanism (pathway) with a lower activation energy.


- (d) Identify a catalyst used in industry. **(1 mark)**

Nickel (margarine manufacture), platinum (diverse uses), Fe_3O_4 (ammonia manufacture), V_2O_5 (sulfuric acid manufacture). ► Only name or formula is required.

- (e) Study the four energy level diagrams (I – IV) ...



- (i) Which diagram best represents the most easily ignitable fuel? **III (1 mark)**
- (ii) Which diagram best represents a fuel with the highest ignition temperature? **I (1 mark)**
- (f) Complete the graph by drawing in a line or curve showing the relative relationship between the kinetic energy of particles and the temperature as they are heated up... **(1 mark)**



► **Temperature is directly proportional to the average kinetic energy of the particles.**