



NORMANHURST BOYS HIGH SCHOOL

MATHEMATICS ADVANCED

2019 Year 11 Course Assessment Task 2

y-day, xx May 2019

General instructions

- Working time – 1 hour.
(plus 5 minutes reading time)
- Write using blue or black pen. Where diagrams are to be sketched, these may be done in pencil.
- NESA approved calculators may be used.
- Attempt **all** questions.
- At the conclusion of the examination, bundle the booklets used in the correct order within this paper and hand to examination supervisors.

SECTION I

- Mark your answers on the answer grid provided (on page 3)

SECTION II

- Commence each new question on a new booklet. Write on both sides of the paper.
- All necessary working should be shown in every question. Marks may be deducted for illegible or incomplete working.

NESA STUDENT #: # BOOKLETS USED:

Marker's use only.

QUESTION	1-5	1	2	3	??	??	Total	%
MARKS	$\overline{5}$	$\overline{10}$	$\overline{6}$	$\overline{4}$	$\overline{11}$	$\overline{7}$	$\overline{43}$	

Section I

5 marks

Attempt Question 1 to 5

Allow approximately 10 minutes for this section

Mark your answers on the answer grid provided (labelled as page 3).

Questions

Marks

1. What is the solution to the equation $\cos\left(\frac{\theta}{2} + 30^\circ\right) = \sin \theta$ for $0^\circ \leq \theta \leq 90^\circ$? 1

(A) $\theta = 20^\circ$ (B) $\theta = 30^\circ$ (C) $\theta = 40^\circ$ (D) $\theta = 50^\circ$

2. What are the values of a and b if $a + b\sqrt{2} \equiv \frac{2}{3 + \sqrt{2}}$? 1

(A) $a = \frac{6}{7}, b = -\frac{2}{7}$ (C) $a = \frac{6}{5}, b = \frac{2}{5}$

(B) $a = \frac{6}{7}, b = \frac{2}{7}$ (D) $a = \frac{6}{5}, b = -\frac{2}{5}$

3. Determine the type of roots for the following equation:

$$4x^2 - 4x + 1 = 0$$


(A) Two distinct, real roots (C) One pair of unreal roots
(B) Irrational roots (D) One pair of equal roots

4. What is $\frac{\tan \theta \sec \theta}{1 + \tan^2 \theta}$ in simplest form?

(A) $\sin \theta$ (B) $\cos \theta$ (C) $\tan \theta$ (D) $\sin \theta \cos \theta$

5. The equation of the graph below takes the form $y = A \cos Bx + 3$.

1



Which of the following are the values of A and B ?

(A) $A = 3, B = -2$

(C) $A = 6, B = 3$

(B) $A = -3, B = 2$

(D) $A = -6, B = \pi$

Answer grid for Questions 1 - 5

- | | | | | | |
|---|---|-----|-----|-----|-----|
| 1 | — | (A) | (B) | (C) | (D) |
| 2 | — | (A) | (B) | (C) | (D) |
| 3 | — | (A) | (B) | (C) | (D) |
| 4 | — | (A) | (B) | (C) | (D) |
| 5 | — | (A) | (B) | (C) | (D) |

Examination continues overleaf...

Question 1 (8 Marks) Commence a NEW booklet. **Marks**

(a) Fully simplify:

i. $8x - 2y + 4x + 9x + z - 2y.$ **1**

.....

.....

ii. $(2p - 5)^2 - (2p + 5)^2.$ **2**

.....

.....

.....

.....

(b) $\frac{5}{2k+6} + \frac{k}{k^2-9}.$ **3**

.....

.....

.....

.....

.....

(c) Solve for x and y : **2**

$$\begin{cases} 7x - 3y = 41 \\ 3x - y = 17 \end{cases}$$

.....

.....

.....

.....

.....

Question 2 (22 Marks)

Commence a NEW booklet.

Marks

- (a) Is the function $f(x) = x^3 - x$ odd or even? Justify your answer with the appropriate reasoning. **2**

.....

.....

.....

.....

- (b) Find the domain and range of the following, writing in interval notation:

- i. $y = 3x^2 - 1$. **1**

.....

.....

.....

.....

- ii. $y = \frac{1}{x-2}$. **2**

.....

.....

.....

.....

- iii. $y = -\sqrt{9-x^2}$. **2**

.....

.....

.....

.....

- (c) Suppose $f : x \mapsto x^2 + 1$ and $g : x \mapsto 3 - x$.

Find:

- i. $f \circ g(2)$. **1**

.....

- ii. $g \circ f(x)$, in simplest form. **2**

.....

- iii. Find the value(s) of x such that $g \circ f(x) = f(x)$. **2**

.....


- (d) Solve for x : $|7 - 3x| = 1$. **2**

.....


Question 2 continues overleaf...

- (e) Sketch the following graphs on separate number planes. Showing all important features, including x and y intercepts.


i. $y = (x - 1)(x + 3)(4 - x)$.

2


ii. $y = -x^2 - 4x + 5$.

2

iii. $y = \sqrt{x+3}$

2

iv. $y = |x-2| + 1$

2

Question 3 (17 Marks)

Commence a NEW booklet.

Marks

- (a) In $\triangle ABC$, $\angle BAC = 50^\circ$, $AC = 40$ cm and $BC = 25$ cm.

3

Find the size of $\angle ABC$ correct to the nearest minute.

.....

.....

.....

.....

.....

- (b) Solve for θ :

$$4\sin^2\theta - 3 = 0 \quad 0^\circ \leq \theta \leq 360^\circ$$

3

.....

.....

.....

.....

.....

- (c) If $\sin x = -\frac{2}{5}$ and $\cos x > 0$, find the exact value of $\tan x$.

2

.....

.....

.....

.....

- (d) Fully simplify:

3

$$\frac{\cos \theta - 2 \cos^3 \theta}{\sin \theta - 2 \sin^3 \theta}$$

.....


.....

.....

.....

.....

- (e) A man walking along a straight road observes a spire bearing 045°T with an angle of elevation of 5° .



After travelling a distance of 5 km, the spire bears 315°T and the angle of elevation is 8° .

- i. Find the size of $\angle ASB$.

1

.....

.....

- ii. Show that $AS = h \tan 85^\circ$, and write a similar expression for BS .

2

.....

.....

.....

- iii. Find the height of the spire, correct to the nearest metre.

3

.....

.....

.....

.....

.....

.....

.....

End of paper.

2019 Mathematics Advanced (Year 11 Course) Assessment Task 2 STUDENT SELF REFLECTION

1. In hindsight, did I do the best I can? Why or why not?

.....
.....
.....

2. Which topics did I need more help with, and what parts specifically?

- Q1, 2 - Functions, absolute value inequality

.....
.....
.....

- Q3 - Trigonometric Ratios

.....
.....
.....

- Q4 - Trigonometric Ratios (Graphs, identities)

.....
.....
.....

- Q5 - Plane Geometry

.....
.....
.....

- Q6 - Applied Trigonometry

.....
.....
.....

- Q7 - Problem solving

.....
.....
.....

3. What other parts from the teacher feedback can I take away to refine my solutions for future reference?

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....



SOLUTIONS

NORMANHURST BOYS HIGH SCHOOL

MATHEMATICS ADVANCED

2019 Year 11 Course Assessment Task 2

y-day, xx May 2019

General instructions

- Working time – 1 hour.
(plus 5 minutes reading time)
- Write using blue or black pen. Where diagrams are to be sketched, these may be done in pencil.
- NESA approved calculators may be used.
- Attempt all questions.

SECTION I

- Mark your answers on the answer grid provided (on page 3)

SECTION II

- All necessary working should be shown in every question. Marks may be deducted for illegible or incomplete working.


NESA STUDENT #: # BOOKLETS USED:

Marker's use only.

QUESTION	1-5	6	7	8	Total	%
MARKS	$\overline{5}$	$\overline{8}$	$\overline{22}$	$\overline{16}$	$\overline{51}$	

5. The equation of the graph below takes the form $y = A \cos Bx + 3$.

1



Which of the following are the values of A and B ?

(A) $A = 3, B = -2$

(C) $A = 6, B = 3$

(B) $A = -3, B = 2$

(D) $A = -6, B = 180^\circ$

Answer grid for Questions 1 - 5

1 -	☐ (A)	☐ (B)	☒ (C)	☐ (D)
2 -	☒ (A)	☐ (B)	☐ (C)	☐ (D)
3 -	☐ (A)	☐ (B)	☐ (C)	☒ (D)
4 -	☒ (A)	☐ (B)	☐ (C)	☐ (D)
5 -	☐ (A)	☒ (B)	☐ (C)	☐ (D)

Examination continues overleaf...

Section II

46 marks

Attempt Question 6 to 8

Allow approximately 50 minutes for this section

Write your answers in the space provided.

Question 6 (8 Marks)

Marks

(a) Fully simplify:

i. $8x - 2y + 4z + 9x + z - 2y.$

1

$$= 17x - 4y + 5z \quad \checkmark$$

ii. $(2p - 5)^2 - (2p + 5)^2.$

2

$$= [2p - 5 + 2p + 5][2p - 5 - 2p - 5] \quad \checkmark$$

$$= 4p(-10)$$

$$= -40p \quad \checkmark$$

iii. $\frac{5}{2k+6} + \frac{k}{k^2-9}.$

3

$$= \frac{5}{2(k+3)} + \frac{k}{(k+3)(k-3)} \quad \checkmark$$

$$= \frac{5(k-3) + 2k}{2(k+3)(k-3)} \quad \checkmark$$

$$= \frac{7k-15}{2(k+3)(k-3)} \quad \checkmark$$

(b) Solve for x and y :

2

$$\begin{cases} 7x - 3y = 41 & \text{--- (1)} \\ 3x - y = 17 & \text{--- (2)} \end{cases}$$

$$\textcircled{2} \times 3,$$

$$9x - 3y = 51 \text{ --- } \textcircled{3}$$

$$\textcircled{3} - \textcircled{1},$$

$$2x = 10$$

$$x = 5$$

✓

Sub into $\textcircled{2}$,

$$15 - y = 17$$

$$y = -2$$

✓

$$\therefore x = 5, y = -2$$

Question 7 (22 Marks)

Marks

- (a) Is the function $f(x) = x^3 - x$ odd or even? Justify your answer with the appropriate reasoning. 2

$$\begin{aligned}
 f(x) &= x^3 - x && \therefore f(x) \text{ is odd } \checkmark \\
 f(-x) &= (-x)^3 - (-x) \\
 &= -x^3 + x \\
 &= -f(x) \quad \checkmark
 \end{aligned}$$

- (b) Find the domain and range of the following, writing in interval notation:

i. $y = 3x^2 - 1$.

2

$$\begin{aligned}
 D &: \{x : x \in \mathbb{R}\} \quad \checkmark \\
 R &: \{y : y \in [-1, \infty)\} \quad \checkmark
 \end{aligned}$$

ii. $y = \frac{1}{x-2}$.

2

$$\begin{aligned}
 D &: \{x : x \in \mathbb{R} \setminus 2\} \quad \checkmark \\
 R &: \{y : y \in \mathbb{R} \setminus 0\} \quad \checkmark
 \end{aligned}$$

iii. $y = -\sqrt{9-x^2}$.

2

$$\begin{aligned}
 D &: \{x : x \in [-3, 3]\} \quad \checkmark \\
 R &: \{y : y \in [-3, 0]\} \quad \checkmark
 \end{aligned}$$

Q7(b) Deduct 1 mark if answers not written in interval notation.

- (c) Suppose $f: x \mapsto x^2 + 1$ and $g: x \mapsto 3 - x$.

$$f(x) = x^2 + 1$$

$$g(x) = 3 - x$$

Find:

i. $(f \circ g)(2) = f(g(2))$

1

$$= f(1)$$

$$= 2$$

ii. $(g \circ f)(x)$, in simplest form.

$$= g(f(x))$$

$$= g(x^2 + 1)$$

$$= 3 - (x^2 + 1)$$

$$= 2 - x^2$$

iii. Find the value(s) of x such that $(g \circ f)(x) = f(x)$.

2

$$g(f(x)) = f(x)$$

$$2 - x^2 = x^2 + 1$$

$$2x^2 = 1$$

$$x = \pm \frac{1}{\sqrt{2}}$$

(d) Solve for x : $|7 - 3x| = 1$.

2

$$7 - 3x = 1$$

or

$$7 - 3x = -1$$

$$3x = 6$$

$$3x = 8$$

$$x = 2$$


$$x = \frac{8}{3}$$

Question 7 continues overleaf...

- (e) Sketch the following graphs on separate number planes. Showing all important features, including x and y intercepts.


i. $y = (x - 1)(x + 3)(4 - x)$.

shape ✓
Critical point / x / y intercepts labelled ✓ 2




ii. $y = -x^2 - 4x + 5 = (x+5)(1-x)$

2




iii. $y = \sqrt{x+3}$

2



iv. $y = |x-2|+1$

2



Question 8 (16 Marks)

Marks

- (a) In $\triangle ABC$, $\angle BAC = 50^\circ$, $AC = 30$ cm and $BC = 25$ cm.

3

Find the size of $\angle ABC$ correct to the nearest minute.

$$\frac{30}{\sin \angle ABC} = \frac{25}{\sin 50^\circ} \quad \checkmark$$

$$\sin \angle ABC = \frac{30 \sin 50^\circ}{25}$$

$$= 0.91925 \dots$$

$$\angle ABC = 66^\circ 49' \text{ or } 113^\circ 11' \quad \checkmark$$

- (b) Solve for θ :

2

$$4 \sin^2 \theta - 3 = 0 \quad 0^\circ \leq \theta \leq 360^\circ$$

$$\sin^2 \theta = \frac{3}{4}$$


$$\sin \theta = \pm \frac{\sqrt{3}}{2} \quad \checkmark$$

$$\theta = 60^\circ, 120^\circ, 240^\circ, 300^\circ \quad \checkmark$$

- (c) If $\sin x = -\frac{2}{5}$ and $\cos x > 0$, find the exact value of $\tan x$.

2

$$\tan x = -\frac{2}{\sqrt{21}} \quad \checkmark$$




(d) Fully simplify:

3

$$\frac{\cos \theta - 2 \cos^3 \theta}{\sin \theta - 2 \sin^3 \theta}$$

$$\begin{aligned}
 &= \frac{\cos \theta (1 - 2 \cos^2 \theta)}{\sin \theta (1 - 2 \sin^2 \theta)} \checkmark \\
 &= \frac{\cos \theta [1 - 2(1 - \sin^2 \theta)]}{\sin \theta (1 - 2 \sin^2 \theta)} \checkmark \\
 &= \frac{\cos \theta (-1 + 2 \sin^2 \theta)}{1 - 2 \sin^2 \theta} \\
 &= -\cos \theta \checkmark
 \end{aligned}$$

(e) A man walking along a straight road observes a spire bearing 045°T with an angle of elevation of 5° .After travelling a distance of 5km, the spire bears 315°T and the angle of elevation is 8° .i. Find the size of $\angle ASB$.

$$\begin{aligned}
 \angle ASB &= 45^\circ + 45^\circ \\
 &= 90^\circ \checkmark
 \end{aligned}$$

ii. Show that $AS = h \tan 85^\circ$ where h is the height of the spire in metres. Write a similar expression for BS .

$$\text{In } \triangle AST, \angle ATS = 90^\circ - 5^\circ = 85^\circ$$

$$\tan 85^\circ = \frac{AS}{h}$$


$$\therefore AS = h \tan 85^\circ$$

In $\triangle BST$,

$$\begin{aligned}
 \angle STB &= 90^\circ - 8^\circ \\
 &= 82^\circ
 \end{aligned}$$

$$\tan 82^\circ = \frac{BS}{h}$$

$$\therefore BS = h \tan 82^\circ \checkmark$$



- iii. Find the height of the spire, correct to the nearest metre.

3

In $\triangle ABC$

$$AS^2 + BS^2 = AB^2$$

$$h^2 \tan^2 85^\circ + h^2 \tan^2 82^\circ = 5000^2 \quad \checkmark$$

$$h^2 (\tan^2 85^\circ + \tan^2 82^\circ) = 5000^2$$

$$h = \frac{5000}{(\tan^2 85^\circ + \tan^2 82^\circ)^{\frac{1}{2}}} \quad \checkmark$$

$$= 371 \text{ m (nearest m)} \quad \checkmark$$

End of paper.